

INTEGRAL SOLUTIONS OF THE HEPTIC EQUATION WITH FIVE VARIABLES

$$x^3 - y^3 + (x^2 + y^2) + z^3 - w^3 = 2 + 11(x - y)p^6$$

S. Vidhyalakshmi, A. Kavitha, M. A. Gopalan

Department of Mathematics,

Shrimati Indira Gandhi College, Trichy-620002, India

Abstract- The non-homogeneous Diophantine equation of degree seven with five variables represented by $x^3 - y^3 + (x^2 + y^2) + z^3 - w^3 = 2 + 11(x - y)p^6$ is analyzed for its non-zero distinct integer solutions. A few interesting relation between the solutions and special numbers namely Polygonal numbers, Pyramidal numbers, centered Polygonal numbers are exhibited.

Index Terms- Integral solutions, heptic, non-homogeneous equation. M.sc 2000 mathematics subject classification: 11D41

I. INTRODUCTION

Diophantine equations, homogeneous and non-homogeneous have aroused the interest of numerous mathematicians since antiquity as can be seen from [1,4]. The problem of finding all integer solutions of a diophantine equation with three or more variables and degree at least three, in general presents a good deal of difficulties. There is vast general theory of homogeneous quadratic equations with three variables. Cubic equations in two variables fall into the theory of elliptic curves which is a very developed theory but still an important topic of current research. A lot is known about equations in two variables in higher degrees. Very few equations with more than three variables and degree at least three is known. It is worth to note that undesirability appears in equations, even perhaps at degree four with fairly small co-efficient. It seems that

much work has not been done in solving higher order Diophantine equations. In [5-26] a few higher order equations are considered for integral solutions. In this communication a seventh degree non-homogeneous equation, with five variables represented by $x^3 - y^3 + (x^2 + y^2) + z^3 - w^3 = 2 + 11(x - y)p^6$ is considered and a few interesting relations among the solutions are presented.

II NOTATIONS

$t_{m,n}$: Polygonal number of rank n with size m

CP_n^m : Centered Pyramidal number of rank n with size m .

S_n : Star number of rank n

J_n : Jacobsthal number of rank n

j_n : Jacobsthal-Lucas number of rank n

$F_{4,n,6}$ = Four dimensional hexagonal figurate number of rank n .

$F_{4,n,8}$ = Four dimensional octagonal figurate number of rank n .

III METHOD OF ANALYSIS

The non-homogeneous heptic equation with five unknowns to be solved for its distinct non-zero integral solutions is

$$x^3 - y^3 + (x^2 + y^2) + z^3 - w^3 = 2 + 11(x - y)p^6 \quad (1)$$

Introduction of the linear transformations,

$$\begin{aligned} x &= u + 1, & y &= u - 1, & z &= v + 1, \\ w &= v - 1 \end{aligned} \quad (2)$$

in (1) leads to

$$4u^2 + 6v^2 = 22p^6 \quad (3)$$

Different methods of obtaining the patterns of integer solutions to (1) are illustrated below:

Pattern:1

Let $p = 4a^2 + 6b^2 \quad (4)$

Write 22 as

$$22 = (4 + i\sqrt{6})(4 - i\sqrt{6}) \quad (5)$$

Using (4), (5) in (3) and applying the method of factorization define

$$2u + i\sqrt{6}v = (4 + i\sqrt{6})(2\alpha + i2\sqrt{6}\beta) \quad (6)$$

where $(2\alpha + i2\sqrt{6}\beta) = (2a + ib\sqrt{6})$ from which we have

$$\left. \begin{aligned} \alpha &= 2^5 a^6 - 15 \cdot 2^3 \cdot 6a^4 b^2 + 306^2 a^2 b^4 + 3 \cdot 6^2 b^6 \\ \beta &= 6 \cdot 2^4 a^5 b - 480 a^3 b^3 = 6^3 a b^5 \end{aligned} \right\} (7)$$

Equating real and imaginary parts in (6), we get

$$\begin{cases} u = 4\alpha - 6\beta \\ v = 8\beta + 2\alpha \end{cases} \quad (8)$$

Using (8) and (2), the values of x,y,z and w are given by

$$\left. \begin{aligned} x(a, b) &= 4\alpha - 6\beta + 1 \\ y(a, b) &= 4\alpha - 6\beta - 1 \\ z(a, b) &= 8\beta + 2\alpha + 1 \\ w(a, b) &= 8\beta + 2\alpha - 1 \end{aligned} \right\} \quad (9)$$

Thus, (4) and (9) represent the non-zero distinct integral solutions to (1)

Properties:

(i) $z(n, 1) + w(n, 1) - x(n, 1) + 1 = 6t_{4,n}[6CP_n^{26} + 25CP_n^6] + 432t_{3,n} - 216t_{4,n}$

(ii) $4x(n, 1) + 3w(n, 1) + 4751 = 22t_{4,n}[6F_{4,n,6} - 2CP_n^9 - 92t_{4,n} + 539]$

(iii) $p(2^n, 2^{n+1}) + 7 = 7j_{2n+2}$

Pattern: 2

Consider (3) as

$$4u^2 + 6v^2 = 22p^6 * 1 \quad (10)$$

Write 1 as

$$1 = \frac{(5+i2\sqrt{6})(5-i2\sqrt{6})}{49} \quad (11)$$

Substituting (4),(5) and (11) in (10) and employing the factorization method, define

$$2u + i\sqrt{6}v = \frac{1}{7} [(4 + i\sqrt{6}) + (5 + i2\sqrt{6})(2\alpha + i\beta\sqrt{6})]$$

Equating real and imaginary parts, we have

$$\left. \begin{aligned} u &= \frac{1}{7}(8\alpha - 78\beta) \\ v &= \frac{1}{7}(26\alpha + 16\beta) \end{aligned} \right\} \quad (12)$$

As our interest is on finding integer solutions, we choose a and b suitably so that u and v are integers. Replace a by 7a and b by 7b in (7). Substituting the corresponding values of α and β in (12) and employing (2),

non-zero integral solutions to (1) are found to be

$$\left. \begin{aligned} x(a, b) &= 7^5(8\alpha - 78\beta) + 1 \\ y(a, b) &= 7^5(8\alpha - 78\beta) - 1 \\ z(a, b) &= 7^5(26\alpha + 16\beta) + 1 \\ w(a, b) &= 7^5(26\alpha + 16\beta) - 1 \end{aligned} \right\} \quad (13)$$

Thus, (4) and (13) are non-zero distinct integer solutions to (1)

Pattern:3

In addition to (11), 1 can be written as

$$1 = \frac{(1+i2\sqrt{6})(1-i2\sqrt{6})}{25} \quad (14)$$

Following the procedure as presented in pattern.2, the corresponding non-zero distinct integral solutions to (1) are given by

$$\left. \begin{aligned} x(a, b) &= -5^5(8\alpha + 54\beta) + 1 \\ y(a, b) &= -5^5(8\alpha + 54\beta) - 1 \\ z(a, b) &= 7^5(18\alpha - 16\beta) + 1 \\ w(a, b) &= 7^5(18\alpha - 16\beta) - 1 \\ p(a, b) &= 4a^2 + 6b^2 \end{aligned} \right\} \quad (15)$$

Properties:

(i) $9x(n,1)+4w(n,1)-5 =$

$$-5^7 \cdot 22[6t_{4,n}(2CP_n^3 - 2CP_{21,n} + 22t_{4,n} + 2) + 432t_{3,n} - 216t_{4,n}]$$

(ii) $8x(1,n)-27w(1,n)+1718715 =$

$$5^7 * 396t_{4,n}[48F_{4,n,8} - 6CP_n^{30} - 3CP_n^4 - 30t_{3,n} - 27t_{4,n} + 5]$$

(iii) $p(2^{n-1}, 2^n) = 3J_{2n} + 3J_{2n+1} - 2$

Pattern:4

(3) can be written as

$$2u^2 + 3v^2 = 11p^6 \quad (16)$$

The substitution of the linear transformations

$$u = X + 3T, \quad v = X - 2T \quad (17)$$

in (16) lead to

$$5X^2 + 30T^2 = 11p^6$$

Replacing p by 5Q in the above equations, we have

$$X^2 + 6T^2 = 34375Q^6 \quad (18)$$

Let $Q = a^2 + 6b^2 \quad (19)$

$$34375 = (25 + i75\sqrt{6})(25 - i75\sqrt{6}) \quad (20)$$

Using (19) and (20) in (18) and applying the method of factorization, define

$$X + i\sqrt{6}T = (25 + i75\sqrt{6})(\alpha + i\sqrt{6}\beta) \quad (21)$$

W

where

$$\left. \begin{aligned} \alpha &= a^6 - 90a^4b^2 + 540a^2b^4 - 216b^6 \\ \beta &= 6a^5b - 120a^3b^3 + 216ab^5 \end{aligned} \right\} \quad (22)$$

Equating real and imaginary parts in (21), we have

$$\left. \begin{aligned} X &= 25\alpha - 450\beta \\ T &= 75\alpha + 25\beta \end{aligned} \right\} \quad (23)$$

Substituting (23) in (17) we get

$$\left. \begin{aligned} u &= 250\alpha - 375\beta \\ v &= -125\alpha - 500\beta \end{aligned} \right\} \quad (24)$$

Using, (24) in (2) the value of x,y,z,w and p are given by

$$x(a, b) = 250\alpha - 375\beta + 1$$

$$y(a, b) = 250\alpha - 375\beta - 1$$

$$z(a, b) = -125\alpha - 500\beta + 1$$

$$w(a, b) = -125\alpha - 500\beta - 1$$

$$p(a, b) = 5(a^2 + 6b^2)$$

Properties:

$$(i)x(n, 1) + 2z(n, 1) = 3750[t_{4,n}(2CP_n^3 - 42t_{3,n} + 216t_{4,n}) + 72t_{3,n} - 36t_{4,n}] + 3$$

$$(ii)y(n, 1) + 2z(n, 1) = 8250[-CP_n^6(S_n - t_{12,n} + Gn_n - 20) + 72t_{3,n} - 36t_{4,n}] + 1$$

$$(iii)x(n, 1) + z(n, 1) + w(n, 1) = -8250[t_{4,n}(2CP_n^3 - 2t_{3,n} + t_{4,n}) + 72t_{3,n} - 36t_{4,n}] + 1$$

Pattern:5

Also, (17) can be considered as

$$u = X - 3T, \quad v = X + 2T \quad (25)$$

Following the procedure as presented in Pattern.4, the corresponding non-zero distinct integral solutions to (1) are given by

$$x(a, b) = -200\alpha - 525\beta + 1$$

$$y(a, b) = -200\alpha - 525\beta - 1$$

$$z(a, b) = 175\alpha - 400\beta + 1$$

$$w(a, b) = 175\alpha - 400\beta - 1$$

$$p(a, b) = 5(a^2 + 6b^2)$$

IV. CONCLUSION

In this paper, we have made an attempt to determine different patterns of non-zero distinct integer solutions to the non-homogeneous heptic equation with five variables. As the heptic equations are rich in variety, one may search for other forms of heptic equation with variables greater than or equal to five and obtain their corresponding properties.

ACKNOWLEDGEMENT

The finical support from the UGC, New Delhi (F.MRP-5123/14(SERO/UGC) dated march 2014) for a part of this work is gratefully acknowledged.

REFERENCES

[1]. Dickson.L.E, History of Theory of Numbers, Vol.11, Chelsea Publishing company,New York (1952).
 [2]. Mordell.L.J, Diophantine eequations, Academic Press, London (1969).
 [3]. Telang,S.G., Number theory, Tata Mc Graw Hill publishing company, New Delhi (1996)
 [4]. Carmichael, R.D., The theory of numbers and Diophantine Analysis, Dover Publications, New York (1959).
 [5]. Gopalan.M.A and Sangeetha.G., On the sextic equations with three unknowns $x^2 - xy + y^2 = (k^2 + 3)^n z^6$, Impact J.Sci.tech., Vol.4, No:4, 89-93,2010
 [6]. Gopalan. M.A, Manju Somnath and Vanitha.N, Parametric Solutions of $x^2 - y^6 = z^2$,Acta ciencia indica, XXXIII,3,1083-1085,2007
 [7]. Gopalan M.A and VijayaSankar.A, Integral Solutions of the sextic equation $x^4 + y^4 + z^4 = 2w^6$, Indian Journal of Mathematics and mathematicalsciences, Vol.6, No2, 241-245.2010
 [8]. Gopalan M.A, Vidhyalakshmi.S and Vijayasankar.A, Integral solutions of Non-Homogeneous sextic equation $xy + z^2 = w^6$, impact j.Sci.tech., Vol.6, No:1, 47-52, 2012
 [9]. Gopalan M.A, Vidhyalakshmi.S and Lakshmi.K, On the non-homogeneous sextic equation $x^4 + 2(x^2 + w) + x^2 y^2 + y^4 = z^4$,IJAMA, 4(2), 171-173, Dec-2012
 [10].Gopalan M.A, Sangeetha.G, On the heptic Diophantine equation with five unknowns $x^4 - y^4 = (X^2 - Y^2)z^5$, Antarctica J.Math., 9(5), 371-375,2012

- [11]. Manjula.M, Gopalan M.A and Vidhyalakshmi.S, On the non-homogeneous heptic equation with five unknowns
 $(x^2 - y^2)(4x^2 + 4y^2 - 6xy) = 11(X^2 - Y^2)z^5$, International Conference on mathematical Methods and computation, PP. 275-278,2014
- [12]. Gopalan M.A, Vidhyalakshmi.S and Lakshmi.K, Integral solution of the non-homogeneous heptic equation with five unknowns
 $x^4 + y^4 - (x - y)z^3 = 2(k^2 + 6s^2)w^2T^5$, SJET, Vol.2, issue.2, 212-218, Mar-2014
- [13]. Gopalan M.A and Sangeetha.G, Integral solutions of Ternary non-homogeneous biquadratic equation
 $x^4 + x^2 + y^2 - y = z^2 - z$, Acta Ciencia Indica, XXXVIIM(4), 799-803,2011
- [14]. Gopalan M.A and Sangeetha.G, Integral solutions of Ternary Quadratic equation
 $x^2 + y^2 + 2xy = z^4$, Antarctica J.math., 7(1), 95-101,2010
- [15]. Gopalan M.A, Manju Somanath and Sangeetha.G, Integral solutions of Ternary non-homogeneous quartic equation
 $x^4 - y^4 = (k^2 + 1)(z^2 - w^2)$, Archimedes J.Math., 1(1), 51-57, 2011
- [16]. Gopalan M.A and Sangeetha.G, Integral solutions of Ternary Quintic Diophantine equation
 $x^2 + y^2 = (k^2 + 1)z^5$, Bulletin of Pure and Applied Sciences, 29E(1), 23-28,2010
- [17]. Gopalan M.A and Janaki.G, Integral solutions of
 $(x^2 - y^2)(3x^2 + 3y^2 - 2xy) = 2(z^2 - w^2)p^3$, Impact J. Sci.,Tech., 4(1),97-102,2010
- [18]. Gopalan M.A and Vijayashankar.A, Integral solutions of Ternary Quintic Diophantine equation
 $x^2 + (2k + 1)y^2 = z^5$, International Journal of mathematical sciences, 9(1-2), 165-169,(Jan-Jun 2010).
- [19]. Gopalan M.A, and Sangeetha.G, Parametric Integral solutions of the heptic equation with five unknowns
 $x^4 - y^4 + 2(x^3 + y^3)(x - y) = 2(X^2 - Y^2)z^5$, Bessel J,Math., 1(1),17-22 ,2011
- [20]. Gopalan M.A and Sangeetha.G, Integral solutions of m^{th} degree non-homogeneous equation with three unknowns
 $a(x - y)^2 + xy = [1 + (4a - 1)k^2]^n z^m$, South East Asian J.Math.&Math.,Sci.,9(3),33-38, 2011
- [21]. Manju somanath, Sangeetha.G and Gopalan M.A, Observations on the higher degree Diophantine equation
 $x^2 + y^2 = (k^2 + a^2)z^m$, Impact J.Sci. Tech., 5(1), 67-70,2010
- [22].Manju somanath, Sangeetha.G and Gopalan M.A, On the non-Homogeneous heptic equation with three unknowns
 $x^3 + (2^p - 1)y^5 = z^7$, Diophantus J.Math.,1(2), 117-121,2012
- [23]. Manju somanath, Sangeetha.V and Gopalan M.A On the non-homogeneous heptic equation with five unknowns
 $(x^2 - y^2)(4x^2 + 4y^2 - 6xy) = 8(X^2 - Y^2)z^5$, International journal of Innovative Research and review, Vol.2, issue-4, 23-26, Dec-2014
- [24]. Manju somanath, Sangeetha.V and Gopalan M.A Integral Solutions of the non-homogeneous heptic equation with five unknowns
 $(x^3 - y^3) - (x^2 + y^2) + z^3 - w^3 = 2 + 5(x - y)(z - w)^2 p^4$, International journal of scientific & research publications Vol.5, issue-1, Jan-2015
- [25]. P.Jayakumar, K.Sangeetha, On the non homogeneous heptic equation with five unknowns
 $(x^2 - y^2)(9x^2 + 9y^2 - 16xy) = 21(X^2 - Y^2)Z^5$
 International journal of science and Research, Vol.3, issue.11, 3163-3165,

2014

- [26]. Manju somanath, Sangeetha.V and Gopalan M.A, On the heptic equations with three unknowns $3(x^2 + y^2) - 5xy = 15z^7$,The IJST, Vol.2, issue.2,26-28, 2014