

Review on Oral dosage forms

Swami Datta¹ Renuka R. Deshpande² Nandkishor B. Bavage³, Vidyasagar Gali⁴, Shyamlila B. Bavage⁵

¹B.Pharmacy Final Year Student, Latur College of Pharmacy Hasegaon, Tq. Ausa, Dist. Latur-413512, Maharashtra, India

²Assistant Professor, Department of Pharmaceutics, Latur College of Pharmacy Hasegaon, Tq. Ausa, Dist. Latur-413512, Maharashtra, India

^{3,4}Department of Pharmaceutical Analysis, Latur College of Pharmacy Hasegaon, Tq. Ausa, Dist. Latur-413512, Maharashtra, India

⁵Department of Pharmacognosy, Latur College of Pharmacy, Hasegaon, Tq. Ausa, Dist. Latur-413512, Maharashtra, India

Abstract- Dosage forms are the vehicles for administering drug in to the body. Oral dosage forms can be solid, liquid, and gaseous; some may be inhalers, pellets, and granules. Route of administration can be oral, topical, rectal, vaginal, and ophthalmic and also suppositories. These dosage forms differ for infant's children and adults. Based on the amount of drug it is prescribed differently for each individual. There is different mechanism of action for different dosage forms.

Index terms- Tablets, Capsule, Drug delivery

INTRODUCTION

Oral Solid indefinite quantity forms (Tablets and Capsules) are a number of the foremost fashionable and convenient ways of drug delivery. They will be made in exceedingly non-sterile surroundings and also the method, instrumentality and technology is well outlined and acknowledged, when quite a hundred years of development. With the high volume of merchandise made in these indefinite quantity forms, it's necessary that the unit operations for his or her production be totally understood, developed and enforced. This course focuses on the basics of every distinct process step (unit operation) with intensive discussions on these technologies needed for the producing and packaging of tablets and capsules.

Dosage forms are classified into solid, liquid and semi-solid forms. Solid dosage forms are of two types- Internal and external. Tablet is a dosage form which contains drug in it and also additives, preservatives, binders, diluents. In case of Buccal tablet, tablets are placed between the cheek and the

gum [1-10]. Capsule is cylindrical in shape. It has two parts: cap and body. Drug is filled in between the cap and body. They are of two types' hard gelatin and soft gelatin capsules. External can be Tooth powders and dusting powders. Tooth powders can be of different type's detergents. Preservatives and colors etc. Liquid dosage forms can be classified as monophasic and biphasic phase means direction.




Figure 1. Different dosage forms

Monophasic dosage forms include syrup. Elixir. and Oral drops. This syrup contains preservatives .additives, binding agent's etc .These syrups contain drug inside in liquid it shows effect easily. Proper dose should be maintained. Biphasic dosage forms can be suspension or emulsion. Suspension contains drug in suspended form, where in emulsion contains drug in the form two immiscible liquids. Emulsions are usually water in oil or oil in water type. Semisolid dosage forms are external and internal. They usually applied topically on skin. They include ointments, lotions, liniments .internal include pessaries and suppositories rectal or vaginal devices.



Figure 2. Capsules

There is another route of administration of drug which includes parenterals. These are sterile preparations which contains drug in the form of ampoules or vials. These injections can be intravenous (to vein) intramuscular (to muscle) or subcutaneous (to skin). Inhaler is also used for asthma patients. Nebulizer is used for reducing blockage in the nostrils for small children [21-40].

Advantages of solid dosage forms include

- Safe, less expensive, immediate relief
- Accurate dose, releases drug over a long period
- Can be used for all age groups, convenient, cheap
- Absorption is fast, pain less
- Less space for storage
- Less spoilage and less microbial contamination
- Action is more rapid in liquid dosage form



Figure 3. Syrup: A liquid form



Figure 4. Injections

Disadvantages of solid dosage forms include

- Bitter in taste, Difficulty in swallowing
- Irritation to gastro intestinal tract–nausea and vomiting
- Unable to use for unconscious patients
- For liquid dosage form storing is problematic
- Types of release drug in the human body
- Controlled release: It delivers drug at a predetermined place
- Sustained release: It releases drug as two portions once at a place and after some time in some other place
- Release rate is not constant [41-60]

Defects of tablets

- Capping; breaking of tablet horizontally
- Sticking: sticking to the die walls [61-88]
- Motling: it is due to improper mixing of color
- Picking: it is due to improper drying and lubrication of granules
- Cracking: cracking occurs due to improper granulation



CONCLUSION

Oral dosage forms can be a tablet, a capsule or drops. This route of administration is safe and results are good. These are cheaper and everyone will be aware of some tablets. The only problem of tablets is acidity. This will be balanced by taking anti-ulcer agents.

REFERENCES

- [1] Bapuji AT, et al. Bioequivalence Testing- Industry Perspective. J Bioequiv Availab. 2010; 2: 098-101.

- [2] Semdé R, et al Bioequivalence in vitro evaluation of some antibacterial generic dosage forms. *J Pharm Belg.* 2005; 60: 51-55.
- [3] Fassin D, et al. The Bamako Initiative-Letter. *Lancet* 1989; 333; 162-163.
- [4] Sagui E. Stroke in sub-Saharan Africa. *Med Trop. (Mars)* 2007; 67: 596-600.
- [5] Sauwakon R and Wondemagegnehu E. Effective drug regulation: A multicounty study. *World Health Organization* 2002: 1-202.
- [6] Polli JE. In vitro studies are sometimes better than conventional human pharmacokinetic in vivo studies in assessing bioequivalence of immediate-release solid oral dosage forms. *AAPS J*, 2008; 10:289-299.
- [7] Anand O, et al. Dissolution testing for generic drugs: an FDA perspective. *AAPS J* 2011; 13: 328-335.
- [8] Abelli C, et al. Equivalence pharmaceutiques des médicaments essentiels génériques. *STP Pharma Pratiques* . 2001; 11: 89-101.
- [9] Vander Meersch A. Quality of reporting of bioequivalence trials comparing generic to brand-name drugs: a methodological systematic review. 2011; *PLoS One* 6: e23611.
- [10] Burgess DJ. Assuring quality and performance of sustained and controlled release parenterals: Workshop report. *AAP S pharm Sci B.* 2002; 4: 13-23.
- [11] Brown CK, et al. FIP/AAPS joint workshop report: dissolution/In vitro release testing of Novel/Special dosage forms. *AAPS Pharm SciTech.* 2011; 12: 782-794.
- [12] Siewert M, et al. FIP/AAPS guidelines to dissolution/In vitro release testing of novel/special dosage forms. *AAPS Pharm Sci Tech.* 2003; 4:43-52.
- [13] Seidlitz A, et al. Biorelevant Dissolution Testing of Drug-Eluting Stents: Experiences with a Modified Flow-Through Cell Setup. *Dissol Technol.* 2011; 18: 26-34.
- [14] Bhardwaj U and Burgess DJ. A novel USP apparatus 4 based release testing method for dispersed systems. *Int J Pharm.* 388: 287-294.
- [15] Rebucci M and Michiels C. Molecular aspects of cancer cell resistance to chemotherapy. *Biochem Pharmacol.* 2013; 85: 1219-1226.
- [16] Wu CP, et al. The emergence of drug transporter-mediated multidrug resistance to cancer chemotherapy. *Mol Pharm.* 2011; 8: 1996-2011.
- [17] Moulder S. Intrinsic resistance to chemotherapy in breast cancer. *Womens Health (Lond Engl)* 2010; 6: 821-830.
- [18] Ullah MF. Cancer multidrug resistance (MDR): a major impediment to effective chemotherapy. *Asian Pac J Cancer Prev.* 2008; 9: 1-6
- [19] Pauwels EK, et al. Multidrug resistance in cancer: its mechanism and its modulation. *Drug News Perspect.* 2007; 20: 371-377.
- [20] Teoh D and Secord AA. Antiangiogenic agents in combination with chemotherapy for the treatment of epithelial ovarian cancer. *Int J Gynecol Canc.* 2012; 22: 348-359.
- [21] Slovin S. Chemotherapy and immunotherapy combination in advanced prostate cancer. *Clin Adv Hematol Oncol.* 2012; 10: 90-100.

AUTHOR DETAIL



Swami Datta Umakant
Student of B.pharmacy 4th year, Latur
College of Pharmacy, Hasegaon



Renuka R. Deshpande,
Assistant Professor, Department of
Pharmaceutics, Latur College of
Pharmacy, Hasegaon