

Structural Elucidation of Some Coordination Compounds with Reference to Thermal Studies

Vijay Kumar¹, Dr. R.P.S. Chauhan²

¹Research Scholar, Department of Chemistry, Magadh University, Bodh-Gaya

²Professor & Head, University Department of Chemistry, Magadh University, Bodh-Gaya

Abstract - This paper reveals that Novel mixed ligand Co(II), Ni(II), Cu(II) and Zn(II) complexes of Schiff base derived through the condensation of furfurylidene-4-aminoantipyrine and 2-aminobenzothiazole with 2-aminophenol have been synthesized and characterized by various analytical and spectral techniques. The spectral results suggest octahedral geometry for Co (II), Ni (II) and Zn (II) complexes. Distorted octahedral geometry has been assigned for Cu (II) complex. The low molar conductance values of the complexes support their neutral nature. The thermal stability of the complexes was determined using TG/DTA studies. The electrochemical behavior of the complexes was studied using cyclic voltammetry. The grain size of the complex was calculated by Scherrer formula using powder XRD. Surface morphology of the complexes was determined by SEM analysis.

Index Terms - Schiff Base, Mixed Ligand Complexes, TG/DTA, XRD

INTRODUCTION

Schiff base ligands have significant importance in chemistry; especially in the development of Schiff base complexes because Schiff base ligands are potentially capable of forming stable complexes with metal ions (Souza et al., 1985). Mixed ligand complexes of transition metals containing ligands with N, S or N, S, O donors are known to exhibit interesting stereochemical, electrochemical and electronic properties (Prabhakaran et al., 2005).

In recent decades, a great deal of interest in the metal complexes of nitrogen-oxygen chelating agents derived from 4-aminoantipyrine Schiff bases have various applications in antifungal, antibacterial, analgesic, sedative, antipyretic, anti-inflammatory and greater DNA binding ability (Kumaran et al., 2013) Heterocycles containing thiazole ring is present in a number of pharmacologically and biologically active compounds. Compounds containing benzothiazole

derivatives were used as antifungal, antiinflammatory, anti-HIV, anticancer, anticarbonic anhydrase, diuretic, hypoglycaemic, antithyroid, antimalarial and in therapeutic fields (Neelakantan et al., 2010) The metal complex-DNA interactions have received much importance for the development of new metal-based chemotherapeutic drugs (Afrati et al., 2009) The numerous biological experiments performed so far suggest that DNA is the primary intracellular target of anticancer drugs because the interaction between small molecules and DNA can induce DNA damages in cancer cells, blocking the division of cancer cells and resulting in the cell death (Li et al., 2009).

By considering the above, in this paper, we synthesized and characterized novel Co (II), Ni (II), Cu (II) and Zn (II) mixed ligand complexes.

MATERIALS AND METHODS

Chemicals and reagents

The chemicals used were of AnalaR grade, furfuraldehyde, 4-aminoantipyrine, 2-aminophenol and 2-aminobenzothiazole were obtained from Sigma Aldrich. Metal (II) acetates were obtained from Merck and were used as received. The solvents used were purchased from Merck and used without further purification.

Physical Measurements

Elemental analysis of ligand and its metal complexes were carried out using Perkin-Elmer elemental analyzer. Molar conductance of the complexes was measured using a coronation digital conductivity meter. IR spectra were recorded using Jasco FTIR-410 spectrometer in KBr pellets from 4000-400 cm^{-1} . ^1H NMR spectra were recorded with Bruker 300 MHz spectrometer using CDCl_3 solvent for ligand and DMSO- d_6 for Zn (II) complex with TMS as internal

standard. DART-MS spectrum was recorded on a JEOL-Accu TOF JMS mass spectrometer.

Magnetic moments were measured by Guoy method and corrected for diamagnetism of the component using Pascal's constants. Electronic spectra were recorded on Thermo Scientific Evolution-200 UV-Visible spectrophotometer in the range 190-1100 nm. ESR spectrum of the Cu (II) complex was recorded at 300 and 77 K in DMSO solution using Varian, USA E-112 ESR spectrometer using tetracyanoethylene (TCNE) as g-marker. Thermal analysis was carried out under nitrogen atmosphere at a heating rate of 10 °C per minute using Perkin Elmer Diamond TG/DTA analyzer. SEM images were recorded in a Hitachi SEM analyzer.

Synthesis of Mixed Ligand Complexes

The Schiff base was prepared using the reported procedure (Leelavathy *et al.*, 2013). The complexes were prepared by following the procedure. A methanolic solutions of Schiff base (0.004 mol), Co(II)/ Ni(II)/ Cu(II)/ Zn(II) metal acetates (0.004 mol) and 2-aminophenol (2-ap) (0.004 mol) were taken in 1:1:1 molar ratio and stirred with heating for about 4 h. The resulting mixture is then cooled to room temperature and the solid product formed was filtered, washed with methanol, and dried over anhydrous calcium chloride.

RESULTS AND DISCUSSION

The mixed ligand complexes synthesized are stable at room temperature. They are soluble in DMF and DMSO. The analytical, physical properties and molar conductance data of the complexes are given in Table: 1. Elemental analysis indicates that the found and calculated values were within acceptable limits (\pm

0.5). The molar conductance data of the mixed ligand complexes fall in the range of 3.3-12.5 $\Omega^{-1}\text{cm}^2\text{mol}^{-1}$, which confirms the present complexes, are non-electrolytic in nature (Geary, 1971).

IR Spectra

A comparative study of IR spectra of the mixed ligand complexes with ligand reveals that several peaks are shifted, vanished or have newly appeared. The infrared spectrum of the Schiff base showed two strong bands, one at 1590 cm^{-1} and another at 1652 cm^{-1} . These bands are assigned to the (-CH=N) and (>C=N) stretching frequency vibrations respectively (Ali *et al.*, 2013).

A shift in these frequencies (Table: 2) is observed in all complexes due to the coordination of metal through nitrogen atoms of the -CH=N and >C=N groups. In all the complexes a broad band is present in the region 3435-3406 cm^{-1} , indicates the presence of coordinated water molecules (Mashaly *et al.*, 2004). The band at 3354 cm^{-1} for $\nu(\text{OH})$ in the free 2-aminophenol ligand disappeared on complexation, indicating coordination of -OH group through deprotonation (Krishnankutty *et al.*, 2007). The absence of the bands due to NH_2 group in the spectra of complexes indicating the involvement of amino group in coordination through deprotonation.

The complexes also display bands in the 439-417, 590-497 cm^{-1} region due to the formation of M-N and M-O bonds respectively (El-Shahawi *et al.*, 2013). From the IR spectra, it is concluded that the Schiff base behaves as a bidentate ligand coordinated to the metal ions via (-CH=N) and (>C=N) groups. The other coordination sites of metal (II) ions were occupied by two water molecules and two positions were occupied by the -OH and NH_2 groups of 2-aminophenol through deprotonation.

Table 1: Physical properties and analytical data of mixed ligand complexes

Compound	Yield (%)	Molecular weight	Elemental analysis (%) Found (Calculated)						Molar conductivity ($\Omega^{-1}\text{cm}^2\text{mol}^{-1}$)
			C	H	N	O	S	M	
$\text{C}_{29}\text{H}_{28}\text{CoN}_6\text{O}_4\text{S}$	67	615.57	56.73 (56.58)	4.38 (4.58)	13.62 (13.65)	10.35 (10.40)	5.17 (5.21)	9.68 (9.57)	3.3
$\text{C}_{29}\text{H}_{28}\text{NiN}_6\text{O}_4\text{S}$	52	615.33	56.70 (56.61)	4.35 (4.59)	13.78 (13.66)	10.60 (10.40)	5.13 (5.21)	9.42 (9.54)	12.5
$\text{C}_{29}\text{H}_{28}\text{CuN}_6\text{O}_4\text{S}$	72	620.18	56.43 (56.16)	4.43 (4.55)	13.61 (13.55)	10.72 (10.32)	5.29 (5.17)	10.21 (10.55)	8.6
$\text{C}_{29}\text{H}_{28}\text{N}_6\text{O}_4\text{SZn}$	58	622.03	56.21 (56.00)	4.59 (4.54)	13.42 (13.51)	10.50 (10.29)	5.35 (5.15)	10.31 (10.51)	5.5

Table 2: IR spectral data of ligand and its complexes (cm⁻¹)

Compound	$\nu(\text{HC}=\text{N})$	$\nu(\text{OH})_{\text{water}}$	$\nu(\text{C}=\text{N})$	$\nu(\text{M}-\text{N})$	$\nu(\text{M}-\text{O})$
Ligand	1590	-	1646	-	-
Co(II) complex	1593	3420	1644	438	590
Ni(II) complex	1572	3406	1644	439	590
Cu(II) complex	1587	3435	1641	417	497
Zn(II) complex	1581	3432	1639	430	588

Electronic Spectra and Magnetic Susceptibility

The electronic absorption spectra of the ligand and its mixed ligand complexes were recorded at room temperature in DMF solution. The absorption assignments and magnetic susceptibility values of the complexes along with the geometry are given in Table 3. The electronic spectra of the ligand show broad band 318 and 370 nm which can be assigned to $\pi-\pi^*$ transitions of the azomethine ($-\text{CH}=\text{N}$) chromophore. On complexation this was shifted to lower wavelength, suggesting the coordination of azomethine nitrogen. In addition, other intense absorption band at higher energy 220-265 nm is due to $\pi-\pi^*$ transition of the benzene ring of the Schiff base ligand. The electronic spectrum of Co(II) complex show two absorption bands one at ~ 1075 nm and another one at 345 nm assignable to ${}^4\text{T}_{1g}(\text{F}) \rightarrow {}^4\text{T}_{2g}(\text{F})$ and ${}^4\text{T}_{1g}(\text{F}) \rightarrow {}^4\text{T}_{1g}(\text{P})$ transitions respectively, which

Table 3: Electronic spectral data and magnetic moment values of Metal complexes

Complex	λ_{max} (nm)	Band assignments	Magnetic moment (BM)	Geometry
Co (II) complex	1075	${}^4\text{T}_{1g}(\text{F}) \rightarrow {}^4\text{T}_{2g}(\text{F})$	5.14	Octahedral
	345	${}^4\text{T}_{1g}(\text{F}) \rightarrow {}^4\text{T}_{1g}(\text{P})$		
Ni (II) complex	387	${}^3\text{A}_{2g}(\text{F}) \rightarrow {}^3\text{T}_{1g}(\text{P})$	3.20	Octahedral
Cu (II) complex	464	${}^2\text{B}_{1g} \rightarrow {}^2\text{E}_{1g}$	1.91	Distorted Octahedral
	287	$\pi-\pi^*$		
Zn (II) complex	335	$n-\pi^*$	Diamagnetic	Octahedral

¹H NMR Spectra

The ¹H NMR spectra of ligand and Zn (II) complex were recorded in DMSO d₆. The signal for azomethine proton ($-\text{CH}=\text{N}-$) in the ligand appears as a singlet at 9.6 ppm (Amer et al., 2013). In the ¹H NMR spectrum of the Zn (II) complex, the azomethine proton signal is shifted downfield compared to the free ligand due to the deshielding of the azomethine group on coordination with Zn (II) ion. The multiple obtained in the 7.2-7.5 ppm range is due to the aromatic protons of the ligand as expected. The signal for pyrazolone ring carbon attached methyl proton ($-\text{CH}_3$) appear as a singlet at 2.5 ppm, while pyrazolone nitrogen

is characteristic for the octahedral Co(II) complex. The magnetic moment value of the Co (II) complex is 5.14 BM. The electronic spectrum of the Ni (II) complex shows one absorption band at 387 nm assignable to ${}^3\text{A}_{2g}(\text{F}) \rightarrow {}^3\text{T}_{1g}(\text{P})$ transition. This is characteristic of six coordinated octahedral Ni (II) complexes. The magnetic moment value is 3.20 BM for the Ni (II) complex confirms six coordinate octahedral geometry. The distorted octahedral Cu (II) complex, display one absorption band at 464 nm, corresponding to ${}^2\text{B}_{1g} \rightarrow {}^2\text{E}_{1g}$ transition. The μ_{eff} for the complex is 1.91 BM, which is characteristic for distorted octahedral geometry around Cu (II). The Zn (II) complex exhibited two intra ligand transitions at 287 and 335 nm and is diamagnetic. According to the empirical formula, an octahedral geometry is proposed for the Zn (II) complex (Nag *et al.*, 2005; Banerjea, 1993; Kettle, 1969).

attached methyl protons ($>\text{N}-\text{CH}_3$) appear as a singlet at δ 3.3 ppm (Aupama et al., 2012).

The peaks at 6.5 and 6.8 ppm are due to furfuryl protons.

Mass Spectra

The mass spectrum of ligand and its Co (II), Ni (II), Cu (II) and Zn (II) mixed ligand complexes were recorded. The molecular ion peak for ligand, L (C₂₃H₁₉N₅O₅) was observed at 413 *m/z*. Whereas the molecular ion peaks of Co (II), Ni (II), Cu (II) and Zn (II) mixed ligand complexes were observed at 617, 616, 621 and 623 *m/z* which confirms the stoichiometry of the metal complexes to be

[ML(H₂O)₂(2-ap)]. The mass spectra of ligand and its complexes exhibited other molecular ion peaks for several fragments. Elemental analysis values are in agreement with the values calculated from the molecular formulae assigned to these complexes which are further supported by DART-mass studies.

ESR Spectra

ESR spectra of the Cu (II) complex was recorded at room temperature and at liquid nitrogen temperature in DMSO. The g_{\parallel} and g_{\perp} values were calculated from the spectrum using tetracyanoethylene (TCNE) free radical as the “g” marker. In the present Cu (II) complex, g_{\parallel} is less than 2.3 is an indication of significant covalent character to M-L bond. The g_{\parallel} and g_{\perp} values are >2.04 , consistent with an elongated tetragonally distorted octahedral stereochemistry. The G factor [$G = (g_{\parallel}-2)/(g_{\perp}-2)$] is >4.0 suggests that the exchange interactions between Cu(II) centers in the solid state are negligible (El-Bindary *et al.*, 2000). The absence of half field signal at 1600 G due to the $\Delta m_s = \pm 2$ transitions, ruling out any Cu-Cu interaction. The g values are in the order $g_{\parallel} > g_{\perp} > 2.0023$ corresponding to the presence of an unpaired electron in the $d_{x^2-y^2}$ orbital. From the observed values, it is clear that $A_{\parallel} = 110 > A_{\perp} = 82$; $g_{\parallel} = 2.13 > g_{\perp} = 2.03 > 2$ and the ESR parameters of the Cu(II) complex suggest that the complex have tetragonally distorted octahedral geometry and the unpaired electron lies predominantly in the $d_{x^2-y^2}$ orbital.

Thermal Studies

The thermal stabilities of ligand and its mixed ligand metal complexes were investigated using TG and DTA under nitrogen atmosphere with a heating rate of 10 °C per minute from 40 °C to 700 °C. The Co (II) and Zn (II) complexes show four stages of decomposition. While the Ni (II) and Cu (II) complexes complex exhibited three stage of decomposition. Thermograms of Co (II), Ni (II), Cu (II) and Zn (II) complexes show weight loss around 151-223 °C indicates the presence of two coordinated water molecules in these complexes (Dhanaraj *et al.*, 2014). The other decomposition stages correspond to the loss of 2-aminophenol ligand and decomposition of organic moieties of the Schiff base ligand. The initial decomposition temperature of Co (II) complex is higher compared to the remaining complexes, indicating it is more thermally stable. The Zn (II)

complex shows the final decomposition at 457 °C, which is the higher final decomposition temperature compared to the rest of the complexes.

Based on the elemental analysis, IR, electronic spectra, magnetic moments, ¹H NMR, mass, ESR spectral data and thermal analysis, the proposed structure of the complexes are given in Figure: 1.




Figure 1: Proposed structure of metal complexes (where M=Co (II), Ni (II), Cu (II) and Zn (II))




Figure 2: Powder XRD pattern of a) Co (II), b) Ni (II), c) Cu (II) and d) Zn (II) complexes




Figure 3: Powder SEM images of a) Co (II), b) Ni (II), c) Cu (II) and d) Zn (II) complexes

Powder XRD

The X-ray diffraction pattern of the mixed ligand complexes is shown in Figure: 2(a-d). From the observed d_{XRD} patterns, the crystalline size of the complexes were calculated from Scherrer's formula $d_{XRD} = 0.9\lambda / \beta \cos\theta$, where λ is the wave length, β is the full-width half maximum of the characteristic peak and θ is the diffraction angle for the hkl plane (Tabassum *et al.*, 2012). In the present study, Co (II), Ni (II), Cu (II) and Zn (II) complexes are amorphous in nature.

SEM

The SEM micrographs of the Co (II), Ni (II), Cu (II) and Zn (II) mixed ligand complexes are shown in Figure: 3(a-d). The SEM micrograph of Co (II) complex exhibit spherical structured particles with small grains. Ni (II) complex exhibited irregularly shaped particles with small grains. The Cu (II) and Zn (II) complexes showed cauli flower like morphology.

CONCLUSION

Mixed ligand Co(II), Ni(II), Cu(II) and Zn(II) complexes with Schiff base derived from furfurylidene-4- amino antipyrine and 2-aminobenzothiazole and 2-aminophenol have been synthesized and characterized by elemental analysis, molar conductance, magnetic moment and spectral data. DART mass spectra gave the molecular weight of the complexes. The physico-chemical data suggest octahedral geometry for Co (II), Ni (II) and Zn (II) complexes and distorted octahedral geometry for Cu (II) complex. Powder XRD studies revealed that all the complexes are amorphous. The complexes exhibit different surface morphologies.

REFERENCES

- [1] Afrati T, Pantazaki AA, Dendrinou-Samara C, Raptopoulou C, Terzis A and Kessissoglou DP (2009). Copper inverse-9-metallacrown-3 compounds interacting with DNA. Dalton Transactions 39 765-775.
- [2] Ali A, Abdullah N and Maah MJ (2013). Synthesis, Characterization and Antioxidant Studies on 4-Phenyl-1,3,5-triazine-2,6-diamine Schiff Bases and Their Nickel (II), Copper (II) and Zinc (II) Complexes, Asian Journal of Chemistry. 25 (6) 3105-3108.
- [3] Amer S, El-Wakiel N and H El-Ghamry (2013). Synthesis, spectral, antitumor, and antimicrobial studies on Cu (II) complexes of purine and triazole Schiff base derivatives. Journal of Molecular Structure 1049 326–335.
- [4] Anupama B, Padmaja M and Kumari CG (2012). Synthesis, Characterization, Biological Activity and DNA Binding Studies of Metal Complexes with 4-Aminoantipyrine Schiff Base Ligand, E-Journal of Chemistry 9(1) 389-400.
- [5] Banerjea D (1993). Coordination Chemistry. Tata McGraw-Hill.
- [6] Dhanaraj CJ and Johnson J (2014). Synthesis, characterization, electrochemical and biological studies on some metal (II) Schiff base complexes containing quinoxaline moiety. Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy 118 624–631.
- [7] El-Bindary AA and El-Sonbati AZ (2000). Synthesis and Properties of Complexes of Copper (II), Nickel (II), Cobalt (II) and Uranyl Ions with 3-(p-Tolylsulphonamido) rhodanine. Polish Journal of Chemistry 74 615–620.
- [8] El-Shahawi MS, Al-Jahdali MS, Bashammakh AS, Al-Sibaai AA and HM Nassef (2013). Spectroscopic and electrochemical characterization of some Schiff base metal complexes containing benzoin moiety. Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy 113 459–465.
- [9] Geary WJ (1971). The use of conductivity measurements in organic solvents for the characterisation of coordination compounds. Coordination Chemistry Reviews 7 8-122. Kettle SFA 1969. Coordination Compounds. ELBS, Essex, UK.
- [10] Krishnankutty K, Sayudevi P and Ummathur MB (2007). Metal complexes of Schiff's bases derived from 3-(aryloxy)-2,4-pentanediones with 2-aminophenol and 2-aminothiophenol, Journal of Serbian Chemical Society 72(11) 1075–1084.
- [11] Kumaran JS, Priya S, Muthukumaran J, Jayachandramani N and Mahalakshmi S (2013). Synthesis, characterization, and biological studies of Cu (II), Co (II), Ni (II) and Zn (II) complexes of tetradentate Schiff base ligand. Journal of

Chemical and Pharmaceutical Research 5(7) 56-69.

antimicrobial studies. Journal of Photochemistry and Photobiology B 114 108–118.

- [12] Leelavathy C and Arulantony S (2013). Synthesis, spectral characterization and biological activity of metal (II) complexes with 4-aminoantipyrine derivatives. *Spectrochim Acta Part A: Molecular and Biomolecular Spectroscopy*. 113 346-355.
- [13] Li Y and Yang Z (2009). DNA binding affinity and antioxidative activity of copper (II) and zinc (II) complexes with a novel hesperetin Schiff base ligand. *Inorganica Chimica Acta* 362 4823–4831.
- [14] Mashaly MM, Abd-Elwahabb ZH and Faheimb AA (2004). Preparation, Spectral Characterization and Antimicrobial Activities of Schiff Base Complexes Derived from 4-Aminoantipyrine. Mixed Ligand Complexes with 2-Aminopyridine, 8-Hydroxyquinoline and Oxalic Acid and their Pyrolytical Products, *Journal of the Chinese Chemical Society* 51 901-915.
- [15] Nag JK, Pal S and Sinha C (2005). Synthesis and characterization of cobalt (II), nickel (II), copper (II), palladium (II) and dioxouranium (VI) complexes of the antipyrine Schiff base of 3-formylsalicylic acid. *Transition Metal Chemistry* 30 523-526.
- [16] Neelakantan MA, Esakkiammal M, Mariappan SS, Dharmaraja J and Jeyakumar T (2010). Synthesis, Characterization and Biocidal Activities of Some Schiff Base Metal Complexes. *Indian Journal of Pharmaceutical Sciences* 72 (2) 216-222.
- [17] Prabhakaran R, Karvembu R, Hashimoto T, Shimizu K and Natarajan K (2005). Formation of structurally different solvated and non-solvated [Ni (PTSC)(PPh₃)] (PTSC = salicylaldehyde-N-phenylthiosemicarbazide anion) crystals from single pot. *Inorganic Chimica Acta* 358 2093-2096.
- [18] Souza P, Garcia-Vazquez JA and Masaguer JR (1985). Synthesis and characterization of copper (II) and nickel (II) complexes of the Schiff base derived from 2-(2-aminophenyl) benzimidazole and salicylaldehyde. *Transition Metal Chemistry* 10 410-412.
- [19] Tabassum S, Zaki M, Arjmand F and Ahmad I (2012). Synthesis of heterobimetallic complexes: In vitro DNA binding, cleavage, and