

# Ragi flour enriched with fish pasta products produced from differentiation between microwave and steam heat treatment cooked defatted Pangasius meat

S. Kothandaperumal

*Department of Fish Processing Technology, Thoothukudi, Schools of Post- Harvest Technology*

**Abstract - Pangasius fish contains more unfit fats. It will cause heart and brine diseases. It contains more essential amino acid and more good fat and nutritional content. It mostly contains high amount of saturated and mono-unsaturated fats and less amount of poly-unsaturated fats. This affects marketability value and due to its bad odour most of people are disliking it. Fats in it are removed or reduced by use the different between heat treatment such as microwave oven method, steam. These methods are used to render fats and converted to pasta products. This pasta product does not contain essential amino acid due to heat treatment. Based on differentiation from between heat treatment methods, the study of nutritional differentiation and sensory evaluation on ragi pasta product is done.**

vitamins and fatty acid composition (Garcia- Arisa et al., 2003). Cooking was a very important factors which determines the final content of nutrition in fish (Badian et al., 2013). Pangasius had high amount of SFA and MUFA which affect the quality of product and also affect the consumer accessibility. Different heat treatment such as microwave heat, grill oven and steaming were followed to remove the above fatty acids, Thereby to improve the nutritional quality and this process is called as Defatting. EPA and DHA are the remaining stable fatty acids during certain type of cooking (Larsen et al., 2011). Vitamins like thiamine is more heat stable during heat treatment (Erkan et al., 2010; Ersoy & Ozeren 2009) and fat-soluble vitamins are less heat-labile than water soluble vitamins especially B group of vitamins. The fishes were cooked in pressurized method. In general, it contains high content of the protein than without cooking meat. There is a no difference in total fat and omega-3 fatty acids content between raw and processed freshwater P. pangasius and marine fish, H. macrura. The marine fish are better in source of protein, fat and omega-3 content, while processed fish shows comparable amount of protein, fat and omega-3 fatty acids content with raw fish. The consumers to prepare a healthy menu in order to retain the protein and omega-3 fatty acids content of fish, The fish meat was cooked by different ways before consumption. There are different Cooking methods used which (boiling, baking, roasting, frying, and grilling) can be both beneficial and detrimental to nutrient content of the foods. On the time of cooking of meat, reaction takes place in changing the biochemical and physical content of portion of fillet, it will either improve or impact the food nutritional quality. The cooking was done to improve the hygienic quality of the food by inactivation or destroyed of pathogenic

## 1.INTRODUCTION

Pangasius sp. are commonly found in the south-east Asian region. It belongs to pangasiidae family. The most common variety of cultured fish is Pangasianodon hypophthalmus. This fish species is also called as Sutchi catfish, striped catfish, or Tra fish. Among all the freshwater species, Pangasius catfish is the world's fastest-growing species in aquaculture. Pangasius sp. are popularly traded as skinless and boneless fillets worldwide along with portions, steaks, fillets, and also as value-added products. Pangasius sp. attains weight of 1.2 to 1.3 kg rapidly within six months and harvested after eight months. There is a tremendous demand for fish-based products, especially value-added products such as ready-to-eat "convenience" products. In general, the purpose of cooking was to improve the taste, flavor and reduction (or) inactivation of pathogenic microorganism. Since fish compositions are more sensitive to heat, light and oxygen, These combined cause leads to loss of nutrition during cooking. During cooking of fish fillets, there is a change in proximate, minerals,

microorganisms and their promoted digestibility and bio-availability of nutrient in the digestive tract. It also improves the sensory quality of food by forming aroma compounds, attractive color, crust, and texture. The heat treatment will affect both chemical and biological function of some protein. On the time of direct heat treatment on portion of fillet, moisture will get evaporated to which affect the chemical composition of portion of fillets. There is no modification of amino acid profile due to cooking methods. The composition of fish fillets are affected by (feeding, sex, biological stage, season). It will affect the total fat, protein, and ashes contents and fatty acids composition of fish. The content of thermo-sensitive compounds, fat-soluble vitamins, or PUFAs are often reduced and the nutritive value of fish can be affected by cooking methods. The concentration of  $\omega 3$  fatty acids in rainbow trout, vendace, and pike flesh increased due to the cooking process (boiling, baking, microwave oven, and frying with sunflower and rapeseed oils). Drying in a hot air oven at 180°C for 90 min and microwave drying at 400 W for 8 min of grass carp resulted in a significant increase of protein content but reduced fat content. Changes in proximate chemical composition were more prominent in fried silver catfish. Heat treatment (boiling at 85-90°C during 10-15 min and roasting under 250-280°C during 15-20 min) will not decrease content of EPA and DHA in humpback salmon. García-Arias et al. have also reported that frying shows the highest water loss, fat gain, and affect the fatty acid composition of sardine. Which methods are suitable from a nutritional point of view, it is essential to recommend, the best cooking method preserving nutritional and healthy attributes of fish muscles. These recommendations should be based on the examine and used in different cooking or heat treatment, it will affect an essential nutrition composition and it will give indicators of the quality attributes of fish muscle. Such as Protein, fat, and ashes are the essential nutritional fish muscle compounds which determine the energetic and nutritional values of the muscle. The content of unsaturated fatty acids could be retained as a healthy indicator of fresh and processed fish muscles. The chemical attributes of fresh fish species and the effects of different cooking methods on their chemical compositions is essential to enhance the good food behavior of consumers. The appropriate cooking methods should be done in preserving nutritional and

healthy attributes of fish muscles. The present study aims to determine the influence of 2 cooking methods (steam-cooking, and microwave cooking) on the proximate chemical composition, fatty acid profile, ash contents, and metal contents of red mullet fillets.

## 2. MATERIALS AND METHODS

### 2.1. Materials

*Pangasius hypophthalmus* were collected from Madurai AM fish farm and fish markets. The collected fishes were kept in insulated iceboxes. Insulated icebox prevents dehydration, and temperature fluctuation thus delays the spoilage of fish. Further, it is easy to handle. Flake ice produced by flake ice machine was used during fish transportation and processing purpose. Size of the ice for 2-3 cm level were produced to kept into the box and fish were spread on ice layer and then carried out further steps.

### 2.2. Method

#### 2.2.1. Preparation of dressed meat

The raw *Pangasius* sp. was collected from the market and washed with water. If any foreign material adhered to the outer surface, it was removed. Weight of the cleaned fish sp. was noted down. Removal of fins, head, evisceration was carried out and further washed in clean water. The weight of dressed meat was noted down.

#### 2.2.2. Cooking of pangasius fillets and pasta preparation

- Raw meat (*pangasius* sp)
- Dressed meat
- Washed with water
- Microwave oven cooking (110°C for 6 minutes)
- Steam cooking (98°C for 15 minutes)
- Reduced content of SFA and MUFA

#### 2.2.2. Sampling procedure

Randomly, samples were chosen and studied at different time and temperature was used to remove fat from the fillets. The time and temperature which gave better removal of fat is suitable for standardization of time and temperature. Samples were collected from raw fillets, microwave cooked meat, steam meat and analysis of fatty acid composition. The focus of study is mainly on saturated and mono-unsaturated fatty acid from defatted fish meats.

### I. Microwave oven cooking

The domestic microwave oven was used for cooking of Pangasius fillets. Fillets were kept in the oven to reach the core temperature of 110°C for 6minutes.

### II Steam cooking

The steam cooker was used for this cooking method. The meat was kept in the cooker to reach the core temperature of 98±2°C for 15 minutes.

#### 2.2.3. Fatty acid composition analysed by Gas chromatography

Fatty acids are very important components of lipid content. GC is most common method in analysing fatty acid composition. The fatty acid is a complex structure and it contains many components of fatty acid such as acylglycerols, cholesterol esters, waxes and glycosphingolipids. It is extracted by process of saponification hydrolysis. It is done by alkaline medium AOAC,1990. The FAMES are extracted by use of the methanol and boron trifluoride. Extraction and methylation are done by folch method are used to obtained the lipid components from the ten gram of fish samples. Esterification was done, take 250g lipid fraction it is dissolved in toluene in the round bottom flask. Then, 4ml sodium hydroxide is added and refluxed for 5-10minutes until droplets of fat disappears. 5ml of methanol is added and then refluxed for another 1min. the content is cooled and then 15ml of saturated sodium chloride solution is added. Then, 5ml of hexane is added and shaken well and then remove the upper layer hexane layer. Repeat the extraction with hexane twice. it is combined with the hexane layer and evaporated to dryness in a rotary flask evaporator set at 55-60°C. The methyl esters in 1ml of HPLC grade hexane for injection in GC. The column at 210°C for 30minutes. tTen, inject 0.5ml of standard FAMES mixture onto the GC. Then, it is started and the separation of FAMES takes 45min. Next, inject 0.5ml of sample FAMES. Identify the individual fatty acid in the sample by comparing the retention time of the individual fatty acid in the standard mixture. Calculate area unit value expressed to percentage of the fatty acid of total lipids is done.

#### 2.2.4. Proximate composition analysis

The protein, fat, ash, water and carbohydrate content were analyzed by AOAC methods (2000). 1 gram of wet sample needed for analysis of protein content, 4.5-5 gram of samples are used to estimate of the fat

content, 2 gram of dry sample are used to estimate of the ash and 10 gram of wet samples are used to analysis the moisture content. The protein content of raw fillet, cooked meat and pasta product was analyzed by the Kjeldahl method (1883). The digester temperature was followed at 300 to 400°C and distillation process was run for 8 minutes. The fat content of raw fillet, cooked meat and pasta product was analysed by Folch methods (1957). The moisture content of raw meat, cooked meat and pasta products were analysed by AOAC methods (2000). The samples were kept in hot air oven for 12 hours at 100°C. Ash content of raw meat, cooked meat and pasta products were analysed by AOAC methods (2000). The samples were kept in muffle furnace for 24 hours at 550°C. The carbohydrate content of raw meat, cooked meat and pasta products were analysed by different methods. The chemicals used for estimating the protein includes sulfuric acid, digestion mixture: copper sulfate-0.1g and potassium sulfate-2.5 g for each sample, sodium hydroxide, 40%, boric acid,4%, mixed indicator: methyl red-0.16g and bromocresol green-80 mg in 100ml of 95% ethanol and standard sulfuric acid 0.1N. Fat composition was analyzed by chemicals includes chloroform and methanol (2:1) and potassium chloride(KCL), 0.74%. and another method of soxhlet used in 80 ml of petroleum ether is used to estimate the fat composition. Crude fiber is estimated by chemicals includes sulfuric acid 1.25% and sodium hydroxide 1.25% and petroleum ether 40-60°C.

#### 2.2.5. Sensory evaluation

Sensory evaluation of pasta product was carried out by trained and untrained panelists using 9 points hedonic scale according to (Yousef et al., 2003; prabhasankar et al., 2009a).

#### 2.3. Statistical analysis

The SPSS 19 (IBM, 2010) statistical package was used for analysis of experimental results. The results were produced in the mean standard deviation.

### 3. RESULT AND DISCUSSION

3.1. Different between microwave and steam heat treatment methods remove the fats from Pangasius meat convert to ragi enriched pasta

There is different between microwave and steam heat treatment to remove fats from fish fillets. In microwave heating process, the heating is done by exposing the fish fillets to radiation of electromagnetic wave come from electromagnetic spectrum with range of wave length 0.3-300GHz (Ahmed & Ramaswamy, 2007). This type of instrument is recently used for widely application of food products in both domestic as well as industrial process (Baffler 1993; Cerretani et al., 2009). During heating of food in microwave oven by interaction of electromagnetic field with chemical constitutes of foods, this interaction raise it is comprising molecular fraction and excitation generated if need for cooking purpose (Inchingolo et al., 2013; Mishra et al., 2012; Domiszewski et al., 2011). In general, fish muscle protein was highly sensitive to microwave oven and texture of meat become it very dry, hard and rubbery appearance due to heated with elevated temperature (Mizrahi 2012). Microwave cooking of meat in higher temperature results in loss of nutrition from fish fillets (Shimi 1992). Cook-chilled products will vary in wide range of 6 – 42 days depending upon the heat treatment (Ahmed & Ramaswamy 2007). The raw fish fillets contain high fat content, but after heat treatments fat content reduced due to production of primary and secondary oxidative products. (Regulska-Iliow & Iliow 2002). Mineral composition was very important for health life. Steaming method provides desirable for sensory property and loss of minimum content of nutrients and also plays a important role in destruction of microorganism. At the time of cooking of fish meat containing water, the physical and chemical properties will get changed. Digestibility of fish will get increased due to denaturation of proteins and PUFAs content get reduced in during heat treatment (Raj et al., 2008; Asmah et al., 2014). (Nurhan, 2007) observed that the cooking process affects the amino acid content. The various fish fatty acid profile can be affected by various cooking methods it was reported given by (Nurhan 2007; Weber et al., 2008; Larsen et al., 2010; Koubaa et al., 2012; Sengor et al., 2013; Asmah et al., 2014; Neff et al., 2014). This type of heat treatment reduced the lipid content and retain vitamins C, and preserve the colour and texture of various food products (Idrus & Yang, 2012). The cooking will reduce the amount of amino acids and PUFAs in the fish meat and also used in determination the super-heated steam cooking are influence into fish

nutritional composition which has include in proximate composition, fatty acid and amino acid composition.

The present research proposed to defatted from Pangasius meat and convert to value added products of ragi enriched with fish pasta and to study the proximate composition based on cooking methods and sensory evaluation differentiation based on cooking methods and different concentration of fish flour inclusion. Proximate composition varies with different between microwave and steam heat treatment methods. After heat treatment, the moisture content will get decreased from raw meat to cooked meat the microwave cooked meat is given best reduction of moisture content from Pangasius meat. protein, fat and ash content will get increased after heat treatment. High amount of protein present in the steam cooked meat and high content of fat has been reduce in the cooked meat and ash content was increased by steam cooked methods. Fatty acid composition of raw meat contains more saturated and mono-unsaturated and less poly-unsaturated fatty acids. After heat treatment SFA and MUFA has been decreased but PUFA has been increased. Microwave cooking method shows best reduction of SFA and MUFA and PUFA was increased by steam cooking method. After heat treatment by steaming methods shows increased mineral composition Which is the best methods to promoted to development of ragi pasta products.

#### 1. Proximate composition of pasta products and sensory evaluation

Pasta added with fish flour reduces the moisture content and carbohydrate and increases the lipid, protein and ash content of fish pasta. Addition of tilapia flour to fish results in the lower amount of carbohydrate and higher amount of protein, fat and lipid content (Monterio MLG, et.al., 2014). This results stated that the content of ash, fat, protein content was increased and moisture and carbohydrate decreased due to addition of fish flour with pasta products compared with control pasta (Hong SR, Yoo B, 2012, USDA-2016). There are different concentration of fish flour which is added with pasta products. There are four concentrate with which is best nutritional content was prepared suitable for consumer performance. The sensory assessment of pasta products samples was carried out by trained and untrained panels and the hedonic scale were used. It

has 1–9 rating scale starting from dislike extremely to like extremely (Yousef et al. 2003; Prabhasankar et al. 2009a). Further, four sensory characteristics were included, color, aroma, flavor and aftertaste and have maximum score point of 10. Total or overall acceptability of sensory attributes at  $10 \times 4 = 40$ . It could be accepted for both intensity, hedonic and overall acceptability (Lee et al.1991). Microwave cooked fish with wheat pasta with which Sample-I has moisture-4.40%, protein-14.31%, fat-0.25%, ash-0.56% and carbohydrate-80.46%. Sample-II has moisture-4.03%, protein-15.87%, fat-0.57%, ash-0.87% and carbohydrate-78.64%. Sample-III has moisture-3.80%, protein-16.97%, fat-1.07%, ash-0.94% and carbohydrate-77.54%. Sample-IV has moisture-3.04%, protein-17.66%, fat-1.17%, ash-1.12% and carbohydrate-77.00%. Sample-V has moisture-2.86%, protein-18.77%, fat-1.36%, ash-1.24% and carbohydrate-75.75%. Sensory evaluation has been studied about the fish which is added with wheat pasta. There are five different samples and average score Sample-I 8.87%, Sample-II 8.88%, Sample-III 8.89%, Sample-IV 8.91% and Sample-V 8.93%. Steam cooked with pasta with which Sample-I has an amount of moisture-4.40%, protein-14.31%, fat-0.25%, ash-0.56% and carbohydrate-80.46%. Sample-II has moisture-4.08%, protein-15.97%, fat-0.65% and carbohydrate-78.58%, ash-0.75%. Sample-III has moisture-3.87%, protein-16.75%, fat-0.88%, ash-0.91% and carbohydrate-77.67%. Sample-IV has moisture-3.46%, protein-18.86%, fat-1.57, ash-1.06% and carbohydrate-76.62%. Sample-V has moisture-3.05%, protein-16.32%, fat-, ash-0.35% and carbohydrate-74.82%. Sensory evaluation has been studied about the fish added with ragi pasta. There are five samples and the added score averages are Sample-I 8.67%, Sample-II 8.70%, Sample-III 8.70%, Sample-IV 8.86% and Sample-V 8.91%.

Table1 Proximate composition of pasta products

Microwa ve cooked Sample	Moisture %	Protein %	Fat %	Ash %	Carbohydrate %
1.	4.40	14.31	0.25	0.56	80.46
2.	3.93	15.68	0.66	0.86	78.86
3.	3.62	16.74	1.06	1.02	77.54
4.	3.04	17.66	1.17	1.12	77.00
5.	2.86	18.77	1.36	1.29	75.75

Steam cooked	Moisture %	Protein %	Fat %	Ash %	Carbohydrate %

Sample					
1.	4.40	14.31	0.25	0.56	80.46
2.	4.03	15.97	0.65	0.75	78.58
3.	3.84	16.75	0.88	0.91	77.67
4.	3.46	17.67	1.17	1.06	76.62
5.	3.05	18.86	1.57	1.35	74.82

#### 4.CONCLUSION

There is different between heat treatment methods such as microwave cooked and steam cooked fish with wheat enriched with pasta. Main purpose of these study is to differentiate among pasta products and it is based on proximate and sensory score value. Overall there is a good score for steam cooked meat with wheat pasta due to no loss in nutrition during cooking the meat. Steam meat contains more nutrient compared with microwave cooked meat. The steam cooked meat with pasta has been selected as the best among three pasta products.

#### REFERENCE

- [1] Jeyakumari, A., Janarthan, G., Chouksey, M.K. and Venkateshwarlu, G., 2016. Effect of fish oil encapsulates incorporation on the physico-chemical and sensory properties of cookies. *Journal of food science and technology*, 53(1), pp.856-863.
- [2] Long, N.V., Yang, Y., Thi, C.M., Van Minh, N., Cao, Y. and Nogami, M., 2013. The development of mixture, alloy, and core-shell nanocatalysts with nanomaterial supports for energy conversion in low-temperature fuel cells. *Nano Energy*, 2(5), pp.636-676.
- [3] Severi, S., Bedogni, G., Manzieri, A.M., Poli, M. and Battistini, N., 1997. Effects of cooking and storage methods on the micronutrient content of foods. *European Journal of Cancer Prevention*, 6, S21-S24.
- [4] Garcia-Arias, M.T., Pontes, E.Á., Garcia-Linares, M.C., Garcia-Fernandez, M.C. and Sanchez-Muniz, F.J., 2003. Cooking–freezing–reheating (CFR) of sardine (*Sardina pilchardus*) fillets.
- [5] Effect of different cooking and reheating procedures on the proximate and fatty acid compositions. *Food chemistry*, 83(3), 349-356.
- [6] Food and Agriculture Organization of the United Nations. *Fats and fatty acids in Human Nutrition*.

- Report of an expert consultation 2010, November 10–14, 2008, Geneva.
- [7] Larsen, R.; Eilertsen, K.E.; Elvevoll, E.O. Health Benefits of Marine Foods and Ingredients. *Biotechnology Advances* 2011, 29, 508–518.
- [8] Turan, H.; Sonmez, G.; Kaya, Y. Fatty Acid Profile and Proximate Composition of the Thornback Ray (*Raja Clavata*, L. 1758) from the Sinop Coast in the Black Sea. *Journal of Fisheries Sciences* 2007, (T1)1, 97–103.
- [10] Testi, S.; Bonaldo, A.; Gatta, P.P.; Badiani, A. Nutritional Traits of Dorsal and Ventral Fillets from Three Farmed Fish Species. *Food Chemistry* 2006, 98, 104–111.
- [11] Unusan, N. Change in Proximate, Amino Acid, and Fatty Acid Contents in Muscle Tissue of Rainbow Trout (*Oncorhynchus Mykiss*) after Cooking. *International Journal of Food Science and Technology* 2007, 42, 1087–1093.
- [12] Kalogeropoulos, N.; Andrikopoulos, N.; Hassapidou, M. Dietary Evaluation of Mediterranean Fish and Molluscs PanFried in Virgin Olive Oil. *Journal of the Science of Food and Agriculture* 2004, 84, 1750–1758.
- [13] Marques, A.; Teixeira, B.; Barrento, S.; Anacleto, P.; Carvalho, M.L.; Nunes, M.L. Chemical Composition of Atlantic Spider Crab *Majabrachy Dactyla*: Human Health Implications. *Journal of Food Composition and Analysis* 2010, 23, 230–237.
- [14] Badiani, A.; Stipa, S.; Bitossi, F.; Pirini, M.; Bonaldo, A.; Gatta, P.P.; Rotolo, M.; Testi, S. True Retention of Nutrients upon Household Cooking of Farmed Portion-Size European Sea Bass (*Dicentrarchus Labrax* L). *LWT–Food Science and Technology* 2013, 50, 72–77.
- [15] Immanuel, G.; Sathasivan, S.; Selva Shankar, V.; Punitha Peter, M.G.; Palavesam, A. Processing and Characterization of Low Cost Balistid Fish *Sufflamen Capistratus* Liver Oil for Edible Purpose. *Food Chemistry* 2009, 115, 430–435.
- [16] Erkan, N.; Selçuk, A.; Ozden, O. Amino Acid and Vitamin Composition of Raw and Cooked Horse Mackerel. *Food Analytical Methods* 2010, (E1) 3, 269–275.
- [17] Ersoy, B.; Ozeren, A. The Effect of Cooking Methods on Mineral and Vitamin Contents of African Catfish. *Food Chemistry* 2009, 115, 419–422.
- [18] Taskaya, L.; Chen Chen, Y.; Beamer, S.; Tou, J.C.; Jaczynski, J. Compositional Characteristics of Materials Recovered from Whole Guttled Silver Carp (*Hypophthalmichthys Molitrix*) Using Iso-Electric Solubilization/Precipitation. *Journal of Agricultural and Food Chemistry* 2009, 57, 4259–4266.
- [19] Badiani, A., Stipa, S., Bitossi, F., Pirini, M., Bonaldo, A., Gatta, P.P., Rotolo, M. and Testi, S., 2013. True retention of nutrients upon household cooking of farmed portion-size European sea bass (*Dicentrarchus labrax* L.). *LWT-Food science and Technology*, 50(1), 72-77.
- [20] AOAC, 2000 AOAC Official methods of analysis.
- [21] Kjeldahl, C., 1883. A new method for the determination of nitrogen in organic matter. *Z Anal Chem*, 22, p.366.
- [22] Folch, J., Lees, M. and Stanley, G.S., 1957. A simple method for the isolation and purification of total lipides from animal tissues. *Journal of biological chemistry*, 226(1), pp.497-509.
- [23] Cerretani, L., Bendini, A., Rodriguez-Estrada, M.T., Vittadini, E. and Chiavaro, E., 2009. Microwave heating of different commercial categories of olive oil: Part I. Effect on chemical oxidative stability indices and phenolic compounds. *Food Chemistry*, 115(4), 1381-1388.
- [24] Domiszewski, Z., Bienkiewicz, G. and Plust, D., 2011. Effects of different heat treatments on lipid quality of striped catfish (*Pangasius hypophthalmus*). *Acta Scientiarum Polonorum Technologia Alimentaria*, 10(3), 359-373.
- [25] Inchingolo, R., Cardenia, V. and Rodriguez-Estrada, M.T., 2013. The effects of microwave heating on edible oils and lipid-containing food. *Lipid Technology*, 25(3), 59-61.
- [26] Mishra, R., Sharma, H.K., Sarkar, B.C. and Singh, C., 2012. Thermal oxidation of rice bran oil during oven test and microwave heating. *Journal of food science and technology*, 49(2), 221-227.
- [27] Ahmed, J., Ramaswamy, H.S. and Raghavan, V.G., 2007a. Dielectric properties of butter in the MW frequency range as affected by salt and temperature. *Journal of Food Engineering*, 82(3), 351-358.

- [28] El-Shimi, N.M., 1992. Influence of microwave and conventional cooking and reheating on sensory and chemical characteristics of roast beef. *Food chemistry*, 45(1), 11-14.
- [29] Regulska-Ilow, B. and Ilow, R., 2002a. Comparison of the effects of microwave cooking and conventional cooking methods on the composition of fatty acids and fat quality indicators in herring. *Food/Nahrung*, 46(6), 383-388.
- [30] Raj, A.J.A., Haniffa, M.A., Seetharaman, S. and Appelbaum, S., 2008. Utilization of various dietary carbohydrate levels by the freshwater catfish *Mystus montanus* (Jerdon). *Turkish Journal of fisheries and aquatic sciences*, 8(1), 31-35.
- [31] Asmah, R., Sumaiyah, S.S. and Nurul, S.R., 2014a. Comparison of protein, total fat, and omega-3 fatty acids content in yellowtail catfish (*Pangasius pangasius*) and long tail shad (*Hilsa (clupea) macrura*) in raw and pressurized fish. *International Food Research Journal*, 21(6), 2147.
- [32] Nurhan, U., 2007a. Change in proximate, amino acid and fatty acid contents in muscle tissue of rainbow trout (*Oncorhynchus mykiss*) after cooking. *International journal of food science & technology*, 42(9), 1087-1093.
- [33] Weber, J., Bochi, V.C., Ribeiro, C.P., Victório, A.D.M. and Emanuelli, T., 2008a. Effect of different cooking methods on the oxidation, proximate and fatty acid composition of silver catfish (*Rhamdia quelen*) fillets. *Food Chemistry*, 106(1), 140-146.
- [34] Larsen, D., Quek, S.Y. and Eyres, L., 2010. Effect of cooking method on the fatty acid profile of New Zealand King Salmon (*Oncorhynchus tshawytscha*). *Food Chemistry*, 119(2), 785-790.
- [35] Koubaa, A., Mihoubi, N.B., Abdelmouleh, A. and Bouain, A., 2012. Comparison of the effects of four cooking methods on fatty acid profiles and nutritional composition of red mullet (*Mullus barbatus*) muscle. *Food Science and Biotechnology*, 21(5), 1243-1250.
- [36] Şengor, G.F.U., Alakavuk, D.U. and Tosun, Ş.Y., 2013. Effect of cooking methods on proximate composition, fatty acid composition, and cholesterol content of Atlantic Salmon (*Salmo salar*). *Journal of aquatic food product technology*, 22(2), 160-167.
- [37] Neff, M.R., Bhavsar, S.P., Braekevelt, E. and Arts, M.T., 2014. Effects of different cooking methods on fatty acid profiles in four freshwater fishes from the Laurentian Great Lakes region. *Food chemistry*, 164, 544-550.
- [38] Asmah, R., Sumaiyah, S.S. and Nurul, S.R., 2014b. Comparison of protein, total fat, and omega-3 fatty acids content in yellowtail catfish (*Pangasius pangasius*) and long tail shad (*Hilsa (clupea) macrura*) in raw and pressurized fish. *International Food Research Journal*, 21(6), 2147.
- [39] Ezhil, C., 2010. Superheated steam drying of foods—A review. *World journal of dairy and food sciences*, 5(2), 214-217.
- [40] Idrus, N.F.M. and Yang, T.A., 2012. Comparison between roasting by superheated steam and by convection on changes in colour, texture and microstructure of peanut (*Arachis hypogaea*). *Food Science and Technology Research*, 18(4), 515-524.
- [41] Anbudhasan, P., Asvini, G., Surendraraj, A., Ramasamy, D. and Sivakumar, T., 2014. Development of functional pasta enriched with omega-3 fatty acids. *Fish Technol*, 51, 242-246.
- [42] Prabhasankar, P., Ganesan, P., Bhaskar, N., Hirose, A., Stephen, N., Gowda, L.R., Hosokawa, M. and Miyashita, K., 2009. Edible Japanese seaweed, wakame (*Undaria pinnatifida*) as an ingredient in pasta: Chemical, functional and structural evaluation. *Food Chemistry*, 115(2), 501-508.
- [43] Yousif, A.M., Cranston, P. and Deeth, H.C., 2003. Incorporation of bovine dry blood plasma into biscuit flour for the production of pasta. *LWT-Food Science and Technology*, 36(3), 295-302..
- [44] Lee, C. C., Love, J. A., & Johnson, L. A. (1993). Sensory and physical properties of cakes with bovine plasma products substituted for egg. *Cereal Chemistry*, 70, 18–21.