

A Study on Association Factors of BMI and Prevalence of Obesity in Management Students from CSI Institute of Legal Studies Cheruvarakonam

Mrs.Jijitha A Joseph¹, Dr. P.Arul deva Paul², Mrs. A.Amala Shalni³

^{1,2,3}*Department of Health and Wellness, Management CSI Institute of Legal Studies Cheruvarakonam, India*

Abstract - Obesity was a vexing problem in the developed economies country. For developing countries like India, morbid obesity has not yet become a public health priority. Successive cross-sectional representative surveys method was used. For this study sixty-five students (N= 64) were selected from Management department students from CSI Institute of Legal Studies Cheruvarakonam. The subjects are Management department students from assorted department in CSI Institute of Legal Studies Cheruvarakonam. Following procedure are used to take the data for BMI from the respective sampling weights of the subject are taken by using of digital weighing machine in kg and height was taken by using of stadiometer scale in cm. The data pertaining BMI were statistically analyses by the use of descriptive frequency percentile method .The obtained BMI association factors (Weight ,Height and Age) are tested for significance at 0.05 level of confidence by Crosstabs Chi-Square Tests. The result Among the Following N=65 subject shows 45% of students comes under the weight range from 56-65kg,29.7%students are 66-75kg,14.1% are <75kg and also 10.9%are >55kg.In Height of the subject are 56.2% of students comes under the height of 166-172cm,28.1% are >165cm,14.1% are 176-185cm and 1.6% of students comes under <185cm. According to the age57%are>27years old, 43.8% are 27-29yeras old ,and 1.6% are <29. The results of Chi-square test shows that there is association between weight and BMI of sample respondent but there is no association were found on sample respondents age and height. From this result it is concluded that increasing the amount of weight causes obesity in the human beings and produced health related disease in nature.

INTRODUCTION

Obesity was a vexing problem in the developed economies country. For developing countries like India, morbid obesity has not yet become a public

health priority. Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have a negative effect on health, leading to reduced life expectancy increased health problems. Obesity increases the likelihood of various diseases, particularly heart disease, type 2 diabetes, obstructive sleep apnea, certain types of cancer, and osteoarthritis. Obesity is most commonly caused by a combination of excessive food energy intake, lack of physical activity, and genetic susceptibility, although a few cases are caused primarily by genes, endocrine disorders, medications, or psychiatric illness. On average, obese people have greater energy expenditure than their thin counterparts due to the energy required to maintain an increased body mass. Dieting and exercising are the main treatments for obesity. Obesity is a leading preventable cause of death worldwide, with increasing rates in adults and children. Authorities view it as one of the most serious public health problems of the 21st century. Based on Third National Family Health Survey. International Institute for Population Sciences, 2005–2006, Mumbai in India, states which topped the list of rates of obesity are Punjab 30.3% males, 37.5% females, Kerala 24.3% males,34% females and Goa 20.8% males,27% females.

DESIGN

Successive cross-sectional representative surveys study, including the association of weight, height and age with BMI. Body mass index was calculated from measured weight and height. Overweight and obesity were defined as follows: morbidly Obese (BMI > 40); obese (BMI 30-40), overweight (BMI 25-30), Normal(BMI 20-25), and Under Weight (BMI < 20). BMI = weight (kg)/ height (cm)

METHODOLOGY & RESULTS OF THE STUDY

For this study sixty-five students (N= 64) were selected from Management department students from CSI Institute of Legal Studies Cheruvarakonam. The subjects are Management department students from assorted department in CSI Institute of Legal Studies Cheruvarakonam. Following procedure are use to take the data for BMI from the respective sampling weights of the subject are taken by using of digital weighing machine in kg and height was taken by using of stadiometer scale in cm. The data pertaining BMI were statistically analyses by the use of descriptive frequency percentile method .The obtained BMI association factors (Weight ,Height and Age) are tested for significance at 0.05 level of confidence by Crosstabs Chi-Square Tests.

Table 1 – Descriptive Statistics of Sample Respondents

S.NO		Weight	Height	Age	BMI
1.	N	64	64	64	64
2.	Mean	2.48	1.91	1.48	2.21
3.	S.D	0.871	0.692	0.535	0.643
4.	Variance	0.760	0.482	0.286	0.414
5.	Range	3	3	2	4

Table 2: Percentile of BMI Association Factors

weight	N	Percentage	Height(cm)	N	Percentage	Age	N	Percentage
Below 55	23	10.9%	Below 165	18	28.1%	Below 27	35	54.7%
56 - 65	29	45.3%	166 - 175	36	56.2%	27 -29	28	43.8%
66 - 75	9	29.7%	176 - 185	9	14.1%	Above 29	1	1.6%
Above 75	9	14.1%	Above 185	1	1.6%	Total	64	100.0%

	Value age	d.f	Sig.	Value Height	d.f	Sig.	Value Weight	d.f	Sig.
Pearson Chi-Square	4.421	6		8.632	9		43.650	9	0.000*
Likelihood Ratio	4.695	6		6.738	9		36.309	9	
Fisher's Exact Test	7.369		0.499	10.772		0.358	31.790		
Linear-by-Linear Association	0.034	1		0.245	1		25.589	1	
N of Valid Cases	64			64			64		

Tot al	64	100.0%	Total	64	100.0%			
--------	----	--------	-------	----	--------	--	--	--

The result Among the Following N=65 subject shows 45% of students comes under the weight range from 56-65kg,29.7%students are 66-75kg,14.1% are <75kg and also 10.9%are >55kg.In Height of the subject are 56.2% of students comes under the height of 166-172cm,28.1% are >165cm,14.1% are 176-185cm and 1.6% of students comes under <185cm. According to the age57%are>27years old, 43.8% are 27-29yeras old and 1.6% are <29.

Table 3: Obesity Range for Management department students

BMI	Frequency	Percent
Below 19	4	6.2
19 - 24	42	65.6
25 - 30	17	26.6
Above 40	1	1.6
Total	64	100.0




Table 4: Results of Chi-Square Tests

In Chi-Square test the significant value of the weight was0.000*so that weight sample respondent to BMI was significant. This was significant at 0.05level of confidence. In age the significant value was 0.499 that higher than the p value so there is no significant difference on age. 0.058 was the significant value of height of the subjects. That value was higher than the p value so there was no significant at 0.05 level of confidence.

CONCLUSION AND FINDS

Based on the observation and findings from this study it is concluded that 1.60% students are in obesity, Where as 6.20% students are in underweight, 26.60 % are in overweight and 65.60% are in normal in management students. The results of Chi-square test shows that there is association between weight and BMI of sample respondent but there is no association were found on sample respondents age and height. From this result it is concluded that increasing the amount of weight causes obesity in the human beings and produced health related disease in nature. In order to obesity prevention is the best ideal method. These who are in overweight obesity and underweight students are suggested to counseled with physical education trainee, practice daily physical activity such as yoga, and maintaining healthy balance diet. BMI normal students are recommended to maintain their BMI in normal as lifelong.

REFERENCE

- [1] Agarwal VK, Basannar DR, Sing RP, Dutt M, Abraham D, Mustafa MS (2006) Coronary risk factors in a rural community. *Indian JPublic Health* 50:19–23
- [2] Blas E, Kurup AS (eds) (2010) Equity, social determinants and publichealth programs. World Health Organization, Geneva
- [3] Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet [Internet]* (0).
- [4] Stevens GA, Singh GM, Lu Y, Danaei G, Lin JK, Finucane MM, et al. National, regional, and global trends in adult overweight and obesity prevalences. *Popul Health*. 2012;10(1):22.
- [5] Kelly T, Yang W, Chen C-S, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes* 2005. 2008 Sep;32(9):1431–7.
- [6] Wang Y, Beydoun MA, Liang L, Caballero B, Kumanyika SK. Will all Americans become overweight or obese? estimating the progression and cost of the US obesity epidemic. *Obes Silver Spring Md*. 2008 Oct;16(10):2323–30.
- [7] Hu FB. *Obesity epidemiology*. Oxford University Press; Oxford; New York: 2008. p. 498.
- [8] Hu FB. Obesity and Mortality: Watch Your Waist, Not Just Your Weight. *Arch Intern Med*. 2007 May 14;167(9):875.
- [9] Alberti KGMM, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the Metabolic Syndrome: A Joint Interim Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation*. 2009 Oct 20;120(16):1640–5.
- [10] Alberti KGM, Zimmet P, Shaw J. The metabolic syndrome—a new worldwide definition. *The Lancet*. 366(9491):1059–62.
- [11] WHO Multicentre Growth Reference Study Group WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Oslo Nor 1992 Suppl*. 2006 Apr;450:76–85.
- [12] De Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ*. 2007 Sep;85(9):660–7.
- [13] Barlow SE, the Expert Committee Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report. *Pediatrics*. 2007 Dec 1;120(Supplement 4):S164–92.
- [14] Flegal KM, Wei R, Ogden CL, Freedman DS, Johnson CL, Curtin LR. Characterizing extreme values of body mass index-for-age by using the 2000 Centers for Disease Control and Prevention growth charts. *Am J Clin Nutr*. 2009 Nov;90(5):1314–20.
- [15] Grummer-Strawn LM, Reinold C, Krebs NF, Centers for Disease Control and Prevention (CDC) Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United States. *MMWR Recomm Rep Morb Mortal Wkly Rep Recomm Rep Cent Dis Control*. 2010 Sep 10;59(RR-9):1–15.