

Fuzzification of Lattice Algebraic Structure

Mr. M.U.makandar

Assistant Professor, PG, MBA Department, KIT's IMER, Kolhapur. Maharashtra

Abstract - In this paper we introduce the notion of fuzzy lattice ordered m normal subgroups and investigated some of its basic properties. In this paper we defined fuzzy lattice ordered m-normal group, arbitrary intersection of fuzzy lattice ordered m-normal groups, m-fuzzy characteristic of fuzzy lattice ordered m group. We also studied the Cartesian product of arbitrary family of fuzzy lattice ordered m-normal groups.

Index Terms - Lattice ordered group, Fuzzy lattice ordered m-group, Fuzzy lattice ordered m normal group, characteristic of fuzzy group, direct product.

1.INTRODUCTION

A fuzzy algebra has become an important branch of research. A. Rosenfeld [9] used the concept of fuzzy set theory due to Zadeh [5]. Since then, the study of fuzzy algebraic substructures is important when viewed from a Lattice theoretic point of view. N. Ajmal and K.V. Thomas [1] initiated such types of study in the year 1994. It was latter independently established by N. Ajmal [1] that the set of all fuzzy normal subgroups of a group constitute a sub lattice of the lattice of all fuzzy subgroups of a given group and is Modular. Nanda [8] proposed the notion of fuzzy lattice using the concept of fuzzy partial ordering. More recently in the notion of set product is discussed in detail and in the lattice theoretical aspects of fuzzy subgroups and fuzzy normal subgroups are explored. G.S.V. Satya Saibaba [3] initiate the study of L-fuzzy lattice ordered groups and introducing the notice of L-fuzzy sub l- groups. J.A. Goguen [4] replaced the valuation set [0,1] by means of a complete lattice in an attempt to make a generalized study of fuzzy set theory by studying L-fuzzy sets. A Solairaju and R. Nagarajan [11] introduced the concept of lattice valued Q-fuzzy sub-modules over near rings with respect to T-norms. DrM.Marudai & V. Rajendran [6] modified the definition of fuzzy lattice and introduce the notion of fuzzy lattice of groups and investigated some of its basic properties. Gu [12] introduced concept of fuzzy groups with operator. Then S.

Subramanian, R Nagarajan & Chellappa [10] extended the concept to m fuzzy groups with operator. In this paper we introduce the notion of fuzzy lattice ordered m normal subgroups and investigated some of its basic properties. In this paper we defined fuzzy lattice ordered m-normal group, arbitrary intersection of fuzzy lattice ordered m-normal groups, m-fuzzy characteristic of fuzzy lattice ordered m group. We also studied the Cartesian product of arbitrary family of fuzzy lattice ordered m-normal groups.

2.PRELIMINARIES

Definition 2.1: Fuzzy group- Let $\mu: X \rightarrow [0, 1]$ be a fuzzy set & $G \in \mathfrak{p}(X) = \text{Set of all fuzzy sets on } X$. A fuzzy set μ on G is called a fuzzy group if i) $\mu(x y) \geq \min \{\mu(x), \mu(y)\}$ ii) $\mu(x^{-1}) \geq \mu(x)$, for all $x, y \in G$.

Definition 2.2: Lattice ordered group - A lattice ordered group is a system (G, \leq) if i) (G, \cdot) is a group ii) (G, \leq) is a lattice . iii) $x \leq y$ implies $a x b \leq a y b$ (compatibility) For $a, b, x, y \in G$

Definition 2.3: Fuzzy lattice ordered group- Let $\mu: X \rightarrow [0, 1]$ be a fuzzy set & G is a lattice ordered set, $G \in \mathfrak{p}(X)$. A function μ on G is said to be a fuzzy lattice ordered group if i) $\mu(x y) \geq \min \{\mu(x), \mu(y)\}$ ii) $\mu(x^{-1}) \geq \mu(x)$ for all $x, y \in G$

Definition 2.4: M group- Let G be a group, M be any set if i) $m x \in G$. ii) $m(x y) = (m x) y = x m y$ for all $x, y \in G, m \in M$. Then G is called a m group.

Definition 2.5: Fuzzy m group- Let $\mu: X \rightarrow [0, 1]$ be a fuzzy set & G be a M group G . A fuzzy set on $G, G \in \mathfrak{p}(X)$ is called a fuzzy m group if i) $\mu(m(x y)) \geq \min \{\mu(m x), \mu(m y)\}$ ii) $\mu(m x^{-1}) \geq \mu(m x)$ for all $x, y \in G, m \in M$

Definition 2.6: A fuzzy lattice ordered m-group- $\mu: X$ to $[0, 1], G \in \mathfrak{p}(X), M \subset X$.

A function μ on G is said to be a fuzzy lattice ordered m-group if

1. (G, \cdot) is a M-group.
2. (G, \cdot, \leq) is a lattice ordered group.
3. $\mu(m(x y)) \geq \min\{\mu(m x), \mu(m y)\}$
4. $\mu((m x)^{-1}) \geq \mu(m x)$
5. $\mu(m x \vee m y) \geq \min\{\mu(m x), \mu(m y)\}$
6. $\mu(m x \wedge m y) \geq \min\{\mu(m x), \mu(m y)\}$ For all $x, y \in G$

Definition 2.7: Fuzzy lattice ordered m-normal subgroup- A fuzzy lattice ordered m-group is said to be a fuzzy lattice ordered m-normal subgroup if $\mu((m x)(m y)) = \mu((m y)(m x))$ for all $x, y \in G, m \in M$

Definition 2.8: M-fuzzy characteristic of lattice ordered m-group

A fuzzy lattice ordered m group A is said to be a m-fuzzy characteristic of lattice ordered m-group if $\mu_A(m x) = \mu_A(f(m x))$ for all $x \in G, m \in M, f \in \text{Aut}(G)$

3.BASIC PROPERTIES OF FUZZY LATTICE ORDERED M-NORMAL GROUP

Proposition 3.1: Intersection of two fuzzy lattice ordered m-normal subgroup is again fuzzy lattice ordered m-normal subgroup.

Proof- Let A and B be two fuzzy lattice ordered m-normal groups on G .

$$\begin{aligned} \mu_{A \cap B}((m x)(m y)) &= \mu_A((m x)(m y)) \wedge \mu_B((m x)(m y)) \\ &= \mu_A((m y)(m x)) \wedge \mu_B((m y)(m x)) \\ &= \mu_{A \cap B}((m y)(m x)) \end{aligned}$$

Proposition 3.2: If $\{A_i\}$ is a family of fuzzy lattice ordered m-normal subgroups of G then $\cap A_i$ is a fuzzy lattice ordered m-normal group of G where $\cap A_i = \{m x, \wedge \mu_{A_i}(m x) / x \in G, m \in M\}$

Proof- $(\cap \mu_{A_i})((m x)(m y)) = \wedge \mu_{A_i}((m x)(m y))$
 $= \wedge \mu_{A_i}(m y m x)$
 $= (\cap \mu_{A_i})(m y m x)$

Proposition 3.3: If A is a m-fuzzy characteristic of fuzzy lattice ordered m group of G then A is fuzzy lattice ordered m-normal subgroup of G .

Proof- Let A be a m-fuzzy characteristic of fuzzy lattice ordered m group of G .

Consider the map $f: G \rightarrow G$ defined by $f(m x) = m y m x (m y)^{-1}$

Clearly $f \in \text{Aut}(G)$

$$\begin{aligned} \text{Now } \mu_A((m x)(m y)) &= \mu_A(f((m x)(m y))) \\ &= \mu_A(m y m x (m y)(m y)^{-1}) \\ &= \mu_A(m y m x) \end{aligned}$$

Proposition 3.4: A fuzzy lattice ordered m-group is a normal if and only if A is constant on conjugate classes of G .

Proof- Let A be a fuzzy lattice ordered m-normal subgroup of G .

$$\mu_A(m y)^{-1}(m x)(m y) = \mu_A((m y)^{-1} m y m x) = \mu_A(m x)$$

Hence A is constant on conjugate classes of G .

Conversely

$$\begin{aligned} \mu_A(m x m y) &= \mu_A(m x m y m x (m x)^{-1}) = \mu_A(m x (m y m x)(m x)^{-1}) \\ &= \mu_A(m y m x) \end{aligned}$$

Therefore, the fuzzy lattice ordered m-group is normal.

Proposition 3.5: For fuzzy lattice ordered m normal group.

$$\mu_A(m y)^{-1}(m x)(m y) = \mu_A(m y)(m x)(m y)^{-1}$$

Proof- Let A be a fuzzy lattice ordered m-normal subgroup of G .

$$\begin{aligned} \mu_A(m y)^{-1}(m x)(m y) &= \mu_A(m x) \\ &= \mu_A((m x)(m y)(m y)^{-1}) \\ &= \mu_A(m y)(m x)(m y)^{-1} \end{aligned}$$

Applications: Lattice structure has been found to be extremely important in the areas of quantum logic, Ergodic theory, Reynold's operations, Soft Computing, Communication system, Information analysis system, artificial intelligences, and physical sciences.

ACKNOWLEDGEMENT

The author is highly grateful to the referees for their valuable comments and suggestions for improving the paper.

REFERENCES

- [1] Ajmal N and K.V.Thomas, The Lattice of Fuzzy subgroups and fuzzy normal subgroups, Inform. sci. 76 (1994), 1 – 11.
- [2] Birkoff, G: Lattice theory 3rd edn. Amer.Math. Soc. Colloquium pub.25 (1984).

- [3] G.S.V. Satya Saibaba. Fuzzy lattice ordered groups, South east Asian Bulletin of Mathematics 32,749-766 (2008).
- [4] J.A. Goguen: L – Fuzzy Sets, J. Math Anal.Appl. 18, 145-174 (1967).
- [5] L. A.Zadeh : Fuzzy sets, Inform and Control, 8, 338-353 (1965).
- [6] M.Marudai & V. Rajendran: Characterization of Fuzzy Lattices on a Group International Journal of Computer Applications with Respect to T-Norms, 8(8),0975 – 8887 (2010)
- [7] Mordeson and D.S. Malik: Fuzzy Commutative Algebra, World Scientific Publishing Co. Pvt. Ltd.
- [8] Nanda, S: Fuzzy Lattices, Bulletin Calcutta Math. Soc. 81 (1989) 1 – 2.
- [9] Rosenfeld: Fuzzy groups, J. Math. Anal. Appl. 35, 512 – 517 (1971).
- [10] S. Subramanian, R Nagarajan & Chellappa, Structure Properties of M-Fuzzy Groups Applied Mathematical Sciences,6(11),545-552(2012)
- [11] Solairaju and R. Nagarajan: Lattice Valued Q-fuzzy left R – Submodules of Neat Rings with respect to T-Norms, Advances in fuzzy mathematics 4(2), 137 – 145 (2009).
- [12] W.X.Gu. S.Y.Li and D.G.Chen, fuzzy groups with operators, fuzzy sets and system,66 (1994), 363-371.