

Exploring the Role of Serum CRP in Early Assessment of Stroke Severity and Functional Recovery

Zain Ul Bashar Khan¹, Dr.Narottam Pal², and Arlagadda Prasad³, Mukund Panukanti⁴

^{1,3,4}Associate professor, Bhaskar Pharmacy College, Moinabad

² Professor and HOD, Bhaskar Pharmacy College, Moinabad

Abstract—Stroke is a leading cause of death and disability globally. C-reactive protein (CRP), an inflammatory marker, increases after stroke and may predict functional outcomes in acute ischemic stroke (AIS). This study aimed to evaluate CRP's role in assessing stroke severity and short-term outcomes. A prospective observational study was conducted, with a mean patient age of 62.3 years. The majority of patients had moderate stroke (58%), and 16% had minor stroke. CRP levels were significantly related to stroke severity ($p=0.001$). Hypertension, dyslipidemia, and smoking were associated with worse short-term outcomes. At discharge, 4% were independent, while 14% were totally dependent. CRP levels can serve as an independent prognostic factor for stroke severity and functional outcomes.

Index Terms—Acute Ischemic Stroke, CRP, Barthel Index

I. INTRODUCTION

Stroke is one of the leading causes of death and the main causes of disability worldwide. As per WHO, 15 million people suffer from stroke every year.^[1] Worldwide, 70% of strokes and 87% of both stroke-related deaths and disability-adjusted life years occur in low and middle-income countries. Stroke mainly affect individuals at the peak of their productive life. Even though it has got high impact on socio- economic development, this growing crisis has received very little attention till date.^[2]

WHO defines stroke as “rapidly developing clinical symptoms/signs of focal or global loss of brain function, with symptoms lasting more than 24 hrs or leading to death, with no apparent cause other than that of vascular origin”.^[3] New definition of stroke has been proposed by the American Stroke Association “CNS infarction is defined as the brain,

spinal cord, or retinal cell death attributable to ischemia based on pathological, imaging, or other objective evidence of cerebral, spinal cord or retinal focal ischemic injury in a defined vascular distribution or clinical evidence of cerebral, spinal cord, or retinal focal ischemic injury based on symptoms persisting ≥ 24 hours or until death, and other etiologies excluded. Ischemic stroke is an episode of neurological dysfunction caused by focal cerebral, spinal, or retinal infarction”.^[4] When blood flow is quickly restored brain tissue can recover fully and the patient's symptoms are only transient is called Transient Ischemic Attack (TIA).^[5]

Studies shows that C-reactive protein, a peripheral marker of inflammation, can also be used as a marker of generalized atherosclerosis.^[6] Acute stroke may trigger an inflammatory response that leads to increased levels of CRP. CRP has the potential to serve as both a prognostic indicator following vascular events as well as a potential predictor of future vascular events due to the association between inflammation and atherosclerosis. CRP is an acute phase reactant produced not only by the liver but also in vascular smooth muscle cells and adiposities.⁽⁷⁻¹⁰⁾

II. MATERIAL AND METHODS

A. Study design:


The clinical study was performed at secondary care hospital, Andhra Pradesh, India from Jan 2023 to Dec 2023. Laboratory data were collected.

III. CRITERIA FOR PATIENT SELECTION: INCLUSION CRITERIA

- 1.Symptom onset within 24 hours of admission
- 2.Age more than 18 years

- 3. Patients with focal neurological deficits and MRI Brain confirmed acute ischemic stroke
- 4. Patients with new onset stroke with past history /newly detected hypertension, diabetes mellitus and dyslipidaemia are included
- 5. Patients with habits of alcohol intake, smoking and OCPs are included

IV. EXCLUSION CRITERIA



- 1. MRI brain confirmed hemorrhagic stroke
- 2. Symptom onset >24 hours
- 3. Patients with recent ACS, Cerebrovascular events or any infections that affect CRP value

STUDY DESIGN: Prospective observational study
 SAMPLE SIZE: 50 patients who met all the inclusion criteria during the study period

RESULTS AND ANALYSIS

Total 50 cases with cerebrovascular accident due to ischemia between age of 23 to 85 years were studied.

Table 1: Distribution Of the Subjects Based on Age Groups

Age groups (Years)	Frequency	Percent
20 to 40	2	4.0
41 to 50	5	10.0
51 to 60	18	36.0
61 to 70	13	26.0
> 70	12	24.0
Total	50	100.0

In this study youngest of the population was 23 years and oldest was 85 years. Maximum number of cases

were in between 51-60 years with a mean age of 62.3 years




Table 2: Risk Factors Of Ais

RISK FACTORS	SEX		
	F	M	TOTAL
HYPERTENSION	11	28	39
DM	9	27	36
DYSLIPIDEMIA	4	11	15
HEART DISEASES	3	7	10
SMOKING	0	18	18
ALCOHOL	0	11	11
OCPs	0	0	0

Table 3: Stroke Severity Based On Nihss Score

NIHSS	SEX		Total
	Females	Males	
Minor	2	6	8
Moderate	9	20	29
Moderate to severe	3	8	11
Severe	0	2	2
Total	14	36	50

Figure 2: Stroke Severity Based on Nihss

In this study 58% cases were alert, 20% were responding to verbal commands, 18% were pain responsive and 4% were unresponsive. 86% of the patients had weakness of limbs and 70% had cranial nerve involvement. Cerebellar ischemia was seen in 14% cases

When stroke severity classified according to NIHSS, study shows 16% had minor stroke, 58% had moderate stroke, 22% had moderate to severe stroke and 4% had severe stroke. Among these females were more affected with moderate stroke and in other

severity classifications males were more affected than females.

Table 5: Crp with Age Groups

Age groups(years)	CRP CATEGORY	
	NEG	POS
20 - 40	1	1
41-50	3	2
51 -60	13	5
61 -70	7	6
> 70	10	2
Total	34	16

In the study total of 34 cases got negative CRP results and 16 had positive CRP results. The age group 51-60 years had maximum number of cases and negative CRP levels. The age group 61-70 years had maximum number of cases with positive CRP levels.

Table 6: Nihss with Crp

NIHSS	CRP		Total
	NEGATIVE	POSITIVE	
Minor	8	0	8
	23.5%	0.0%	16.0%
Moder- ate	24	5	29
	70.6%	31.2%	58.0%
Moder- ate too severe	2	9	11
	5.9%	56.2%	22.0%
Severe	0	2	2
	0.0%	12.5%	4.0%
Total	34	16	50
	100.0%	100.0%	100.0%

CRP and stroke severity by NIHSS score was significantly related in this study (Chi- square value 23.46, p value=0.001). When CRP is less than 6mg/L, maximum number of cases comes under minor to moderate stroke. When CRP values more than 6mg/L, cases had moderate- severe to severe stroke. This indicates that, the increase in CRP levels were directly proportional to stroke severity.

Dependency of the patients at discharge evaluated by Barthel index. Out of total 50 cases 4% cases were independent at the time of discharge, 28% had mild dependency, 34% had moderate dependency, 20% had severe dependency and 14% were totally depended. In

elderly age group most of the patients had moderate to severe dependency.

In this study 2 cases had mild dependency at admission, and both improved to independent on discharge. Out of 24 cases who were moderately dependent at the time of admission, 54.2% got improved to mild dependency and rest have same outcome at the time of discharge. In severely dependent group there were 16 cases, 50% cases had no change in dependency on discharge, 37.5% improved to moderate and 6.3% become totally dependent. Out of 8 cases with total dependency at admission 25% got slight improvement and rest remained the same. Most of the mild and moderate dependent cases on admission had good short- term functional outcomes. Those who were severely dependent on admission had made no significant improvements in short-term functional outcomes.

CRP and stroke severity by NIHSS score was significantly related in this study (Chi- square value 23.46, p value=0.001). When CRP is less than 6mg/L, maximum number of cases comes under minor to moderate stroke. When CRP values more than 6mg/L, cases had moderate- severe to severe stroke. This indicates that, the increase in CRP levels were directly proportional to stroke severity.

According to the findings, hypertension, dyslipidaemia, and smoking are all associated with dependency after an acute ischemic stroke. Higher grades of dependency were seen in those who had any of these risk factors. Since the sample size is small, further studies with a bigger sample size are needed to assess the relationship between other risk factors and dependency.

In this study, when serum CRP levels were compared with short-term functional outcome, they showed a positive correlation. That is, when CRP levels are less than 6 mg/L, functional outcomes are better, with the majority of patients having mild, moderate, or independent status; when CRP levels are greater than 6 mg/L, functional outcomes are poor, with the majority of patients having severe or total dependency (chi-square value 46.06 and p value 0.001).

In this study group, only 19 patients underwent thrombolysis according to ASA criteria, of whom the maximum number of patients had a better outcome. One of the thrombolysed patients was independent; eight each had mild or moderate dependency; and two were severely dependent. Among non-thrombolysed patients, 10 had severe dependency, 5 had total dependency, and 9 had moderate dependency. It concludes that with

thrombolysis, outcomes can be improved.

V. CONCLUSION

In this study, functional outcomes were better when CRP levels were below the normal range (6 mg/L), while they were worse when CRP levels were above the normal range.

CRP levels in the very early phase of acute ischemic stroke can be used as an independent prognostic factor for short term outcome and for assessing the severity of stroke.

REFERENCES

- [1] Youn CS, Choi SP, Kim SH, Oh SH, Jeong WJ, Kim HJ, Park KN. Serum highly selective C-reactive protein concentration is associated with the volume of ischemic tissue in acute ischemic stroke. *The American journal of emergency medicine*. 2012 Jan 1;30(1):124-8.
- [2] Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, Moran AE, Sacco RL, Anderson L, Truelsen T, O'Donnell M. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *The lancet*. 2014 Jan 18;383(9913):245-55.
- [3] Warlow CP. Epidemiology of stroke. *The Lancet*. 1998 Oct 1;352: S1-4.
- [4] Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ, Culebras A, Elkind MS, George MG, Hamdan AD, Higashida RT, Hoh BL. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013 Jul;44(7):2064-89.
- [5] Easton JD, Saver JL, Albers GW, Alberts MJ, Chaturvedi S, Feldmann E, Hatsukami Functional Outcome in an Acute Ischemic Stroke TS, Higashida RT, Johnston SC, Kidwell CS, Lutsep HL. Definition and evaluation of transient ischemic attack: a scientific statement for healthcare professionals from the American Heart Association/American Stroke Association Stroke Council. *Stroke*. 2009 Jun 1;40(6):2276-93.
- [6] Whiteley W, Chong WL, Sengupta A, Sandercock P. Blood markers for the prognosis of ischemic stroke: a systematic review. *Stroke*. 2009 May 1;40(5): e380-9.
- [7] Di Napoli M, Papa F, Bocola V. C-reactive protein in ischemic stroke: an independent prognostic factor. *Stroke*. 2001 Apr;32(4):917- 24.
- [8] Montaner J, Fernandez-Cadenas I, Molina CA, Ribó M, Huertas R, Rosell A, Penalba A, Ortega L, Chacón P, Alvarez-Sabín J. Poststroke C-reactive protein is a powerful prognostic tool among candidates for thrombolysis. *Stroke*. 2006 May 1;37(5):1205-10.
- [9] Topkian R, Strasak AM, Nussbaumer K, Haring HP, Aichner FT. Prognostic value of admission C-reactive protein in stroke patients undergoing iv thrombolysis. *Journal of neurology*. 2008 Aug;255(8):1190-6.
- [10] Rost NS, Wolf PA, Kase CS, Kelly-Hayes M, Silbershatz H, Massaro JM, D'Agostino RB, Franzblau C, Wilson PW. Plasma concentration of C-reactive protein and risk of ischemic stroke and transient ischemic attack: the Framingham study. *Stroke*. 2001 Nov 1;32(11):2575-9.