

Formulate And Evaluate Antioxidant Gargle

Alshifa

Maharana Pratap School of Pharmacy

Abstract- The research of mouth gargle on for anti oxidation is the helpful in the oxidation activity. Early in the course of the disease, accurate diagnosis of sore throat can be challenging and requires a strong clinical suspicion, a careful examination, and the right testing. Anti oxidation of herbal mouth gargle on gives about a brief relief in the soreness, pain and fatigue. Gargles formulations contains volatile oils which are been used to treat different types of pains such as reduce pain and discomfort associated with a sore throat caused by bacterial and viral infections, mouth ulcers or other allergic reactions. Whether it is also has been treated as in the treatment of acute infections of the lining of the mouth and throat. It comprises a blend of volatile oils, including ginger and clove oil. Typically, these oils are used to treat pain, migraines, anti-inflammatory, anxiety, sinus congestion, and other ailments. This oral painkiller, which is made entirely of natural ingredients, gives off a pleasant sensation that helps to reduce the pain. The results of the study demonstrated that a herbal mouth gargle formulation containing herbal oils might be regarded as a promising skin enhancer to increase the therapeutic efficacy of medications. Specifically, the potential role of clove oil as an anti-inflammatory agent as well as its potential to improve how well ginger oil reduces inflammation.

INTRODUCTION

Sore Throat Infection

The painful feeling of soreness together with throat irritation becomes pronounced when swallowing occurs. Above all viral infections such as flu and cold dominate as the main leading causes of sore throat infections. Antibiotic medications are needed to treat bacterial strep throat infections. Several things other than fungi and parasites result in sore throat including smoke and allergens. A sore throat infection shows symptoms which include fever together with nausea and headache as well as vomiting and loss of appetite.

Cells require antioxidants to defend against free radical damage because this damage leads to aging conditions along with diseases including cancer and heart disease. Plant-based foods containing

antioxidants can primarily be found in colorful varieties and they function to counteract oxidative stress. Professional guidance must accompany the misuse of antioxidants as these nutrients hold essential functions but excessive supplementation provides no proven advantages.

Mechanism of Action of Antioxidants

Free radicals receive neutralization by antioxidants through their HAT and SET variations. The mechanism of action for phenolic compounds includes donating hydrogen atoms and binding metals. The properties of antioxidants regarding effectiveness are influenced by their concentration levels as well as by their chemical nature in combination with environmental conditions. The antioxidant capacity testing methods evaluate either electron transfer or hydrogen transfer mechanisms.

Curcumin (Turmeric)

Curcumin which derives from turmeric functions as a polyphenol compound that acts as an antioxidant and anti-inflammatory agent thus treating arthritis and metabolic syndrome and oxidative stress. The poor bioavailability of curcumin hinders its effectiveness although addition of piperine increases the amount absorbed by the body. Curcumin modifies genetic activity and regulates cholesterol metabolism while enhancing insulin reception and mental processing which indicates its potential as therapeutic medicine.

Benefits of Curcumin

The joint swelling and stiffness in arthritis patients could potentially decrease with curcumin consumption. Studies indicate that curcumin shows potential in treating eye diseases that affect the immune system and controls symptoms of hay fever specifically by decreasing sneezing and congestion. Scientific evidence shows potential anticancer properties of curcumin though continued research must verify its capacity to protect against or treat cancer.

Oral Drug Delivery System

The oral delivery route serves as a popular non-invasive approach for drugs since it boosts patient adherence even though digestive restrictions hinder drug absorption rates. The targeted delivery effects of drug administration come from alternative routes which include IV, IM, ID, IN and transdermal delivery methods. The selection process for optimal drug delivery systems highly depends on drug formulation techniques together with absorption mechanisms.

Challenges Associated with Oral Delivery

The methods that use the mouth to give drugs experience complications from GI tract enzyme breakdown along with insufficient drug solubility as well as chemical breakdown during stomach transit. The jejunum and ileum part of the small intestine provides the best drug absorption yet mucosal barriers along with inconsistent GI conditions result in inconsistent drug availability and therapeutic outcomes.

MATERIALS AND METHODS

Chemical List for the preparation

S.No.	Ingredients	Manufacture
1.	Curcumin	In Laboratory
2.	Ginger oil	Balsons Laboratories Naraina Delhi
3.	Clove oil	Balsons Laboratories Naraina Delhi
4.	Neem oil	Cargo Enterprises
5.	Honey	In Laboratory

Methods

Fourier Transform Infrared Spectroscopy (FTIR)
Using a spectrophotometer with the model number ATR-FTIR Perkin-Elmer 1005, samples of curcumin were collected in KBr pellets and scanned in the IR ranges of 2900 to 3050 cm⁻¹ and 1000 to 4000 cm⁻¹, respectively.

NMR Characterization

To illustrate the oil-related characteristics of the mouth gargles formulation, herbal oils were added to mouth gargles and deuterated methanol at room temperature. Two-dimensional NMR spectra of a roll on containing herbal oils remained logged via a Varian 400 mhz mass spectrometer.

Ultraviolet spectroscopy

Spectral and measures of absorbance using a GBC Cintra-10 UV/ Visible Spectrophotometer with 1400 nm/min scan speed, 1.006 nm data interval, and 2.0 nm constant slit width. Oils are soluble in methanol and dichloromethane because of this.

Method for the preparation of standard curve

A 25ml volumetric flask containing 0.025g of Curcumin was carefully weighed, then the substance was dissolved in a DCM-Me OH (3:2v/v) solvent mixture and the volume was adjusted to the desired amount. The final solution contained 1.0 mg of curcumin per milliliter.

Preparation of Calibration Curve at Curcumin

Aliquots of 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 0.9, 1.0, 1.1, 1.2, ml of std solution were transferred hooked on 10 ml volumetric flasks and the volume completed up with solvent system. The transmission density of the resulting resolutions was read at 283nm 362nm against a solvent blank.

FORMULATION AND DEVELOPMENT

- i. Firstly, take all the ingredients.
- ii. Then, take oil formulations of ginger, neem and clove into a beaker and mix them properly with a continuous stirring.
- iii. After that, add honey and curcumin into the solution and stir them properly.
- iv. After that leave the container for 7days at room temperature for the evaluation parameter study.

Composition of the mouth gargles with curcumin

S.no	Ingredients	Concentration%(w/v)		
		CGF1	CGF2	CGF3
1	Curcumin	13	10	10
2	Clove oil	4	7	6
3	Neem oil	5	7	4
4	Aleovera	12	10	8
5	Honey	3	8	5
6	Ginger oil	20	10	25

Evaluation of Herbal Mouth gargles

Visual observation and digital pH meter evaluation determined the color and odor and measured the pH of the herbal mouth gargle. The testing for microbial growth required agar plate incubation and monitoring. A stability assessment based on ICH guidelines was

combined with an antibacterial *Streptococcus mutans* test that evaluated zone of inhibition through agar well diffusion tests.

SUMMARY AND CONCLUSION

The preparation and evaluation of the curcumin mouth gargle formulation was the primary goal of the work. With the use of several stabilizers, this formulation will focus on oral distribution while minimizing adverse effects through the use of mouth gargles. These less hazardous oils were used to create this mouth gargle composition.

A complete investigation will be needed based on the animal research, even if the preliminary findings from in vitro release and permanency revisions show that this preparation is effective. The real mode of action for this kind of dosage form can then be discovered. The homogenization process and stabilizers needed to be carefully chosen. The stability of mouth gargles with the different excipients used in our experiment is validated by NMR and UV analysis, showing that the formulation method is appropriate for the production of good quality of mouth gargle

Prior to formulation optimization, thorough research is needed to create a product that is safer, more efficient, and affordable for human use.

REFERENCES

- [1] Chen, J.; Yang, J.; Ma, L.; Li, J.; Shahzad, N.; Kim, C.K. Structure-antioxidant activity relationship of methoxy, phenolic hydroxyl, and carboxylic acid groups of phenolic acids. *Sci. Rep.* 2020, 10, 2611–2620.
- [2] Wach, A.; Sa, J.; Szlachetko, J. Comparative study of the around-Fermi electronic structure of 5d metals and metal-oxides by means of high-resolution X-ray emission and absorption spectroscopies. *J. Synchrotron Rad.* 2020, 27, 689–694.
- [3] S. Gopi, S. Thomas, A.B. Kunnumakkara, B.B. Aggarwal, A. Aggarwal, Evaluation and clinical comparison studies on liposomal and non-liposomal ascorbic acid (vitamin C) and their enhanced bioavailability, *J. Liposome Res.* (2020) 1–9
- [4] S. Dewanjee, P. Chakraborty, B. Mukherjee, V. De Feo, Plant-based antidiabetic nano formulations: the emerging paradigm for effective therapy, *Int. J. Mol. Sci.* 21 (6) (2020) 2217.
- [5] V.K. Maurya, S. Kumar, A.K. Prasad, M.L.B. Bhatt, S.K. Saxena, Structure-based drug designing for potential antiviral activity of selected natural products from Ayurveda against SARS-CoV-2 spike glycoprotein and its cellular receptor, *Virusdisease* 31 (2) (2020) 179–193.
- [6] Nakagawa, Y.; Mukai, S.; Yamada, S.; Murata, S.; Yabumoto, H.; Maeda, T.; Akamatsu, S. The Efficacy and Safety of Highly-Bioavailable Curcumin for Treating Knee Osteoarthritis: A 6-Month Open-Labeled Prospective Study. *Clin. Med. Insights Arthritis Musculoskelet. Disord.* 2020
- [7] Ahmed Nasef, N.; Loveday, S.M.; Golding, M.; Martins, R.N.; Shah, T.M.; Clarke, M.; Coad, J.; Moughan, P.J.; Garg, M.L.; Singh, H. Food matrix and co-presence of turmeric compounds influence bioavailability of curcumin in healthy humans. *Food Funct.* 2019, 10, 4584–4592.
- [8] A. Hadi, M. Pourmasoumi, E. Ghaedi, A. Sahebkar, The effect of Curcumin/ Turmeric on blood pressure modulation: a systematic review and meta-analysis, *Pharm. Res.* 150 (2019), 104505.
- [9] J. Huang, S. Qin, L. Huang, Y. Tang, H. Ren, H. Hu, Efficacy and safety of *Rhizoma curcumae longae* with respect to improving the glucose metabolism of patients at risk for cardiovascular disease: a meta-analysis of randomised controlled trials, *J. Hum. Nutr. Diet.* 32 (5) (2019) 591–606.
- [10] P.B. Goncalves, N.C. Romeiro, Multi-target natural products as alternatives against oxidative stress in chronic obstructive pulmonary disease (COPD), *Eur. J. Med. Chem.* 163 (2019) 911–931.
- [11] Eggers M, Koburger-Janssen T, Eickmann M, Zorn J. In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/ mouthwash against respiratory and oral tract pathogens. *Infect Dis Ther* 2018;7:249-59.
- [12] Murata, Y., Isobe, T., Kofuji, K. and Nishida, N., “Preparation of fast dissolving films for oral dosage from natural polysaccharides.” *Materials*. 2010 Bansal P, Sujlana A, Pannu P, Kour R, Dental discomfort Questionnaire: correlated with clinical manifestations of advanced dental caries in young children, *J Dent Specialitation*, 2017.

- [13] Renuka s, Muralidharannp. Comparison in benefits of herbal mouthwashes with chlorhexidine Mouthwash: a review. *Asian J Pharm Clin Res*, Vol 10, Issue 2, 2017.
- [14] Kulkarni P, Singh DK, Jalaluddin M, Mandal A. Comparative evaluation of antiplaque efficacy between essential oils with alcohol-based and chlorhexidine with nonalcohol-based mouthrinses. *J Int Soc Prev Community Dent*. 2017;7:S36–41.
- [15] T. Venkataraman, M.B. Frieman, The role of epidermal growth factor receptor (EGFR) signaling in SARS coronavirus-induced pulmonary fibrosis, *Antivir. Res.* 143 (2017) 142–150.
- [16] Centers for Disease Control and Prevention, 2016. Wide-ranging Online Data for Epidemiologic Research (WONDER). CDC, National Center for Health Statistics, Atlanta, GA Accessed 11 July 2016.
- [17] Gopi, S. A Novel Composition of Curcumin with Enhanced Bioavailability. U.S. Patent US20160151440 A1, 2 June 2016.
- [18]] G. Kaur, M. Invally, M. Chintamaneni, Influence of piperine and quercetin on antidiabetic potential of curcumin, *J. Complement Integr. Med.* 13 (3) (2016) 247–255.
- [19] Fahrner, R.L. Compositions and Methods for Increasing the Bioavailability of One or More Compounds. WO Patent WO2016057839A1, 14 April 2016.
- [20] L.G.Vijayaalakshmi, R.V.Geetha. Comparison of Herbal Mouth Wash with Conventional Mouth Wash in Use in Reducing Streptococcus Mutans - An In vitro Study. *J. Pharm. Sci. & Res.* Vol. 7(7), 2015, 485-486. F.M Viera, M.C Macial, F.R Nascimento and V.P Rodrigues, Plant species used in dental disease: activity evaluation, *Journal of ethanopharmacology*. Vol. 155 (3), 1441-1449, 2014
- [21] Lake, S., Wood, E., Buxton, J., Dong, H., Montaner, J., Kerr, T., 2015b. Prescription opioid use and non-fatal overdose in a cohort of injection drug users. *Am. J. Drug Alcohol Abuse* 41, 257–263.
- [22] Havens, J.R., Leukefeld, C.G., DeVeaugh-Geiss, A.M., Coplan, P., Chilcoat, H.D., 2014.
- [23] Geetha R.V, Anitha Roy. In Vitro Evaluation of Anti bacterial Activity of Ethanolic root extract of Glycyrrhizaglabra on Oral microbes. *Int. J. Drug Dev. & Res.*, October-December 4(4), 2012, 161-165.
- [24] Purdue Pharma L.P. OxyContin® – oxycodone Fadden, D., Kulkarni, N. and Sorg, A. “Fast dissolving orally consumable films containing a modified starch for improved heat and moisture resistance.” WO/2004/096193, 2004.
- [25] KulkarniKumar, L., Sorg, A. “Fast dissolving orally consumable films containing a sucralose as a sweetener.” WO/2004/096192, 2004.
- [26] Chandak C. Master of Sci. Thesis, “Orally dissolving film of selegiline hydrochloride” London Island University, May 2009
- [27] Kumar P, Ansari S.H, Ali .J . Herbal remedies for the treatment of pridental disease: A patient review. *Recent pat drug deliver formula* 2009;3:221-8.
- [28] Aulton. M.E. *Pharmaceutics: the science of dosage form design*, 2nd Edition, Churchill Livingstone, Elsevier Ltd. UK, 2002,
- [29] Shuaib Mohd., Ali M., Naquvi K. J. J. *Chem. Pharm. Res.*, 2007, 5(8),42-47.
- [30] Kariwa H, Fujii N, Takashima I. Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions and chemical reagents. *Dermatology* 2006;212(Suppl 1):119-23.
- [31] El Ashry E., Rashed N., Salma O., Saleh A. *Pharmazie*, 2003, 58(3),163-168Bergamini, C.M.; Gambetti, S.; Dondi, A.; Cervellati, C. Oxygen, Reactive Oxygen Species and Tissue Damage. *Curr. Pharm. Des.* 2004, 10, 1611–1626
- [32] Piotrowska, A. *Modulacyjny Wpływ Witaminy D na Działanie Reaktywnych form Tlenu w Keratynocytach*. Ph.D. Thesis, Medical University of Gdańsk, Gdańsk, Poland, 2014.
- [33] Satoh, K.; Kadofuku, T.; Sakagami, H. Effect of Trolox, a synthetic analog of alpha-tocopherol, on cytotoxicity induced by UV irradiation and antioxidants. *Anticancer Res.* 1997, 17, 2459–2463.
- [34] Heim, K.E.; Tagliaferro, A.R.; Bobilya, D.J. Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. *J. Nutr. Biochem.* 2002, 13, 572–584
- [35] Mughesh, G.; Panda, A.; Singh, H.B.; Puneekar, N.S.; Butcher, R.J. Glutathione peroxidase-like antioxidant activity of diaryl diselenides: A

- mechanistic study. *J. Am. Chem. Soc.* 2001, *123*, 839–850.
- [36] H.W. Chen, H.T. Kuo, C.Y. Chai, J.L. Ou, R.C. Yang, Pretreatment of curcumin attenuates coagulopathy and renal injury in LPS-induced endotoxemia, *J. Endotoxin Res* 13 (1) (2007) 15–23.
- [37] H.W. Chen, H.T. Kuo, C.Y. Chai, J.L. Ou, R.C. Yang, Pretreatment of curcumin attenuates coagulopathy and renal injury in LPS-induced endotoxemia, *J. Endotoxin Res* 13 (1) (2007) 15–23.
- [38] Yamamura Y, Kawakami J, Santa T, Kotaki H, Uchino K, Sawada Y, Tanaka N, Iga T. Pharmacokinetic profile of glycyrrhizin in healthy volunteers by a new high performance liquid chromatographic method. *J Pharm Sci* 1992;81:1042
- [39] Persson B-A, Lagerstrom P-0, Grundevik I. Determination of omeprazole and metabolites in plasma and urine. *Scand J Gastroenterol* 1985;20(suppl 108):71-7.
- [40] W. H. Barr, Symposium on Formulation Factors Affecting Therapeutic Performance of Drug Products, Washington, D. C., Apr. 1969; Drug Information Bull., 1969.