

Antibacterial Activity of Medicinal Plants against Pathogens causing Intricate Urinary Tract Infections

Dr. D. Sripriya

Department of Biotechnology, School of Herbal Studies and Naturo Sciences, Dravidian University, Kuppam – 517 426, A.P., India.

Abstract—Seventeen Indian folklore medicinal plants were investigated to evaluate antibacterial activity of aqueous, ethanol and acetone extracts against 66 multidrug resistant isolates of major urinary tract pathogens (*Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Enterococcus faecalis*) by disc diffusion method. Ethanol extract of *Zingiber officinale* and *Punica granatum* showed strong antibacterial activity against *Escherichia coli*. Ethanol extracts of *Terminalia chebula* and *Ocimum sanctum* exhibited antibacterial activity against *Klebsiella pneumonia*. Ethanol extract of *Cinnamomum, cassia* showed maximum antibacterial activity against *Pseudomonas aeruginosa* while ethanol extract of *Azadirachta indica* and *Ocimum sanctum* exhibited antibacterial activity against *Enterococcus faecalis*. The results support the folkloric use of these plants in the treatment of urinary tract infections by the tribal of Seshachalam Hills region of South India.

Keywords—Urinary tract infection (UTI), Multidrug resistant, Antibacterial activity, Indian medicinal plants.

I. INTRODUCTION

Urinary tract infections (UTI) are most common form of bacterial infections, affecting people throughout their lifespan. The pathogenesis of complicated and uncomplicated UTI is complex and influenced by many host biological behavioral factors and

properties of the infecting urea pathogens. Leading etiological agents of UTI's include *Escherichia coli*, *Candida albicans*, *Enterococcus faecalis*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia* and *Proteus mirabilis*. The incidence of acute uncomplicated UTI is estimated to exceed 0.5 episodes per annum among women between 18-30 years. Despite the existence of potent antibiotics, resistant or multi resistant strains are continuously appearing, imposing the need for a permanent search and development of new drugs. For Centuries plants have been used throughout the world as drugs and remedies for various diseases. These drugs serve as prototype to develop more effective and less toxic medicines. Hence, an attempt has been made to evaluate antibacterial activity of seventeen folklore medicinal plants used by tribal in Seshachalam Hills region of South India against urinary tract pathogens.

Plant parts were collected on the basis of the information provided in the ethnobotanical survey of India and local medicine men of Seshachalam Hills region (AP), India. Each specimen/plant material was labeled, numbered, a noted with the date of collection, locality and their medicinal uses were recorded (Table 1). Fresh plant material was washed thoroughly, air dried and then homogenized to fine powder and stored in airtight containers.

Table 1: Selected Indian medicinal plants for antibacterial activity

Sr.No.	Botanical Name	common name	Family	Part used	Medicinal uses
1.	<i>Zinzibar officinale</i>	Allam	Zingiberaceae	Rhizome	Analgesic, sedative, antipyretic and antibacterial.
2.	<i>Cinnamomum verum</i>	Dalchina chakka	Lauraceae	Bark	Antibacterial, circulatory, respiratory, uterotonic and stomachic
3.	<i>Terminalia chebula</i>	Karakkaya	Combretacea e	Fruit	Laxative, stomachic, tonic and alternative.
4.	<i>Plantago ovata</i>	Avalokanam	Plataginaceae	Husk	Constipation, colitis, irritable bowel, cystitis

5.	<i>Acacia nilotica</i>	Nalla tumma	Fabaceae	Leaf	Treating premature ejaculation
6.	<i>Pimpinella anisum</i>	Anas Puvvu	Apiaceae	Seed	Antiseptic, antispasmodic, aromatic, carminative, digestive, galactogogue, pectoral, stimulant
7.	<i>Centella asiatica</i>	Brahmi	Apiaceae	Stem	Tonic, sedative, alterative.
8.	<i>Ocimum sanctum</i>	Tulsi	Laminaceae	Leaf	Cures cough, cold, chronic, catarrh, nausea, respiratory problem, skin diseases, antibacterial
9.	<i>Azadirachta Indica</i>	Neem/ Vepa	Meliaceae	Fruit	Skin disease, blood disorder, antibacterial.
10.	<i>Phyllanthus fraternus</i>	Nela Usiri	Euphorbiaceae	Leaf	Jaundice, liver disease, fever, genitourinary disease, edema
11.	<i>Coriandrum sativum</i>	Dhaniyalu	Apiaceae	Seed	Flatulence, colic, joint pain, antiseptic
12.	<i>Abutilon indicum</i>	Atibala	Malvaceae	Stem	Demulcent, aphrodisiac, laxative, astringent and diuretic, expectorant, antiinflammatory and analgesic
13.	<i>Punica granatum</i>	pomegranate	Lythraceae	Seed	Anthelmintic (esp. tapeworm), diarrhea, dyspepsia
14.	<i>Syzygium cumini</i>	Neredu	Myrtaceae	Bark	Astringent, stomachic, carminative, antiscorbutic, diuretic.
15.	<i>Cyperus scariosus</i>	Tungagaddalaveru	Cyperaceae	Root	Astringent, diaphoretic, desiccant, cordial and stomachic
16.	<i>Andrographis paniculata</i>	Nelavemu	Acanthaceae	Bark	Laxative, vulnerary, antipyretic, antiperiodic, antiinflammatory, expectorant, depurative, antibacterial
17.	<i>Mangifera indica</i>	Mamidi	Anacardiaceae	Leaf	Supplement of sexual potency, antiallergic, hypoglycaemic and antidiabetic.

Plant material (10 g) was extracted with 100 ml distilled water by subjecting it slow heat for 6 h; filtered through muslin cloth and centrifuged at 5000×g for 15 min. The supernatant was collected and concentrated to one-fourth of the total volume.

Organic solvent extraction of the plant materials was prepared according to the method described by Nair et al. with some modifications. 10 g of plant material was crushed and blended into powder using an electric blender with each organic solvent i.e. ethanol


and acetone separately. The blended material was subjected to agitation on rotary shaker (190-220 rpm) for 30 min, filtered with muslin cloth; centrifuged (5000×g for 15 min), and concentrated to one-fourth of the total volume.

The test organisms included the gram-positive *E. faecalis* (7 isolates) and gram-negative *E. coli* (34 isolates), *K. pneumonia* (15 isolates), and *P. aeruginosa* (10 isolates). All the 66 UTI isolates were obtained from the bacterial germplasm culture collection, Bacteriology laboratory, Kakatiya University, Warangal, Telangana, India. The bacteria were grown in the nutrient broth at 37° and maintained on nutrient agar slants at 4°.

In vitro antibacterial activity was examined for aqueous, ethanol and acetone extracts from 17 different traditionally used medicinal plants. Antimicrobial activities of these extracts were evaluated by Agar disc diffusion method. For all the bacterial strains, overnight cultures grown in broth were adjusted to an inoculum size of 10⁶ CFU/ml for inoculation of the agar plates. A lawn of UTI bacterial culture was made on Mueller Hinton agar plates and sterile filter paper discs soaked with 10 µl (concentration 50 mg/ml) each of aqueous, ethanol and acetone extract was placed on it. Disc soaked in distilled water, ethanol and acetone served as control. Following an incubation period of 24 h at 37°, antibacterial activity was evaluated by inhibition zones of bacterial growth. The results are represented as average zone of inhibition of all the isolates of individual species.


Fifty-one aqueous, ethanol and acetone extracts of the plant belonging to 17 families were tested against gram-positive and gram-negative urinary tract pathogenic bacteria. Ethanolic extract showed considerably more activity than the acetone and aqueous extract. Maximum antibacterial activity was shown by ethanol extract of *Z. officinale*, *P. granatum* and acetone extract of *T. chebula* against *E. coli* with the inhibition of 14 mm zone size (fig. 1). Similar results were observed by previous scientist against pathogenic bacteria. Ethanol extracts of *T. chebula* and *O. sanctum* were found to be active with the inhibition of 16 mm against *K. pneumonia* (fig. 2) while ethanol extract of *C. cassia* showed significant activity with 18 mm of clear inhibition zone against *P. aeruginosa* (fig. 3). Previous reports from china have revealed that various parts of *C. cassia* such as the fruit, oil, inner bark and leafy twigs possessed good antimicrobial activity. The ethanol extract of *C. asiatica* and *O. sanctum* show broad-spectrum activity against *E. faecalis* with the inhibition zone of 19 mm (fig. 4). Best antimicrobial activity was found in case of ethanol extract of *Z. officinale*, *P. granatum*, *T. chebula*, *O. sanctum*, *C. cassia*, *C. asiatica* and acetone extract of *T. chebula* against multidrug resistant UTI pathogens. This probably explains the use of these plants by indigenous people against a number of infections since generations. The plants studied here had shown that they are potentially rich in antimicrobial compounds and have also been extensively used by the tribal. The millenarian use of these plants in folk medicine suggests that they represent an economic and safe alternative for treatment of Urinary Tract Infections.

Fig: 1




Antibacterial activity of plant extracts against *Escherichia coli*. Antibacterial activity of aqueous, ethanol and acetone extracts of plants against *Escherichia coli*.

Fig: 2




Antibacterial activity of plant extracts against *Klebsiella pneumonia*. Antibacterial activity of aqueous, ethanol and acetone extracts of plants against *Klebsiella pneumonia*.

Fig: 3



Antibacterial activity of plant extracts against *Pseudomonas aeruginosa*, Antibacterial activity of aqueous, ethanol and acetone, extracts of plants against *Pseudomonas aeruginosa*.

Fig: 4



Antibacterial activity of plant extracts against *Enterococcus faecalis*, Antibacterial activity of aqueous, ethanol and acetone extracts of plants against *Enterococcus faecalis*.

REFERENCES

- [1] Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Am J Med.* 2002;113:5S–13S. [PubMed] [Google Scholar]
- [2] Barnett BJ, Stephens DS. Urinary tract infection: an overview. *Am J Med Sci.* 1997;314:245–9. [PubMed] [Google Scholar]
- [3] Svanborg C, Godaly G. Bacterial virulence in urinary tract infection. *Infect Dis Clin N Am.* 1997;11:513–29. [PubMed] [Google Scholar]
- [4] Hooton TM, Scholes D, Hughes JP, Winter C, Roberts PL, Stapleton AE, et al. Prospective study of risk factors for symptomatic urinary tract infection in young women. *N Eng J Med.* 1996;335:468–74. [PubMed] [Google Scholar]
- [5] UNESCO: Culture and Health. Orientation-Texts World Decade for Culture Development 1988-1997. Paris, France: 1996. Document CLT/DEC/PRO. [Google Scholar]
- [6] Tetyana P, Prozesky EA, Jager AK, Meyer JJ, Staden J. Some medicinal properties of *Cussonia* and *Schefflera* species used in traditional medicine. *South Afr J Botany.* 2002;68:51–4. [Google Scholar]
- [7] Nair R, Kalariya T, Chhanda S. Antibacterial activity of some selected Indian medicinal flora. *Turk J Biol.* 2005;29:41–7. [Google Scholar]
- [8] Rios JL, Recio MC, Vilar A. Screening methods for natural products with antimicrobial activity: a review of the literature. *J Ethnopharmacol.* 1988;23:127–49. [PubMed] [Google Scholar]
- [9] Collins CH, Lynes PM, Grange JM. *Microbiological methods.* 7th ed. Britain: Butterworth- Hinemann Ltd; 1995. pp. 175–90. [Google Scholar]
- [10] Bauer AW, Kirby WM, Scherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol.* 1966;45:493–6. [PubMed] [Google Scholar]
- [11] Konning GH, Ayare C, Ennison B. Antimicrobial activity of some medicinal plants from Ghana. *Fitoterapia.* 2004;75:65–7. [PubMed] [Google Scholar]
- [12] Aquil F, Khan MS, Owais M, Amad I. Effect of certain bioactive plant extracts on clinical isolates of β -lactamase producing methicillin resistant *Staphylococcus aureus*. *J Basic Microbiol.* 2005;45:106–14. [PubMed] [Google Scholar]
- [13] Holetz F, Pessini G, Sanches N, Cortez D, Nakamura C, Filho D. Screening of some plants used in the Brazilian folk medicine for the treatment of infectious diseases. *Mem Inst Oswaldo Cruz, Rio de Janeiro.* 2002;97:1027–31. [PubMed] [Google Scholar]
- [14] Bonjar GHS, Nik AK, Heydari MR, Ghasemzadeh MH, Farrokhi PR, Moein MR, et al. Anti-Pseudomonas and Anti-Bacilli activity of some medicinal plants of Iran. *DARU.* 2003;11:157–63. [Google Scholar]
- [15] Sripriya D. and Naik MRB. Ethno Medicinal plants of Seshachalam Biosphere Reserve, Andhra Pradesh, India. *International Journal of Pharmacy and Biological Sciences.* 2019b, 9, 2, 18-22. [Google Scholar]
- [16] Sripriya D. and Naik MRB. Ethnomedicobotanical studies of high Potential medicinal plants of seshachalam biosphere reserve, A.P, India. *International Journal of Engineering & Science Research.* 2019a, 9, 5, 22-27. [Google Scholar]
- [17] Sripriya D. Ethnobotanical uses of some Plants by Tribes in A.P, India. *International Journal of Pharmacy and Biological Sciences.* 2017a, 7, 4, 67-74. [Google Scholar]
- [18] Sripriya D. Traditionally Used Medicinal Plants for wound healing in the Chittoor Dt., Andhra Pradesh (India). *International journal of Life Sciences.* 2017b, 5, 4, 706 -708. Online. [Google Scholar]
- [19] D. Sripriya and Sandhya Sri. D (2023) “A Simple Analysis On Herbal Cosmetics” *International Journal of Science and Research Archive.* | Volume 08 | Issue 01 | Jan | 2023 | 001 – 016. eISSN: – 2582-8185, <https://doi.org/10.30574/ijrsra.2023.8.1.0318>, <https://ijrsra.net/> [Google Scholar]