

To Formulation and Evaluation of Herbal Hand Sanitizer and Comparative Assessment with Commercial Formulations

Rutika Dinkar Gaikwad¹, Mrs. Snehal Daud², Ritesh Sachin Daund³,
Rohit Sahebrav Gadhe⁴, Dr. D. K. Vir⁵
^{1,2,3,4,5}Shree Goraksh college of pharmacy and research centre

Abstract—Hand hygiene plays a crucial role in preventing the transmission of infectious diseases. Alcohol-based hand sanitizers are widely used due to their rapid antimicrobial action [1]. However, frequent use may lead to skin dryness and irritation [2]. Herbal hand sanitizers incorporating natural ingredients such as neem (*Azadirachta indica*) offer antimicrobial benefits with improved skin compatibility [3]. The present study focuses on the formulation of a neem-based herbal hand sanitizer and its evaluation for physicochemical properties, antimicrobial activity, and stability. The formulation includes isopropyl alcohol, aloe vera gel, neem extract, glycerine, and essential oils. Comparative analysis with commercial sanitizers was performed. The results indicate that the herbal sanitizer shows effective antimicrobial activity along with better moisturizing properties, making it a suitable alternative to conventional products.

Aim: To formulate and evaluate a neem-based herbal hand sanitizer

Objectives:

- To prepare herbal sanitizer formulation
- To evaluate physicochemical properties
- To assess antimicrobial activity
- To compare with commercial sanitizer

I. INTRODUCTION

Hand hygiene is one of the most effective measures to prevent infections caused by bacteria, viruses, and fungi [1]. The use of alcohol-based sanitizers has increased significantly, especially after global pandemics [17].

Alcohol-based sanitizers are effective when alcohol concentration is above 60% as they denature proteins and destroy microbial cell membranes [2]. However, repeated use may lead to:

- Skin dryness
- Irritation
- Dermatitis

Herbal formulations have gained attention due to their natural origin and reduced side effects [6].

Neem (*Azadirachta indica*) is widely known for:

- Antibacterial activity [4]
- Antifungal properties [5]
- Anti-inflammatory effects

Thus, combining alcohol with herbal components provides both immediate and long-term benefits.



Figure 1. Hand hygiene plays a crucial role in preventing the transmission of infectious diseases.

II. LITERATURE REVIEW

Sr. No.	Author(s)	Year	Title of Study	Journal (UGC / Peer-reviewed)	Key Findings
1	WHO Guidelines	2009	Hand Hygiene in Health Care	WHO Report	Recommended 75% alcohol sanitizer formulation for effective microbial control
2	Kampf & Kramer	2004	Hand Hygiene Review	Journal of Hospital Infection	Alcohol $\geq 60\%$ is essential for antimicrobial action
3	Subapriya & Nagini	2005	Medicinal Properties of Neem	Current Medicinal Chemistry	Neem shows antibacterial, antifungal, anti-inflammatory effects
4	Biswas et al.	2002	Biological Activities of Neem	Current Science	Neem contains bioactive compounds like azadirachtin
5	Khan & Raghav	2021	Neem & Aloe Herbal Hand Wash	Research Journal of Pharmacy and Technology	Developed herbal formulation with antimicrobial activity (RJPT)
6	Sharma et al.	2024	Review on Herbal Sanitizers	International Journal GRANTHAALAYAH	Herbal sanitizers effective when alcohol concentration is optimized
7	Jadhav et al.	2025	Herbal Hand Sanitizer Formulation	International Journal of Pharmaceutical Sciences	Neem + Aloe showed good antimicrobial and skin-friendly results
8	Sridhar et al.	2024	Aloe-based Herbal Sanitizer	International Journal of Research and Review	Aloe enhances antimicrobial effect and skin hydration
9	Tulsawani et al.	2024	Herbal-Alcohol Sanitizer Study	Antonie van Leeuwenhoek Journal	Standardized herbal sanitizer showed strong antimicrobial activity
10	Chaudhari et al.	2022	Herbal Sanitizer Review	Int J Adv Res Chem Sci	Aloe + neem improves sanitizer safety and efficacy
11	Siddhartha et al.	2023	Herbal Sanitizer Evaluation	World Journal of Pharmaceutical Research	Showed inhibition zones ~20–25 mm
12	Patankar et al.	2023	Neem vs Commercial Sanitizer	Int Research Journal of Pharmacy & Biosciences	Neem sanitizer effective even where commercial failed
13	Kumar et al.	2025	Alcohol-Free Herbal Sanitizer	JETIR Journal	Essential oils enhance antimicrobial activity
14	Shelke et al.	2026	Tulsi-Aloe Sanitizer	IJSDR (UGC CARE)	Herbal formulations showed improved skin compatibility
15	Gurav et al.	2025	Herbal Sanitizer Evaluation	Int Journal of Research in Science & Technology	Confirmed antimicrobial efficiency of herbal combinations
16	Rathod et al.	2025	Herbal Mouthwash Study	International Journal of Scientific Innovation	Neem & aloe effective antimicrobial agents (IJSCI)
17	Ankade et al.	2025	Herbal Lotion Formulation	IJIRT (UGC indexed)	Aloe & neem improve skin protection (IJIRT)

III. MATERIALS AND METHODS

3.1 Materials Used

1. Isopropyl Alcohol (99%)

Function: Primary antimicrobial agent Isopropyl alcohol is the main active ingredient responsible for the germicidal activity of the sanitizer. It acts by denaturing proteins and disrupting the cell membranes of microorganisms, leading to rapid cell death [1]. Alcohol concentrations between 60–95% are considered most effective, with 70–75% providing optimal antimicrobial action [2]. In this formulation, isopropyl alcohol constitutes approximately 75% of the total volume, ensuring effective elimination of

bacteria, viruses, and fungi. It provides instant sanitization without the need for water.



Figure 2. Isopropyl alcohol is the main active ingredient responsible for the germicidal activity of the sanitizer

2. Aloe Vera Gel

Function: Moisturizer and soothing agent

Aloe vera gel is incorporated to counteract the drying effect of alcohol. It contains polysaccharides, vitamins, and amino acids that help in skin hydration and repair [4]. Additionally, aloe vera forms a gel base that improves the viscosity and spread ability of the formulation. It also exhibits mild antimicrobial and anti-inflammatory properties, enhancing the overall effectiveness of the sanitizer [4].



Figure 3. aloe vera forms a gel base that improves the viscosity and spread ability of the formulation.

3. Neem Extract / Neem Oil (*Azadirachta indica*)

Function: Herbal antimicrobial and anti-inflammatory agent. Neem extract contains bioactive compounds such as azadirachtin, nimbidin, and flavonoids, which exhibit strong antibacterial, antifungal, and antiviral properties [3,6]. It enhances the antimicrobial spectrum of the sanitizer and provides additional protection against pathogens. Neem also has anti-inflammatory and skin-healing properties, making the formulation safer for frequent use [3].



Figure 4: Neem also has anti-inflammatory and skin-healing properties, making the formulation safer for frequent use

4. Glycerine (Glycerol)

Function: Humectant and skin protectant

Glycerine acts as a humectant, meaning it attracts moisture to the skin and prevents dehydration caused by alcohol [2,4]. It improves the texture and smoothness of the formulation and helps maintain skin hydration. WHO-recommended sanitizer formulations also include glycerine (~1.45%) for this purpose [2]. In this formulation, 2% glycerine is used for enhanced moisturizing effect.




Figure 5. glycerine is used for enhanced moisturizing effect.

5. Essential Oil (Tea Tree / Lemon Oil)

Function: Fragrance and secondary antimicrobial agent Essential oils are added to improve the odor and user acceptability of the sanitizer. Tea tree oil possesses strong antibacterial and antifungal properties, while lemon oil contains limonene, which has antimicrobial activity [7]. These oils also provide a refreshing fragrance and may contribute to additional microbial inhibition.



Figure 6. essential oils are added to improve the odor and user acceptability of the sanitizer

6. Distilled Water

Function: Solvent and volume adjustment

Distilled water is used to adjust the final volume of the formulation to 100 mL and to ensure uniform mixing of all ingredients. It helps maintain the desired concentration of alcohol and improves the consistency of the formulation.



Figure 7. Distilled water is used to adjust the final volume of the formulation

6. Equipment Used

- Analytical balance – for accurate measurement of ingredients
- pH meter – for determining pH of formulation
- Viscometer – for measuring viscosity
- Sterile Petri dishes – for antimicrobial testing
- Incubator – for microbial culture growth
- Glassware (beakers, measuring cylinders) – sterilized before use

Preparation of Extracts

Neem Extract Preparation

Fresh neem leaves (25 g) were washed and boiled in 100 mL distilled water for 15 minutes. The solution was filtered to obtain a clear aqueous extract containing active phytoconstituents [3].

Aloe Vera Gel Preparation

Fresh aloe vera leaves were peeled, and the inner gel was collected and homogenized to obtain a smooth gel suitable for formulation [3].

7. Formulation Procedure (SOP)

- Measure 75 mL isopropyl alcohol in a clean beaker

- Add 2 mL glycerine and mix gently
- Add 20 mL aloe vera gel and stir uniformly
- Add 2 mL neem extract and mix thoroughly
- Add 1 mL essential oil
- Adjust volume to 100 mL using distilled water
- Stir for 2–3 minutes to obtain homogeneous mixture
- Measure pH (~6.0) and adjust if necessary
- Transfer to sterile container and allow to stand for 24 hours



Figure 8. neem extract

Controls Used

- WHO formulation (75% isopropanol + glycerol) [2]
- Blank formulation (without alcohol)

Safety Precautions

- Perform preparation in a well-ventilated area
- Avoid open flames (alcohol is highly flammable)
- Use gloves and protective equipment
- Conduct patch test before use
- Maintain neem concentration $\leq 3\%$ to avoid irritation [5]

3.2 Formulation (100 ml)

Ingredient	Quantity
Isopropyl Alcohol	75 ml
Aloe Vera Gel	20 ml
Neem Extract	2 ml
Glycerine	2 ml
Essential Oil	1 ml

3.3 Method of Preparation

The formulation was prepared using standard mixing techniques [12]:

1. Isopropyl alcohol was measured accurately
2. Aloe vera gel was added slowly with continuous stirring
3. Neem extract and glycerine were incorporated
4. Essential oil was added for fragrance
5. The mixture was stirred to obtain a uniform consistency
6. The formulation was stored in sterile containers



Figure 8. formulation of hand sanitizer

IV. EVALUATION PARAMETERS

4.1 Physical Appearance: The prepared formulation was evaluated for color, odor, and texture [13].

4.2 pH Determination: The pH was determined using a digital pH meter and found to be within the acceptable range of 5.5–7, suitable for skin application [15].

4.3 Viscosity: Viscosity was measured using a Brookfield viscometer to ensure appropriate consistency [19].

4.4 Spread ability: Spread ability was evaluated to determine ease of application on the skin [19].

4.5 Antimicrobial Activity: The antimicrobial activity was tested using the agar well diffusion method against common pathogens such as *E. coli* and *S. aureus* [10].

4.6 Stability Study: The formulation was subjected to stability studies at different temperatures to observe changes in physical and chemical properties [20].

V. RESULTS AND DISCUSSION

The formulated herbal sanitizer exhibited good physical properties, including acceptable colour, Odor, and consistency. The pH was found to be within the skin-friendly range [15]. The presence of neem extract contributed significantly to antimicrobial activity due to its bioactive compounds such as nimbidin and azadirachtin [4]. Aloe vera improved the moisturizing effect and reduced dryness caused by alcohol [11]. The antimicrobial activity of the herbal formulation was found to be comparable to commercial sanitizers [6]. Stability studies indicated that the formulation remained stable under normal storage conditions [20].

TABLE 5.1: Physical Evaluation Parameter

Parameter	Observation	Result
Color	Light green	Acceptable
Odor	Mild herbal fragrance	Pleasant
Texture	Smooth gel	Good consistency
Clarity	Transparent / slightly translucent	No impurities
Phase Separation	Not observed	Stable

TABLE 5.2: Physicochemical Evaluation

Parameter	Observed Value	Standard Range	Inference
pH	6.1 ± 0.1	5.5 – 6.5	Skin-friendly
Viscosity	1200 ± 50 cP	1000 – 2000 cP	Ideal gel consistency
Spread ability	5.0 ± 0.2 cm	4 – 6 cm	Good spreading
Alcohol Content	74–75% v/v	$\geq 60\%$	Effective

TABLE 5. 3: Antimicrobial Activity
(Zone of Inhibition):

Microorganism	Herbal Sanitizer (mm)	Commercial Sanitizer (mm)	Blank (mm)
<i>Staphylococcus aureus</i>	19.5 ± 0.9	21.0 ± 1.0	0
<i>Escherichia coli</i>	18.2 ± 1.0	20.5 ± 1.2	0

TABLE 5.4: Minimum Inhibitory Concentration (MIC):

Microorganism	MIC (Herbal Sanitizer)	Interpretation
<i>S. aureus</i>	2% v/v	Highly effective
<i>E. coli</i>	2–3% v/v	Effective

TABLE 5. 5: Stability Study

Parameter	Initial	After 30 Days (40°C)	Observation
Color	Light green	No change	Stable
pH	6.1	6.0	Slight variation
Odor	Mild	No change	Stable
Phase Separation	None	None	Stable

TABLE 5.6: Skin Irritation Test

Volunteer No.	Observation	Score
1-10	No redness / irritation	0

Score Interpretation:

0 = No irritation

1 = Mild

2 = Moderate

TABLE 5.7: Comparative Evaluation:

Parameter	Herbal Sanitizer	Commercial Sanitizer
Antimicrobial Activity	Good	Very Good
pH	Skin-friendly (6.1)	Slightly alkaline
Moisturizing Effect	Excellent	Moderate
Skin Irritation	None	Mild
Cost	Low	

VI. ADVANTAGES

- Natural and safe ingredients [12]
- Skin-friendly formulation [11]
- Cost-effective [14]
- Eco-friendly [18]



Figure 9. The prepared sanitizer exhibited satisfactory physicochemical properties, including a light green appearance, smooth gel consistency,

VII. LIMITATIONS

- Shorter shelf life compared to synthetic products [20]
- Variability in herbal extract quality [18]
- Requires proper storage conditions

VIII. FUTURE WORK

Viral Testing: Assess efficacy against viruses (e.g. influenza, coronavirus) since alcohol should be effective, but herbal components' contribution is unknown.

Stability (Long-term): Conduct 6-12month stability studies at various temperatures to confirm shelf life.

Sensory Evaluation: Conduct user studies to refine scent and texture (some may prefer stronger fragrance or different consistency).

Ingredient Variations: Test other herbs (e.g. Tulsi or peppermint oil) for enhanced properties or appeal.

Sustainability: Explore biodegradable packaging and eco-friendly preservatives to further improve the product's profile



Figure 10. study concludes that the neem-based herbal hand sanitizer is an effective, stable, and skin-compatible formulation

IX. CONCLUSION

The present study successfully focused on the formulation and evaluation of a neem-based herbal hand sanitizer using isopropyl alcohol, aloe vera gel, neem extract, glycerine, and essential oil. The formulation was designed to combine the rapid antimicrobial action of alcohol with the therapeutic and skin-friendly properties of herbal ingredients.

The prepared sanitizer exhibited satisfactory physicochemical properties, including a light green appearance, smooth gel consistency, and pH of 6.1, which lies within the acceptable skin-friendly range. The viscosity and spread ability were found to be optimal, ensuring ease of application and adequate coverage on the skin surface.

The antimicrobial evaluation demonstrated significant activity against both Gram-positive (*Staphylococcus aureus*) and Gram-negative (*Escherichia coli*) bacteria, with inhibition zones comparable to commercial alcohol-based sanitizers. This confirms that the formulation is effective in eliminating pathogenic microorganisms. The presence of neem extract enhanced the antimicrobial spectrum, while aloe vera and glycerine improved moisturizing properties and reduced the drying effect of alcohol. Stability studies indicated that the formulation remained stable under accelerated conditions, showing no significant changes in color, pH, or phase separation. Additionally, the skin irritation test confirmed that the formulation is safe for topical application and suitable for frequent use.

Overall, the study concludes that the neem-based herbal hand sanitizer is an effective, stable, and skin-compatible formulation. It offers a promising natural alternative to conventional sanitizers by combining antimicrobial efficacy with enhanced user safety and comfort. This formulation can be further developed for large-scale production and commercial application in the field of personal hygiene.

REFERENCES

- [1] World Health Organization, Guidelines on Hand Hygiene in Health Care. Geneva: WHO Press, 2009.
- [2] G. Kampf and A. Kramer, "Epidemiologic background of hand hygiene," J. Hosp. Infect., 2004.
- [3] Centers for Disease Control and Prevention (CDC), Hand Hygiene Guidelines, 2020.
- [4] J. M. Boyce and D. Pittet, "Guideline for hand hygiene in healthcare settings," MMWR, 2002.
- [5] A. F. Widmer, "Replace hand washing with alcohol rub?" Clin. Infect. Dis., 2000.
- [6] K. Biswas et al., "Biological activities of neem," Curr. Sci., 2002.
- [7] R. Subapriya and S. Nagini, "Medicinal properties of neem," Curr. Med. Chem., 2005.
- [8] S. C. Gupta et al., "Neem: A plant of medicinal importance," J. Ethnopharmacol., 2017.
- [9] M. A. Alzohairy, "Therapeutics role of neem," Evid. Based Complement. Altern. Med., 2016.
- [10] O. Koul et al., "Neem bioactive compounds," J. Sci. Ind. Res., 2004.
- [11] A. Surjushe et al., "Aloe vera: A short review," Indian J. Dermatol., 2008.
- [12] J. H. Hamman, "Composition and applications of Aloe vera," Molecules, 2008.
- [13] K. Eshun and Q. He, "Aloe vera: A valuable ingredient," Crit. Rev. Food Sci., 2004.
- [14] S. Choi and M. H. Chung, "Aloe vera effects on wound healing," J. Ethnopharmacol., 2003.
- [15] M. Khan and A. Raghav, "Herbal hand wash formulation," Res. J. Pharm. Technol., 2021.
- [16] A. Sharma et al., "Herbal sanitizer review," Asian J. Pharm. Res., 2018.
- [17] S. Chaudhari et al., "Herbal sanitizer formulation," Int. J. Adv. Res. Chem. Sci., 2022.
- [18] B. Siddartha et al., "Evaluation of herbal sanitizer," World J. Pharm. Res., 2023.
- [19] P. Jadhav et al., "Formulation of herbal sanitizer," Int. J. Pharm. Sci., 2025.
- [20] A. Patankar et al., "Comparative study of herbal vs commercial sanitizer," IRJPB, 2023.
- [21] M. E. Aulton, Aulton's Pharmaceutics: Design and Manufacture of Medicines, 2018.
- [22] L. Lachman et al., Theory and Practice of Industrial Pharmacy, 2009.
- [23] L. V. Allen, Pharmaceutical Dosage Forms, 2016.
- [24] H. C. Ansel, Introduction to Pharmaceutical Calculations, 2014.
- [25] M. M. Cowan, "Plant products as antimicrobial agents," Clin. Microbiol. Rev., 1999.
- [26] A. Nostro et al., "Plant extracts antimicrobial activity," Lett. Appl. Microbiol., 2000.
- [27] K. A. Hammer et al., "Essential oils antimicrobial properties," J. Appl. Microbiol., 1999.
- [28] H. N. Bassolé, "Essential oils antimicrobial review," Molecules, 2012.
- [29] C. F. Carson et al., "Tea tree oil antimicrobial activity," Clin. Microbiol. Rev., 2006.
- [30] S. Burt, "Essential oils antibacterial properties," Int. J. Food Microbiol., 2004.

- [31] A. K. Meena et al., "Herbal formulation study," IJPSR, 2020.
- [32] S. Rani et al., "Herbal sanitizer evaluation," Asian J. Pharm. Clin. Res., 2020.
- [33] S. Das et al., "Formulation and evaluation of gel," IJPSR, 2021.
- [34] G. Pandey et al., "Herbal medicine study," J. Herbal Med., 2021.
- [35] S. Verma et al., "Sanitizer formulation," Int. J. Pharm., 2022.
- [36] D. Patel et al., "Pharmaceutical gel formulation," Int. J. Res. Pharm. Sci., 2021.
- [37] S. Shelke et al., "Herbal sanitizer formulation," IJSDR (UGC CARE), 2026.
- [38] P. Gurav et al., "Evaluation of herbal sanitizer," IJRST, 2025.
- [39] P. Rathod et al., "Neem-based antimicrobial study," IJSCI, 2025.
- [40] R. Ankade et al., "Herbal cosmetic formulation," IJIRT (UGC indexed), 2025.