

Management of Gara Visha Through Ayurvedic Principles A Case Report

Dr.Saurabh R. Dhole¹, Dr.Rajendra Urade², Dr. Pankaj B. Sharnangat³, Dr. Lakshmi S. Burange⁴

¹Final year MD student, Department of Agadtantra evum Vidhivaidyak, Bhausaheb Mulak Ayurved Mahavidyalay Nandanwan Nagpur.440024. Maharashtra.

²Guide And Professor, Department of Agadtantra evum Vidhivaidyak, Bhausaheb Mulak Ayurved Mahavidyalay Nandanwan, Nagpur.

^{3,4}Final year MD student, Department of Agadtantra evum Vidhivaidyak , Bhausaheb Mulak Ayurved Mahavidyalay Nandanwan Nagpur. 440024.Maharashtra.

Abstract—Background: Gara Visha is described in Ayurveda as a form of artificially prepared or compounded toxic substance arising from the combination of different materials. Unlike rapidly acting acute poisons, its manifestations may develop gradually and can involve impairment of Agni, Dosha–Dhatu disturbance and systemic symptoms. Ayurvedic management therefore emphasizes identification and avoidance of causative factors along with appropriate Visha Chikitsa, Agni correction and restoration of physiological balance.

Case Presentation: A 34-year-old male presented with recurrent nausea, anorexia, abdominal discomfort, generalized weakness, heaviness, headache, excessive fatigue, disturbed sleep and intermittent skin itching for approximately four months. The symptoms were associated with irregular dietary habits and frequent consumption of incompatible, stale and highly processed food. Ayurvedic examination revealed Agnimandya, Ama-related features and predominantly Kapha-Pitta involvement with Vata association. Based on the history and clinical presentation, the condition was interpreted within the Ayurvedic framework of Gara Visha.

Intervention: Management consisted of Nidana Parivarjana, Deepana-Pachana, mild Koshta Shodhana followed by Shamana treatment. Trikatu Churna, Guduchi, Haridra, Avipattikara Churna and Bilwadi Agada were administered in staged fashion together with Pathya-Apathya regulation.

Outcome: Progressive reduction in gastrointestinal and systemic symptoms was observed over four weeks. Appetite and bowel habits improved, while nausea, heaviness, fatigue, headache and itching substantially decreased. No clinically apparent adverse effects were reported during treatment.

Conclusion: This case illustrates a possible Ayurvedic approach to a chronic symptom complex interpreted as Gara Visha. Correction of Agnimandya, reduction of Ama, elimination of suspected causative exposures and appropriate Visha-oriented Shamana therapy were associated with symptomatic improvement. Controlled clinical research is required before therapeutic efficacy can be inferred.

Index Terms—Gara Visha; Visha Chikitsa; Agada Tantra; Ayurveda; Agnimandya; Ama; Bilwadi Agada; case report.

I. INTRODUCTION

Agada Tantra is the branch of Ayurveda concerned with toxicology and describes the identification, prevention and management of toxic exposures. Ayurvedic literature broadly discusses toxic conditions under concepts such as Sthavara Visha, Jangama Visha and Kritima Visha. Gara Visha occupies an important position among descriptions of artificially produced toxic states.¹

The classical concept of Gara Visha refers to toxicity resulting from combinations of substances which may produce illness after an interval rather than necessarily causing an immediate severe toxic syndrome. Contemporary Ayurvedic scholarship has also explored Gara Visha in relation to complex biological responses to combined substances; such comparisons remain conceptual rather than established biomedical equivalences.²

Modern lifestyles expose individuals to numerous dietary combinations, processed foods, additives, environmental contaminants and inappropriate eating patterns. However, it would be scientifically inappropriate to classify all such exposures automatically as Gara Visha. The Ayurvedic diagnosis requires interpretation of the patient's history, exposure pattern and clinical manifestations.³

A published conceptual analysis describes Garavisha as an artificial poison produced through combinations of substances and discusses a possible conceptual relationship with hapten-carrier interactions and hypersensitivity. Such interpretations provide hypotheses for research but do not establish direct equivalence between Gara Visha and a particular modern toxicological disorder.⁴

The present case describes Ayurvedic management of a chronic gastrointestinal and constitutional symptom complex interpreted clinically through the framework of Gara Visha.

II. AIM

To document the clinical application of Ayurvedic principles in the management of a patient presenting with symptoms interpreted as Gara Visha.

III. OBJECTIVE

1. To assess the clinical manifestations according to Ayurvedic principles of Visha, Dosha, Agni and Ama.
2. To formulate a treatment protocol based on Nidana Parivarjana, Deepana-Pachana, Koshta Shodhana and Shamana Chikitsa.
3. To document changes in major clinical symptoms during four weeks of management.

IV. CASE PRESENTATION

A 34-year-old male patient attended the outpatient department with complaints of reduced appetite, recurrent nausea, abdominal discomfort, generalized weakness and heaviness for approximately four months.

He additionally reported intermittent headache, excessive fatigue, disturbed sleep, occasional itching and an unpleasant taste in the mouth.

The symptoms initially occurred intermittently but had gradually increased during the preceding two months.

4.1 Chief Complaints

Complaint	Duration
Loss of appetite	4 months
Generalized weakness	4 months
Abdominal discomfort	3 months
Nausea	3 months
Heaviness of body	3 months
Headache	2 months
Intermittent itching	2 months
Disturbed sleep	2 months

4.2 History of Present Illness

The patient was apparently healthy approximately four months before presentation. His occupational schedule resulted in irregular meal timing, frequent skipping of breakfast and late-night meals.

He reported frequent intake of reheated food, packaged snacks, restaurant food, excessively spicy preparations and irregular combinations of dairy products with sour or salty foods.

Gradually, he developed reduced appetite and abdominal heaviness after meals. These symptoms were subsequently accompanied by nausea, fatigue, headache and occasional itching.

No history suggestive of an immediately life-threatening poisoning episode was reported.

V. PAST HISTORY

There was no known history of diabetes mellitus, hypertension, tuberculosis, chronic hepatic disease, chronic renal disease or major gastrointestinal disorder.

No major surgical history was reported.

The patient denied known drug allergy.

VI. PERSONAL HISTORY

Parameter	Finding
Appetite	Reduced
Diet	Mixed
Dietary pattern	Irregular
Bowel	Occasionally constipated
Micturition	Normal
Sleep	Disturbed
Addiction	Nil significant
Exercise	Minimal
Occupational activity	Predominantly sedentary

VII. GENERAL EXAMINATION

The patient was conscious, cooperative and oriented.

Parameter	Finding
Pulse	78/min
Blood pressure	118/76 mmHg
Respiratory rate	16/min
Temperature	Afebrile
Pallor	Absent
Icterus	Absent
Cyanosis	Absent
Clubbing	Absent
Pedal edema	Absent
Lymphadenopathy	Absent

Systemic examination did not reveal significant cardiovascular or respiratory abnormalities. Mild nonspecific epigastric discomfort was elicited without guarding or rigidity.

VIII. ASHTAVIDHA PARIKSHA

Pariksha	Observation
Nadi	Kapha-Pitta predominance
Mutra	Samyak
Mala	Occasional Vibandha
Jihva	Sama, mildly coated
Shabda	Prakrita
Sparsha	Anushna
Drik	Prakrita
Akriti	Madhyama

IX. DASHAVIDHA PARIKSHA

Parameter	Assessment
Prakriti	Pitta-Kapha
Vikriti	Kapha-Pitta with Vata association
Sara	Madhyama
Samhanana	Madhyama
Pramana	Madhyama
Satmya	Madhyama
Satva	Madhyama
Ahara Shakti	Avara
Vyayama Shakti	Avara-Madhyama
Vaya	Madhyama

The most significant findings were Mandagni, reduced Ahara Shakti and features suggestive of Sama Dosha.

X. DIAGNOSTIC ASSESSMENT

Routine investigations were considered primarily to exclude common biomedical causes of chronic fatigue and gastrointestinal symptoms.

Illustrative baseline investigations

Investigation	Result
Hemoglobin	13.4 g/dL
Total leukocyte count	7,600/mm ³
Platelet count	2.45 lakh/mm ³
Fasting blood glucose	92 mg/dL
Serum creatinine	0.9 mg/dL
Total bilirubin	0.7 mg/dL
AST	28 U/L
ALT	31 U/L
TSH	2.6 mIU/L

No major abnormality was identified in the screening investigations.

Potential biomedical differential diagnoses included functional dyspepsia, gastritis, food intolerance, nutritional deficiency, hepatobiliary disease and other causes of chronic constitutional symptoms.

Because Gara Visha is an Ayurvedic diagnostic construct rather than a laboratory-confirmed biomedical toxicological diagnosis, the Ayurvedic assessment was made primarily from the history and symptom complex.

XI. AYURVEDIC DIAGNOSTIC CONSIDERATION

The following clinical characteristics supported interpretation through the Gara Visha framework:

- Chronic and gradually developing symptom pattern
- Mandagni
- Aruchi
- Avipaka-like gastrointestinal disturbance
- Gaurava
- Daurbalya
- Shiroruja
- Intermittent Twak-related manifestations
- History of repeated unsuitable dietary exposure
- Sama Dosha features


Ayurvedic diagnosis

Gara Visha-associated clinical state with Agnimandya and Ama predominance involving Kapha-Pitta with secondary Vata association.

XII. SAMPRAPTI

Repeated intake of unsuitable food and incompatible dietary combinations was considered the principal Nidana in this case.

The proposed pathological sequence was:



XIII. SAMPRAPTI GHATAKA

Component	Assessment
Dosha	Kapha-Pitta predominant; Vata Anubandha
Dushya	Rasa, Rakta
Agni	Jatharagni Mandya
Ama	Present
Srotas	Annavaha, Rasavaha, Raktavaha
Srotodushti	Sanga
Udbhava Sthana	Amashaya
Adhithana	Koshta with systemic involvement
Rogamarga	Abhyantara
Vyadhi Swabhava	Chirakari

XIV. CHIKITSA SIDDHANTA

The therapeutic strategy was formulated according to the following sequence:

Nidana Parivarjana → Deepana → Pachana → Koshta Shuddhi → Visha-oriented Shamana → Agni maintenance → Pathya regulation

Treatment was individualized according to Bala, Agni, Koshta and clinical presentation.

XV. THERAPEUTIC INTERVENTION

Phase I: Deepana-Pachana

Duration: Day 1–5

Medicine	Illustrative dose	Anupana	Frequency
Trikatu Churna	1 g	Lukewarm water	Twice daily before food
Guduchi Churna	3 g	Lukewarm water	Twice daily

The initial objective was to improve Agni and reduce Sama features before administering the principal Shamana regimen.

Phase II: Mridu Koshta Shodhana

On the basis of Koshta and Bala, mild bowel purification was planned rather than intensive Panchakarma.

Avipattikara Churna – 5 g at bedtime with lukewarm water for three nights.

The patient subsequently reported easier bowel evacuation and reduction in abdominal heaviness.

Phase III: Shamana and Visha-oriented Management

Duration: Day 6–28

Medicine	Illustrative dose	Anupana	Frequency
Bilwadi Agada	500 mg	Honey	Twice daily
Guduchi Churna	3 g	Lukewarm water	Twice daily
Haridra Churna	2 g	Lukewarm water	Twice daily
Avipattikara Churna	3–5 g	Lukewarm water	At bedtime when required

The medicines and doses shown here represent the constructed case protocol and should not be interpreted as a universal treatment regimen.

XVI. PATHYA-APATHYA

The patient was advised to consume freshly prepared, warm and easily digestible meals.

Light preparations such as Mudga Yusha and other easily digestible foods were preferred during the initial phase.

He was instructed to maintain regular meal timing and avoid overeating.

The following were restricted:

- Stale and repeatedly reheated foods
- Excessively fried foods
- Highly processed packaged foods
- Excessive sour and spicy foods
- Irregular food combinations
- Heavy meals at night
- Eating before complete digestion of the previous meal
- Alcohol and other potential intoxicants

Adequate sleep, mild physical activity and stress reduction were also advised.

XVII. CLINICAL TIMELINE

Time	Clinical event
~4 months before consultation	Reduced appetite and generalized weakness began
~3 months before consultation	Nausea, abdominal discomfort and heaviness developed
~2 months before consultation	Headache, sleep disturbance and intermittent itching developed
Day 0	Ayurvedic and general clinical assessment
Day 1–5	Deepana-Pachana initiated
Day 6	Shamana/Visha-oriented treatment initiated
Day 14	Appetite and nausea improved
Day 21	Marked reduction in heaviness and fatigue
Day 28	Major symptoms substantially reduced
Follow-up	Dietary regulation and Agni maintenance advised

XVIII. ASSESSMENT CRITERIA

Symptoms were graded using a simple four-point clinical scale:

0 – Absent

1 – Mild

2 – Moderate

3 – Severe

Symptom	Baseline	Day 14	Day 28
Loss of appetite	3	1	0
Nausea	2	1	0
Abdominal discomfort	2	1	0
Generalized weakness	3	2	1
Body heaviness	3	1	0
Headache	2	1	0
Itching	2	1	0
Disturbed sleep	2	1	1

The largest improvements were observed in appetite, nausea, abdominal discomfort and heaviness.

Generalized weakness and sleep disturbance improved more gradually.

No clinically apparent adverse drug reaction was reported during the observational period.

XIX. DISCUSSION

Gara Visha represents an important toxicological concept described under Agada Tantra. Unlike an immediately acting severe poison, the conceptual framework is particularly relevant to insidious clinical manifestations attributed to repeated or compounded exposure. 5

The term should nevertheless be used carefully in contemporary clinical reporting. Symptoms such as anorexia, nausea, weakness, heaviness, headache and itching are nonspecific and may arise from numerous biomedical disorders. Therefore, exclusion of serious hepatic, renal, hematological, metabolic, infectious or gastrointestinal pathology is important before attributing a chronic symptom complex to an Ayurvedic diagnostic category. 6

In the present case, irregular dietary behavior and repeated consumption of unsuitable foods were associated temporally with gradual development of

gastrointestinal and systemic symptoms. From the Ayurvedic perspective, these exposures were considered to disturb Agni.⁷

Role of Agnimandya and Ama

Agni occupies a central position in Ayurvedic physiology and pathology. Persistent impairment of digestive capacity can lead to improper transformation of ingested material and the development of Ama.⁸

The coated tongue, reduced appetite, postprandial heaviness, nausea and fatigue in this patient were clinically interpreted as manifestations associated with Mandagni and Ama.

Consequently, directly administering heavy or extensive formulations without first correcting digestion was considered inappropriate.⁹

Rationale for Deepana-Pachana

Trikatu was selected during the initial phase for its traditional Deepana-Pachana application.

The objective was not to treat an assumed biomedical poisoning directly but to improve digestive function according to Ayurvedic assessment and prepare the patient for subsequent management.

Guduchi was incorporated because of its traditional application in conditions involving Jwara, Pitta disturbance and systemic morbidity and because it is frequently employed as a Rasayana and supportive drug in Ayurvedic practice.

Role of Mridu Shodhana¹⁰

After improvement in appetite and reduction in Sama features, mild bowel purification was employed.

The intention was to facilitate Koshta Shuddhi while avoiding excessive depletion in a patient already reporting weakness.

Avipattikara Churna was selected because the clinical picture contained Pitta-associated gastrointestinal symptoms along with bowel irregularity.

Role of Bilwadi Agada¹¹

Bilwadi Agada is a classical Agada formulation traditionally used in Visha-related clinical contexts. Its selection in the present case was based on the Ayurvedic interpretation of the symptom complex rather than confirmation of a particular chemical toxin. This distinction is important when presenting traditional toxicological concepts in contemporary scientific literature.

Role of Haridra¹²

Haridra was incorporated particularly because of intermittent skin symptoms and the Kapha-Pitta/Rakta involvement considered in the Ayurvedic assessment. Importance of Nidana Parivarjana¹³

Avoidance of suspected causative exposures was arguably one of the most important components of management.

The patient was specifically instructed to discontinue stale, reheated, highly processed and incompatible dietary patterns and adopt regular freshly prepared meals.

Thus, the clinical improvement cannot be attributed to the medicines alone. Dietary modification, normalization of meal timing, spontaneous fluctuation of symptoms and nonspecific effects may all have contributed.

XX.POSSIBLE MODE OF ACTION ACCORDING TO AYURVEDIC PRINCIPLES

The proposed therapeutic sequence can be summarized as:

Deepana

→ improvement of Agni

Pachana

→ reduction of Ama

Mridu Shodhana

→ Koshta Shuddhi and facilitation of Dosha elimination

Agada-based Shamana

→ management within the traditional Visha Chikitsa framework

Kapha-Pitta Shamaka measures

→ correction of predominant Dosha disturbance

Nidana Parivarjana

→ prevention of continued exposure

Pathya

→ maintenance of Agni and reduction of recurrence.

Collectively, these interventions were intended to achieve Agni Deepana, Ama Pachana, Dosha Shamana, Srotoshodhana and restoration of physiological balance.

XXI. CONTEMPORARY PERSPECTIVE

Attempts have been made in Ayurvedic scholarship to understand Gara Visha through contemporary immunological and toxicological concepts.

For example, a published conceptual paper compared the Ayurvedic description of Garavisha with hapten-carrier interactions and hypersensitivity mechanisms. Such interpretations may provide a basis for hypothesis generation, but they should not be treated as proof that Gara Visha is identical to hypersensitivity or any single biomedical diagnosis.

Similarly, chronic exposure to food additives, environmental pollutants or chemical mixtures may produce biological effects, but a direct equivalence between these phenomena and Gara Visha has not been established.

The scientific value of the traditional concept may therefore lie in generating research questions concerning repeated low-level exposures, interactions between substances, individual susceptibility and delayed manifestations.

XXII. STRENGTHS OF THE CASE

The case demonstrates a structured Ayurvedic assessment integrating Nidana, Dosha, Dushya, Agni, Ama and Srotas.

Treatment was administered sequentially rather than using multiple interventions simultaneously from the first day.

Dietary causative factors were specifically addressed through Nidana Parivarjana.

Symptom progression was documented at defined follow-up intervals.

XXIII. LIMITATIONS

This is a single case observation and cannot establish efficacy or causality.

The symptoms attributed to Gara Visha were nonspecific.

No specific chemical toxicant was identified or quantitatively measured.

Multiple interventions were used; therefore, the individual therapeutic contribution of each medicine cannot be determined.

Dietary and lifestyle modification may have contributed substantially to the outcome.

The symptom scale used was clinician-developed rather than a validated patient-reported outcome measure.

Long-term follow-up was limited.

Accordingly, the clinical response should be considered hypothesis-generating rather than confirmatory evidence.

XXIV. PATIENT PERSPECTIVE

The patient reported that abdominal heaviness and nausea began decreasing during the first two weeks of treatment. He perceived improvement in appetite and daily energy and found adherence to freshly prepared meals particularly helpful. At the end of four weeks, he reported that most of his initial symptoms had either disappeared or become mild and expressed willingness to continue the advised dietary regimen. For publication, this paragraph should be replaced by the actual patient's own perspective.

XXV. CONCLUSION

This case illustrates a structured Ayurvedic approach to a chronic gastrointestinal and constitutional symptom complex interpreted through the concept of Gara Visha. Management focused on Nidana Parivarjana, correction of Agnimandya, Ama Pachana, mild Koshta Shodhana and subsequent Visha-oriented Shamana therapy. Clinically meaningful symptomatic improvement was observed during the four-week period in this constructed case scenario. However, because this is a single-case framework involving multiple simultaneous interventions, no causal conclusion regarding efficacy can be drawn. Future research should develop clearly defined diagnostic criteria for Gara Visha, standardized outcome measures and prospective clinical protocols. Investigation of traditional Gara Visha concepts alongside modern toxicology may also help determine whether particular aspects have measurable biomedical correlates.

XXVI. DECLARATION OF PATIENT CONSENT

For an actual submission, the authors should certify that appropriate written informed consent was obtained from the patient for publication of anonymized clinical information. Identifying details should not be disclosed.

XXVII. ETHICAL CONSIDERATION

Patient confidentiality should be maintained throughout the report. Institutional requirements regarding ethics committee review or exemption for case reports should be followed before submission.

XXVIII. CONFLICT OF INTEREST

The authors declare no conflict of interest.

XXIX. SOURCE OF FUNDING

Nil

REFERENCES

- [1] Government of India, Ministry of AYUSH. The Ayurvedic Pharmacopoeia of India. New Delhi: Government of India.
- [2] Government of India, Ministry of AYUSH. The Ayurvedic Formulary of India. New Delhi: Government of India.
- [3] Datta S, et al. Physiological concept of hapten-carrier adduct vis-à-vis Garavisha. AYU. 2018; 38:37–41.
- [4] Gagnier JJ, Kienle G, Altman DG, Moher D, Sox H, Riley DS, CARE Group. The CARE guidelines: consensus-based clinical case reporting guideline development. J Clin Epidemiol. 2014;67(1):46–51.
- [5] Riley DS, Barber MS, Kienle GS, Aronson JK, von Schoen-Angerer T, Tugwell P, et al. CARE guidelines for case reports: explanation and elaboration document. J Clin Epidemiol. 2017; 89:218–235.
- [6] Government of India, Ministry of Health and Family Welfare, Department of AYUSH. The Ayurvedic Pharmacopoeia of India. Part I. Vol 1. New Delhi: Department of AYUSH; 2001.
- [7] Government of India, Ministry of Health and Family Welfare, Department of AYUSH. The Ayurvedic Pharmacopoeia of India. Part I. Vol 2. New Delhi: Department of AYUSH; 2001.
- [8] Government of India, Ministry of Health and Family Welfare, Department of AYUSH. The Ayurvedic Formulary of India. Part I. 2nd ed. New Delhi: Department of AYUSH; 2003.
- [9] Government of India, Ministry of AYUSH. The Ayurvedic Formulary of India. Part II. 2nd ed. New Delhi: Ministry of AYUSH; 2011.
- [10] Datta S, Das B, Sen S. Physiological concept of hapten-carrier adduct vis-à-vis Garavisha. AYU. 2017;38(1-2):37-41.
- [11] Patwardhan B, Warude D, Pushpangadan P, Bhatt N. Ayurveda and traditional Chinese medicine: a comparative overview. Evid Based Complement Alternat Med. 2005;2(4):465-73.
- [12] Patwardhan B, Vaidya ADB, Chorghade M. Ayurveda and natural products drug discovery. Curr Sci. 2004;86(6):789-99.
- [13] Patwardhan B, Mashelkar RA. Traditional medicine-inspired approaches to drug discovery: can Ayurveda show the way forward? Drug Discov Today. 2009;14(15-16):804-11.