

Role Of Homoeopathic Management in Recurrent Tonsillitis

Dr. Shivangi Meva¹, Dr. Vimal Rachchh², Dr. Ashish Sapariya³

¹MD Homoeopathy Part- I Scholar, Department of Organon of Medicine, B.G. Garaiya Homoeopathic Medical College and Hospital, Saurashtra University, Rajkot, Gujarat, India

^{2,3}MD Hom Associated Professor, Department of Organon of Medicine, B.G. Garaiya Homoeopathic Medical College and Hospital, Saurashtra University, Rajkot, Gujarat, India

Abstract—Recurrent tonsillitis is characterized by repeated episodes of inflammation and infection of the palatine tonsils and is particularly common among children and adolescents. It can lead to recurrent sore throat, fever, dysphagia, cervical lymphadenopathy, tonsillar enlargement, absence from school or work, and repeated courses of antibiotics. The condition may be associated with viral or bacterial infections, particularly Group A Streptococcus, as well as environmental and host-related factors. Conventional management includes symptomatic treatment, appropriate antimicrobial therapy when bacterial infection is suspected or confirmed, and tonsillectomy in selected patients with sufficiently frequent or severe episodes. Homoeopathy is used as a complementary system of medicine with the aim of individualizing treatment according to the patient's total symptom picture and constitutional characteristics. Traditional homoeopathic practice commonly considers. However, the clinical evidence remains limited. **Aim:** The aim of this article is to describe the role of homoeopathic management in recurrent tonsillitis, discuss the possible causes and predisposing factors, outline commonly described homoeopathic remedies used according to the individual symptom picture, and review the available evidence regarding homoeopathy as a complementary approach in recurrent tonsillitis disease.

Index Terms—Recurrent tonsillitis, tonsillitis, homoeopathy, homoeopathic remedies, Belladonna, Hepar sulphuris, Mercurius solubilis, Phytolacca, Baryta carbonica, Calcarea carbonica, recurrent throat infection, tonsillar inflammation.

I. INTRODUCTION

Tonsillitis refers to inflammation of the palatine tonsils and is a common condition, particularly in children and young adults. It may occur as an isolated

acute episode or may become recurrent when episodes occur repeatedly over time. Patients may present with sore throat, painful swallowing, fever, enlarged or erythematous tonsils, exudation, headache, malaise and enlarged cervical lymph nodes. Most acute episodes are caused by viruses, while bacterial infection, particularly Group A streptococcal infection, accounts for an important proportion of cases. Nnhs.com.in

Recurrent tonsillitis may have a considerable effect on quality of life because repeated episodes can result in school or occupational absenteeism, sleep disturbance, repeated healthcare visits and repeated exposure to antimicrobial therapy. Management therefore requires not only treatment of the individual acute episode but also assessment of the frequency, severity and documentation of recurrent episodes. In selected children, tonsillectomy may be considered when there are at least seven documented episodes in one year, five episodes annually for two consecutive years, or three episodes annually for three consecutive years, with appropriate clinical documentation.

II. CAUSES AND PREDISPOSING FACTORS OF RECURRENT TONSILLITIS

Incomplete resolution of previous infection may contribute to repeated symptoms, although recurrent episodes should not automatically be interpreted as persistence of the same infection. Some individuals are particularly susceptible to repeated upper respiratory infections because of age, exposure patterns and individual host factors. Children attending school or daycare may experience frequent respiratory infections because of close contact with other children.

Environmental factors can also contribute to recurrent throat complaints. Exposure to cigarette smoke, air pollution, dust and other respiratory irritants may aggravate upper-airway symptoms. Poor hand hygiene and close contact with individuals experiencing respiratory infections can facilitate transmission.

Anatomical and local factors may also play a role. Enlarged tonsils, deep tonsillar crypts and accumulation of debris may predispose some individuals to persistent or recurrent local symptoms. Associated nasal disease, allergic rhinitis and chronic mouth breathing can contribute to throat irritation and may coexist with recurrent tonsillar disease.

Host susceptibility and associated conditions are also important considerations. Nutritional status, sleep, general health and coexisting upper respiratory tract problems may influence an individual's susceptibility to infection. However, recurrent sore throat does not always represent recurrent bacterial tonsillitis, and appropriate clinical assessment is important before attributing every episode to tonsillar infection.

III. HOMOEOPATHIC MANAGEMENT OF RECURRENT TONSILLITIS

The traditional homoeopathic approach emphasizes individualization of treatment. The remedy is selected according to the characteristic symptom picture rather than simply prescribing one medicine for every patient with recurrent tonsillitis. The following remedies are commonly described in homoeopathic materia medica for tonsillar and throat complaints.

1. Belladonna

Belladonna is traditionally considered when tonsillitis has a sudden and intense onset. The throat may appear bright red, swollen and inflamed, with marked sensitivity and fever. The patient may experience throbbing pain and difficulty swallowing. It is traditionally associated with acute inflammatory presentations characterized by rapid onset and prominent fever.

2. Hepar sulphuris calcareum

Hepar sulphuris is traditionally considered in patients with marked sensitivity of the throat and tonsils. Throat pain may be severe, particularly on swallowing, and the patient may be sensitive to cold air, cold drinks or exposure to cold. Recurrent

suppurative tendencies are traditionally emphasized in its materia-medica indications.

3. Mercurius solubilis

Mercurius solubilis is traditionally considered when tonsillar inflammation is accompanied by excessive salivation, offensive breath, coated tongue and difficulty swallowing. The tonsils may be enlarged and inflamed, sometimes with exudation. A tendency for symptoms to be worse at night is also traditionally described.

4. Baryta carbonica

Baryta carbonica is traditionally considered in patients with a tendency toward recurrent enlargement and inflammation of the tonsils, particularly in children. Repeated tonsillar problems, enlarged tonsils and susceptibility to recurrent throat infections are characteristic indications described in homoeopathic materia medica.

5. Calcarea carbonica

Calcarea carbonica is traditionally considered as a constitutional remedy in individuals with a recurring tendency toward tonsillar and respiratory infections. In homoeopathic practice, its selection is based on the broader constitutional symptom picture rather than recurrent tonsillitis alone.

6. Silicea

Silicea is traditionally considered when there is a recurrent tendency toward suppurative processes and chronic or repeated tonsillar complaints. It may be selected when the overall constitutional picture corresponds to the remedy.

IV. ROLE OF HOMOEOPATHY IN RECURRENT TONSILLITIS

The proposed role of homoeopathy in recurrent tonsillitis is primarily that of an individualized complementary approach aimed at addressing the patient's recurrent symptom pattern. A recent systematic review published in 2026 evaluated homeopathy specifically for tonsillitis and identified five randomized controlled trials. The review reported that some individual studies found reductions in recurrence, antibiotic consumption and improvement in quality of life; however, the studies differed

substantially in their populations, interventions and outcome measures, preventing reliable meta-analysis. This finding is important because it indicates that there is some research activity in this area, but it does not establish homoeopathy as a proven treatment for recurrent tonsillitis. Earlier Cochrane evidence examining homeopathic medicines for acute respiratory tract infections in children found little or no consistent benefit compared with placebo and concluded that the evidence did not support their efficacy for acute respiratory infections.

V. CONCLUSION

Recurrent tonsillitis is a common and clinically important condition that can significantly affect quality of life, particularly in children and adolescents. Its causes include viral and bacterial infections, repeated exposure to infectious agents, environmental factors, local anatomical factors and individual susceptibility.

Current evidence concerning homoeopathy for tonsillitis is evolving. Therefore, further well-designed randomized controlled trials with standardized outcome measures are required before firm conclusions can be made regarding the efficacy of homoeopathic treatment for recurrent tonsillitis. Homoeopathy, when used, should be regarded as complementary rather than a replacement for evidence-based medical care, particularly where bacterial infection, airway compromise or indications for tonsillectomy are present.

REFERENCES

- [1] Dhingra, P. L., and S. Dhingra, *Diseases of Ear, Nose and Throat & Head and Neck Surgery*, 7th ed. New Delhi: Elsevier, 2018.
- [2] Kumar, P., and M. Clark, *Kumar & Clark's Clinical Medicine*, 10th ed. Elsevier, 2020.
- [3] Boericke, W., *Pocket Manual of Homoeopathic Materia Medica & Repertory*. New Delhi: B. Jain Publishers, 2007.
- [4] Allen, H. C., *Keynotes and Red Line Symptoms of the Materia Medica with Nosodes*. New Delhi: B. Jain Publishers, 2002.
- [5] Kent, J. T., *Repertory of the Homoeopathic Materia Medica*. New Delhi: B. Jain Publishers, 2004.