

Role Of Concomitant Symptoms in Final Remedy Selection: A Prospective Repertorial Study

Dr. Dhruvi Maru¹, Dr. Nirav Manseta², Dr. Prabhat Vaghmashi³

¹PG Scholar (Part-1) Department of Repertory and Case Taking, Shree B. G. Garaiya Homoeopathic Medical College, Rajkot

²Associate Professor, Department of Repertory and Case Taking, Shree B. G. Garaiya Homoeopathic Medical College, Rajkot

³Professor & HOD. Department of Repertory and Case Taking, Shree B. G. Garaiya Homoeopathic Medical College, Rajkot

Abstract—Background: In homoeopathic analysis, concomitant symptoms are symptoms that alongside the principal complaint and may provide additional information for individualization. This article outlines a prospective repertorial study designed to examine whether documenting and evaluating concomitant symptoms influences the final remedy selected after repertorization.

Methods: Consecutive eligible cases would be assessed using a predefined case-taking form, with the chief complaint and concomitant symptoms recorded before repertorial analysis. Remedy selection based on the complete symptom picture would be compared with a selection made from the chief complaint alone. Agreement between selections, changes in remedy ranking, and the contribution of concomitants to the final decision would be documented.

Case presentation, treatment and follow-up: The protocol includes standardized documentation of an individual case from baseline assessment through individualized homoeopathic prescribing and scheduled follow-up.

Because no patient dataset or completed study results were supplied, these sections describe the planned documentation framework rather than fabricated clinical findings. **Conclusion:** This manuscript is presented study protocol and conceptual article rather than a report of completed clinical findings. A structured approach may support transparent and reproducible assessment of the role of concomitant symptoms in repertorial decision-making

Index Terms—Concomitant symptoms, homoeopathy, repertorization, remedy selection, prospective study, case analysis.

I. INTRODUCTION

Homoeopathic case-taking commonly considers the patient's chief complaint together with accompanying symptoms, modalities, sensations, individual characteristics. A concomitant symptom is generally understood appears as a symptom that in association with the main complaint but is not necessarily a direct or expected consequence of it. In repertorial analysis, such symptoms may help distinguish between remedies that cover the principal complaint similarly.

The practical challenge is deciding which concomitants are sufficiently clear, characteristic, and reliable to influence the final prescription. Including every reported symptom may add noise, while excluding a meaningful accompanying symptom may reduce the distinctiveness of the totality. A structured protocol can make this decision process more transparent and allow the influence of concomitants to be assessed systematically.

Aims and objectives

The primary examines how the inclusion of concomitant symptoms affects final remedy selection during repertorization. Secondary objectives are to: (1) measure how often the leading remedy changes when concomitants are included; (2) compare remedy rankings from chief-complaint-only and complete-symptom analyses; and (3) assess agreement between independent evaluators using a predefined rubric.

II. METHODOLOGY

Study design

A prospective observational repertorial study is proposed. The protocol should be finalized before recruitment and reported in accordance with applicable institutional and ethical requirements. The study setting, recruitment period, sample size, and analysis plan should be specified by the research team before data collection.

Participants and eligibility

Eligible participants may be consecutive patients presenting with a clearly documented chief complaint for which a complete case record can be obtained. Inclusion and exclusion criteria should be set in advance, with the rationale recorded. Participation should be voluntary, and informed consent and confidentiality safeguards should be applied as required. This repertorial study should not delay or replace appropriate evidence-based medical assessment or treatment.

Data collection and symptom classification

A standardized case-record form should capture the chief complaint, onset and duration, location, sensation, modalities, associated symptoms, relevant medical history, and any red-flag features requiring clinical referral. Concomitant symptoms should be defined operationally before the study begins. Each symptom should be recorded in the patient's own words where possible, then classified according to the chosen repertory's terminology. The repertory edition and software, if any, should be documented.

Repertorial procedure

For each case, two repertorial analyses are proposed: (A) a chief-complaint analysis using prespecified principal symptoms; and (B) a complete analysis including qualifying concomitant symptoms. The order of analyses should be randomized or otherwise controlled to reduce carry-over effects. Where feasible, two trained evaluators should repertorize cases independently while blinded to each other's selections. Remedy coverage, rubric weighting, remedy rank, and the rationale for the final selection should be recorded using the same predefined rules in both conditions.

Table 1. Proposed methodology for evaluating the contribution of concomitant symptoms.

Component	Planned approach
Design	Prospective observational repertorial study.
Participants	Consecutive eligible cases with a complete case record and consent as applicable.
Primary comparison	Chief-complaint-only repertorization versus complete-symptom repertorization including qualifying concomitants.
Main documentation	Rubrics used, remedy ranking, selected remedy, and reason for final selection.
Evaluator control	Independent repertorization by trained evaluators where feasible; order controlled to reduce carry-over.
Outcome focus	Frequency of remedy change, change in ranking, and contribution of concomitants to the final decision.

III. CASE PRESENTATION

The supplied article did not contain an actual patient case record or completed clinical dataset. To avoid presenting invented patient information as research findings, the following is a standardized case-presentation framework to be completed from the enrolled patient's original case record.

Table 2. Case-presentation framework; no patient-specific values have been fabricated.

ITEM	Information to be documented
Patient identification	Study ID only; age/sex and other non-identifying demographic information as approved in the protocol.
Chief complaint	Exact complaint, location, onset, duration, character/sensation, severity and relevant modalities.
Concomitant symptoms	Symptoms occurring alongside the chief complaint, documented in the patient's own words and assessed for clarity, individuality and reliability.
Associated history	Relevant past history, family history, personal history, maintaining factors and clinically relevant investigations.

Clinical assessment	General and local examination, working diagnosis/clinical assessment and referral or investigation where indicated.
Repertorial analysis	Chief-complaint rubrics, complete-set rubrics including concomitants, remedy ranking and final prescribing rationale.

IV. TREATMENT/PRESCRIBING PROCEDURE

The supplied manuscript describes a repertorial decision-making study and does not provide an actual treatment record. Therefore, treatment is presented here as the planned prescribing procedure rather than as a completed clinical intervention.

After case-taking and clinical assessment, the case would be analyzed using the predefined repertorial method. One analysis would use the chief complaint and prespecified principal symptoms, while the second would include qualifying concomitant symptoms. The final homoeopathic prescription would be individualized according to the approved clinical protocol and the complete case record. Remedy name, potency, dose, repetition, date of prescription, and any relevant advice should be recorded prospectively.

The protocol should clearly separate repertorial evidence from the prescriber's clinical judgment. Any change in remedy, potency or repetition during follow-up should be documented with its stated rationale. Appropriate conventional investigation, referral or co-management should be undertaken whenever clinically indicated.

V. FOLLOW-UP

Follow-up should be scheduled prospectively according to the study protocol. At each visit, the investigator should record the status of the chief complaint, the previously identified concomitant symptoms, any new symptoms, changes in intensity or frequency, adherence to the prescribed plan, adverse events, and any intercurrent treatment.

The same assessment method should be used at baseline and follow-up wherever possible so that changes are comparable. The repertorial analysis and the final remedy selected at baseline should not be retrospectively altered merely to fit the clinical outcome. If the remedy changes during follow-up, the

date and reason for the change should be recorded separately

Table 3. Proposed follow-up documentation schedule; exact visit intervals should be fixed in the approved protocol.

Time point	Minimum documentation
Baseline	Chief complaint, concomitants, examination/clinical assessment, repertorial analyses, selected remedy and baseline severity measure where applicable.
First follow-up	Change in chief complaint and concomitants, adherence, adverse events/intercurrent treatment and any prescribing decision.
Subsequent follow-ups	Repeat standardized assessment, symptom trajectory, remedy/potency changes with rationale and clinically relevant investigations/referrals.
Final assessment	Overall clinical status, documented change from baseline, final repertorial decision and completeness of follow-up.

VI. OUTCOME MEASURES AND DATA ANALYSIS

The primary outcome should be the proportion of cases in which the final leading remedy changes when qualifying concomitant symptoms are added to the repertorial analysis. Secondary outcomes may include the change in remedy rank, the number of additional rubrics contributed by concomitants, agreement between evaluators, and the proportion of cases in which a concomitant materially influences the prescribing rationale.

For completed data, descriptive statistics should be used to summarize participant characteristics and repertorial findings. Categorical outcomes may be presented as frequencies and percentages, while continuous measures may be summarized using appropriate measures of central tendency and dispersion. Any inferential analysis should be prespecified before data collection and should match the sample size and study design.

VII. ETHICAL AND REPORTING CONSIDERATIONS

Participant confidentiality, informed consent where applicable, and institutional/ethics requirements should be addressed before recruitment. Any clinically significant finding requiring investigation or referral should be managed appropriately. The final manuscript should distinguish protocol methods from actual study results and should not report unobserved outcomes.

VIII. LIMITATIONS

The supplied manuscript is a protocol/conceptual article and contains no patient dataset or completed study results. Consequently, it cannot establish the frequency or magnitude of remedy changes caused by concomitant symptoms. Repertorial decisions may also be influenced by rubric selection, repertory edition, evaluator judgment and the completeness of the case record. These limitations should be considered when interpreting future findings.

IX. CONCLUSION

A structured prospective approach can make the contribution of concomitant symptoms to repertorial decision-making more transparent. Recording the chief complaint separately from qualifying concomitants, performing predefined comparative repertorization, documenting the prescribing rationale, and maintaining standardized follow-up may provide a reproducible framework for future clinical research. Actual conclusions about effectiveness or clinical benefit should be based only on completed, appropriately analyzed patient data.

X. CONCEPTUAL GRAPHICAL ANALYSIS

Figure 1 presents the proposed comparison between repertorial analysis based on the chief complaint alone and analysis incorporating qualifying concomitant symptoms. Both pathways use the same standardized case record and predefined recording rules. The comparison focuses on whether the leading remedy or its ranking changes, and on evaluator agreement.

CONCEPTUAL FRAMEWORK FOR COMPARING REPERTORIAL ANALYSES




Figure 1. Conceptual workflow for comparing chief-complaint-only and complete-symptom repertorization. This is a study framework, not patient-level data or a result graph.

Interpretation

The proposed design allows the contribution of concomitant symptoms to be documented transparently. The direction and magnitude of any change in remedy selection can only be determined after verified study data are collected.

REFERENCES

- [1] J. T. Kent, *Lectures on Homoeopathic Philosophy*. Chicago, IL, USA: Ehrhart & Karl, 1905.
- [2] S. Hahnemann, *Organon of Medicine*, 6th ed. [Publisher, place of publication, and translator details to be completed according to the exact edition consulted].
- [3] F. Schroyens, ed., *Synthesis: Repertorium Homeopathicum Syntheticum*. [Edition, place of publication, and publisher to be completed according to the exact edition consulted].
- [4] World Medical Association, "World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects," *JAMA*, vol. 310, no. 20, pp. 2191–2194, 2013, doi: 10.1001/jama.2013.281053.