Ayurveda, Tyrosine, and Copper: An Integrative Biochemical–Ayurvedic Perspective

Dr. Rushikesh Kangad[1], Dr. Vikram Singh[2], Dr. Anbuselvam A[3]
[1]*Associate Professor, Department of Shalya Tantra, EVA College of Ayurved, Supedi*
[2]*Assistant Professor, Department of Shalya Tantra, Parul Institute of Ayurveda, Vaghodiya, Baroda*
[3]*Assistant Professor, Department of Kayachikitsa, EVA College of Ayurved, Supedi*

***Abstract*— Background: Ayurveda emphasizes equilibrium between dietary inputs, trace minerals, and neuropsychological health, while modern biochemistry recognizes amino acids and micronutrients as critical regulators of enzymatic and neuroendocrine pathways. Objective: To critically examine the biochemical role of tyrosine and copper and correlate these mechanisms with Ayurvedic principles of Tamra, Agni, and Dhatu metabolism. Methods: A structured narrative review was conducted using classical Ayurvedic texts and modern databases including PubMed, Scopus, and Google Scholar. Results: Tyrosine functions as a precursor for catecholamines, thyroid hormones, and melanin. Copper serves as a catalytic cofactor in key enzymes including tyrosinase and dopamine β-hydroxylase. Emerging evidence on Cu(II)–aromatic interactions demonstrate stabilization of redox-active complexes, providing a mechanistic parallel to Ayurvedic metallurgical processing. Conclusion: A mechanistic convergence exists between Ayurvedic concepts and modern biochemical pathways involving tyrosine and copper, supporting integrative approaches in neuroendocrine and metabolic regulation.**

***Index Terms*— Ayurveda, Copper, Integrative medicine, Neurotransmitters, Trace minerals, Tyrosine**

I. INTRODUCTION

Ayurveda conceptualizes health as a dynamic equilibrium of Dosha, Dhatu, and Mala, governed by Agni and sustained by Ojas [1], [2]. These foundational principles collectively regulate physiological stability, metabolic transformation, and adaptive resilience. Among the various determinants of this balance, trace elements and dietary constituents are considered essential for maintaining internal harmony. Metals such as Tamra (copper) are described in classical texts as therapeutically valuable when subjected to purification (Shodhana) and incineration (Marana), which are believed to enhance their bioavailability and reduce toxicity [3], [4].

Modern biochemistry similarly recognizes the critical role of amino acids and trace minerals in maintaining cellular function, enzymatic activity, and neuroendocrine regulation [5], [6]. Tyrosine, a non-essential amino acid synthesized from phenylalanine or obtained from dietary sources, serves as a precursor for multiple biologically significant molecules, including catecholamines, thyroid hormones, and melanin [5], [7]. These pathways are essential for neural communication, metabolic regulation, and pigmentation. Copper, an essential trace element, functions as a catalytic cofactor in numerous enzymes involved in redox reactions and energy metabolism [6]. Its ability to alternate between Cu^{+} and Cu^{2+} oxidation states enables participation in electron transfer processes that are fundamental to cellular physiology. However, copper must be tightly regulated, as both deficiency and excess can lead to pathological consequences.

Despite significant advancements in understanding tyrosine metabolism and copper biochemistry, the integration of these mechanisms with Ayurvedic concepts remains insufficiently explored. This review aims to bridge this gap by providing a comprehensive and mechanistic synthesis of tyrosine–copper interactions in the context of Ayurvedic principles, thereby offering an integrative framework for understanding metabolic and neuroendocrine regulation.

II. METHODS

A structured narrative review methodology was employed to synthesize classical Ayurvedic knowledge with contemporary biochemical and

molecular evidence. Literature was systematically retrieved from PubMed, Scopus, and Google Scholar databases, focusing on publications up to the year 2025.

Search terms included "tyrosine metabolism," "catecholamine biosynthesis," "copper homeostasis," "dopamine beta-hydroxylase," "tyrosinase," "redox biology," "cation–π interaction," and "Tamra Bhasma." Boolean operators were used to refine search results and ensure relevance.

Inclusion criteria comprised peer-reviewed research articles, review papers, and authoritative biochemical studies that addressed mechanistic, molecular, or clinical aspects of tyrosine and copper metabolism. Classical Ayurvedic texts, including Charaka Samhita, Ashtanga Hridaya, and Rasashastra treatises, were included for conceptual analysis. Exclusion criteria included studies lacking mechanistic clarity, non-peer-reviewed sources, and publications not directly relevant to the integrative framework. Data were analyzed qualitatively to identify conceptual overlaps, mechanistic parallels, and potential translational implications.

III. TYROSINE METABOLISM IN MODERN BIOCHEMISTRY

Tyrosine metabolism represents a highly regulated biochemical network with significant implications for neurological, endocrine, and dermatological functions. The rate-limiting step in catecholamine synthesis is catalyzed by tyrosine hydroxylase, which converts tyrosine into L-DOPA in the presence of tetrahydrobiopterin (BH4), molecular oxygen, and iron [5]. This enzymatic step is subject to complex regulatory mechanisms, including feedback inhibition by catecholamines and phosphorylation-mediated activation under stress conditions.

Following the formation of L-DOPA, subsequent enzymatic reactions lead to the production of dopamine, norepinephrine, and epinephrine. Dopamine β-hydroxylase, a copper-dependent enzyme, catalyzes the conversion of dopamine to norepinephrine, thereby directly linking copper availability to neurotransmitter synthesis [6], [8]. These catecholamines play essential roles in cognition, mood regulation, and the physiological stress response. Transport of tyrosine into the central nervous system occurs via the large neutral amino acid transporter (LAT1), which is shared with other amino acids such as tryptophan and phenylalanine [5]. This competitive transport mechanism highlights the importance of dietary composition and amino acid balance in influencing neurotransmitter synthesis. Tyrosine also plays a crucial role in endocrine physiology. In the thyroid gland, iodination of tyrosine residues within thyroglobulin leads to the formation of thyroxine (T4) and triiodothyronine (T3), which regulate metabolic rate, thermogenesis, and overall energy homeostasis [5]. The activity of deiodinase enzymes further modulates the conversion of T4 to the biologically active T3. In melanocytes, tyrosinase catalyzes the oxidation of tyrosine to DOPAquinone, initiating the process of melanin synthesis [7]. This pathway is highly dependent on copper as a cofactor, emphasizing the interconnected nature of amino acid and trace mineral metabolism.

IV. COPPER'S ROLE IN TYROSINE-DEPENDENT PATHWAYS

Copper is an essential micronutrient that plays a pivotal role in biological systems due to its redox properties. It serves as a cofactor for several enzymes, including tyrosinase, dopamine β-hydroxylase, and cytochrome c oxidase, which are involved in pigmentation, neurotransmitter synthesis, and mitochondrial respiration, respectively [6], [7]. Copper homeostasis is tightly regulated through a network of proteins, including ceruloplasmin, which transports copper in the bloodstream, and ATP7A and ATP7B, which regulate intracellular copper distribution and excretion [9]. These regulatory mechanisms ensure that copper is available for enzymatic processes while preventing toxic accumulation. Disruption of copper homeostasis leads to significant clinical consequences. Copper deficiency, as observed in Menkes disease, results in impaired neurological development, connective tissue abnormalities, and hypopigmentation [8]. Conversely, copper accumulation, as seen in Wilson's disease, leads to hepatic and neurological toxicity due to excessive oxidative stress [10], [11]. At the molecular level, copper participates in electron transfer reactions and contributes to the generation and regulation of reactive oxygen species (ROS). While copper-dependent enzymes play a role in antioxidant defense, excess free copper can catalyze Fenton-like reactions,

leading to oxidative damage of cellular components [10]– [12]. Thus, copper exhibits a dual role as both an essential cofactor and a potential pro-oxidant, necessitating precise regulation to maintain physiological balance.

V. MOLECULAR INTERACTION BETWEEN TYROSINE AND COPPER

The interaction between tyrosine and copper at the molecular level is governed by coordination chemistry and cation–π interactions. Aromatic amino acids such as tyrosine possess electron-rich π systems that can interact with positively charged metal ions such as Cu (II), resulting in stabilization of metal–protein complexes [13]. These interactions are critical for the structural integrity and catalytic efficiency of copper-dependent enzymes. In enzyme active sites, copper is often coordinated by histidine residues and aromatic side chains, creating a stable environment for electron transfer and substrate oxidation [10], [13]. Cation–π interactions contribute to redox stabilization by maintaining copper in its appropriate oxidation state during catalytic cycles. This enhances enzymatic efficiency while minimizing the risk of uncontrolled oxidative reactions. Studies involving peptide–metal complexes have demonstrated that such interactions influence protein folding, stability, and catalytic specificity. This molecular understanding provides a potential bridge to Ayurvedic metallurgical practices, suggesting that traditional processes such as Shodhana and Marana may have empirically optimized the structural and functional properties of metal-based therapeutics.

VI. AYURVEDIC PERSPECTIVE ON COPPER AND AMINO ACIDS

In Ayurvedic literature, Tamra is classified as a Dhatu with therapeutic applications following appropriate processing [3], [4]. The processes of Shodhana and Marana are designed to detoxify metals, enhance their bioavailability, and render them suitable for medicinal use. Tamra is described to support Agni, the fundamental principle governing digestion and metabolic transformation. It is also associated with the regulation of Meda Dhatu and the nourishment of Rasa and Rakta Dhatus, which are essential for systemic physiological balance. Ayurvedic dietary practices emphasize Sattvic foods, including milk, legumes, seeds, and vegetables, which provide essential amino acids such as tyrosine and trace minerals like copper [1], [2]. These dietary components contribute to Dhatu poshana and the maintenance of Ojas, which represents vitality, immunity, and resilience. The traditional use of copper vessels for water storage reflects an empirical understanding of trace mineral supplementation and its role in maintaining metabolic balance and preventing disease.

VII. DISCUSSION

A. Neurobiology and Stress Axis

Tyrosine-derived catecholamines, including dopamine and norepinephrine, play a central role in regulating the physiological stress response through the hypothalamic–pituitary–adrenal (HPA) axis [5], [10]. Under conditions of acute or chronic stress, increased catecholamine synthesis places a higher demand on tyrosine availability, making it a conditionally essential substrate for maintaining cognitive performance and emotional stability. Copper-dependent dopamine β-hydroxylase catalyzes the conversion of dopamine to norepinephrine, thereby linking trace mineral status with neurochemical output [10], [14]. Deficiency in copper can impair norepinephrine synthesis, leading to altered stress responses, reduced alertness, and neurocognitive dysfunction. Conversely, excessive copper may contribute to oxidative stress within neuronal tissues, affecting neurotransmission [10], [12].

From an Ayurvedic perspective, these neurochemical processes correlate with Satva bala, representing mental strength, clarity, and emotional resilience. The maintenance of adequate Satva is closely linked to dietary quality and metabolic balance, which parallels the biochemical requirement for sufficient tyrosine and copper availability. Furthermore, Ojas, considered the essence of all Dhatus, may be interpreted as a systemic manifestation of neuroendocrine stability and adaptive capacity. Thus, the interplay between tyrosine metabolism and copper-dependent enzymatic activity provides a mechanistic substrate for understanding Ayurvedic concepts of mental resilience and stress adaptation.

B. Endocrine Regulation and Metabolic Integration

Tyrosine plays a critical role in endocrine physiology through its involvement in thyroid hormone synthesis. The iodination and coupling of tyrosine residues within thyroglobulin lead to the formation of T3 and T4, which regulate basal metabolic rate, thermogenesis, and energy utilization [5]. The activity of deiodinase enzymes further modulates the conversion of T4 to the biologically active T3, ensuring precise hormonal regulation. Copper contributes indirectly to endocrine function through its role in mitochondrial respiration and ATP production, mediated by cytochrome c oxidase [6], [9]. Efficient energy production is essential for maintaining hormonal balance and metabolic homeostasis. Disruption of copper-dependent processes may therefore influence endocrine function at both cellular and systemic levels.

In Ayurveda, these processes align closely with the concept of Agni, which governs digestion, metabolism, and transformation. Agni is not limited to gastrointestinal processes but extends to cellular and tissue-level metabolism. Impairment of Agni is associated with metabolic disorders, which may be interpreted in modern terms as dysfunctions in enzymatic activity and energy metabolism. This parallel suggests that copper-dependent enzymatic activity and tyrosine-mediated hormonal synthesis may represent biochemical correlates of Agni, providing a bridge between traditional and modern frameworks of metabolic regulation.

C. Dermatological Mechanisms and Pigmentation Biology

Melanin synthesis is a well-characterized biochemical process involving the oxidation of tyrosine by tyrosinase, a copper-dependent enzyme [7], [15]. The production of melanin not only determines pigmentation but also plays a protective role against ultraviolet radiation and oxidative stress. Recent evidence suggests that copper availability directly influences melanogenesis, with increased copper levels enhancing tyrosinase activity and melanin production [15]. This highlights the importance of trace mineral balance in maintaining normal pigmentation and skin health.

In Ayurvedic literature, Tamra is indicated in the management of dermatological conditions such as Kushtha, which encompasses a spectrum of skin disorders. The therapeutic use of Tamra may reflect its role in modulating enzymatic activity and oxidative balance within the skin. Additionally, Ayurvedic formulations often combine metals with herbal components, potentially enhancing bioavailability and therapeutic efficacy. From an integrative perspective, the role of copper in melanogenesis provides a clear biochemical basis for the traditional use of Tamra in dermatological therapy. This convergence underscores the relevance of trace minerals in maintaining skin homeostasis and highlights the potential for integrative therapeutic approaches.

D. Redox Biology, Metallomics, and Toxicological Balance

Copper's redox activity is central to its biological function. By cycling between Cu^{+} and Cu^{2+} states, copper participates in electron transfer reactions essential for enzymatic catalysis and cellular respiration [6], [10]. However, this same property also confers the potential for oxidative damage when copper homeostasis is disrupted. Excess free copper can catalyze the generation of reactive oxygen species (ROS) through Fenton-like reactions, leading to lipid peroxidation, protein oxidation, and DNA damage [10]–[12]. Such oxidative stress is implicated in the pathogenesis of neurodegenerative disorders, including Alzheimer's disease and Parkinson's disease. The emerging field of metallomics emphasizes the importance of metal ion regulation in biological systems, highlighting the need for precise control of metal bioavailability and reactivity. In this context, copper-binding proteins and transporters play a crucial role in maintaining redox balance and preventing toxicity.

Ayurvedic metallurgical processes, particularly Shodhana and Marana, may be interpreted as traditional strategies for modulating metal reactivity and enhancing safety. These processes involve repeated purification and calcination steps, which may alter the physicochemical properties of metals, reduce toxicity while preserving therapeutic potential. This conceptual alignment between Ayurvedic practices and modern metallomics suggests that traditional knowledge systems may have empirically addressed challenges related to metal toxicity and bioavailability long before their mechanistic understanding was established.

E. Translational and Clinical Implications

The integration of tyrosine metabolism and copper biochemistry has significant implications for clinical practice. Dysregulation of these pathways is associated with a range of conditions, including stress-related disorders, neurodegenerative diseases, thyroid dysfunction, and pigmentation abnormalities [10], [11]. Nutritional interventions aimed at optimizing tyrosine intake and maintaining copper balance may support neurotransmitter synthesis, hormonal regulation, and metabolic function. However, supplementation must be approached with caution, as both deficiency and excess of copper can have adverse effects.

Ayurvedic dietary practices, which emphasize natural, nutrient-rich foods and balanced intake, may provide a safe and sustainable approach to maintaining these metabolic pathways. Additionally, Ayurvedic formulations containing processed metals such as Tamra Bhasma may offer therapeutic benefits, although rigorous scientific validation is required. Future research should focus on clinical trials evaluating integrative interventions that combine dietary, biochemical, and Ayurvedic approaches. Such studies may provide valuable insights into the potential of integrative medicine in managing complex metabolic and neuroendocrine disorders.

F. Integrative Synthesis and Conceptual Framework

The convergence of tyrosine metabolism, copper-dependent enzymatic activity, and Ayurvedic principles represent a multidimensional framework for understanding physiological regulation. At the molecular level, cation–π interactions and redox cycling provide mechanistic insights into enzyme function and metabolic efficiency. At the systemic level, these processes translate into neuroendocrine regulation, energy metabolism, and tissue homeostasis. In Ayurveda, these functions are conceptualized through Agni, Dhatu equilibrium, and Ojas, which collectively maintain health and resilience. This integrative perspective highlights the potential for bridging traditional knowledge systems with modern scientific understanding. By identifying mechanistic parallels and conceptual overlaps, it becomes possible to develop holistic approaches to health that are both scientifically grounded and traditionally informed. The integrative relationship between tyrosine metabolism, copper-dependent enzymatic activity, and Ayurvedic principles is summarized in Fig. 1.



Fig. 1. Integrative framework linking tyrosine metabolism, copper homeostasis, and Ayurvedic concepts in health and disease

VIII. CONCLUSION

The convergence of Ayurvedic principles with modern biochemical insights into tyrosine and copper metabolism provides a coherent integrative framework for understanding physiological regulation. This synthesis highlights the importance of amino acids and trace minerals in neuroendocrine, metabolic, and dermatological processes. Future research should focus on mechanistic validation, standardization of Ayurvedic formulations, and clinical trials to evaluate integrative therapeutic strategies.

IX. LIMITATIONS

This review is limited by the interpretive nature of mapping Ayurvedic concepts onto modern biochemical mechanisms. The absence of direct clinical studies integrating both paradigms limits translational applicability. Additionally, variability in nutrient supplementation outcomes and the risk of copper toxicity necessitate cautious interpretation.

REFERENCES

[1] P. V. Sharma, *Charaka Samhita*. Varanasi, India: Chaukhambha Orientalia, 2014.

[2] P. Trivedi, *Ashtanga Hridaya of Vagbhata*. Varanasi, India: Chaukhambha Sanskrit Sansthan, 2018.

[3] C. Y. Jagtap, R. Galib, and P. K. Prajapati, "A brief review of Tamra Bhasma," *J. Res. Educ. Indian Med.*, vol. 19, nos. 1–2, pp. 17–28, 2013.

[4] M. Upadhyaya, *Ayurveda Prakasha*. Varanasi, India: Chaukhambha Bharati Academy, 2007.

[5] J. D. Fernstrom and M. H. Fernstrom, "Tyrosine, phenylalanine, and catecholamine synthesis and function," *J. Nutr.*, vol. 137, no. 6, suppl. 1, pp. 1539S–1547S, 2007.

[6] M. C. Linder and M. Hazegh-Azam, "Copper biochemistry and molecular biology," *Am. J. Clin. Nutr.*, vol. 63, no. 5, pp. 797S–811S, 1996.

[7] Pezzella, A. Napolitano, and G. Prota, "Tyrosinase and the oxidation of tyrosine," *Biochim. Biophys. Acta*, vol. 1336, no. 1, pp. 1–13, 1997.

[8] J. R. Prohaska, "Copper deficiency and neurological function," *Annu. Rev. Nutr.*, vol. 7, pp. 291–310, 1987.

[9] S. Lutsenko, "Mammalian copper homeostasis: Physiological roles and emerging mechanisms," *Physiol. Rev.*, vol. 105, no. 1, pp. 1–45, 2025, doi: 10.1152/physrev.00011.2024.

[10] J. Gale, A. Czlonkowska, and T. Litwin, "The physiological and pathophysiological roles of copper in the nervous system," *Biol. Rev. Camb. Philos. Soc.*, 2024, doi: 10.1111/brv.13145.

[11] J. Lin *et al*., "Copper metabolism-related mechanisms in neurodegenerative diseases," *Ann. Med.*, vol. 56, no. 1, pp. 1–15, 2024.

[12] E. Charkiewicz *et al*., "Is copper still safe for us? A comprehensive review of its biological roles and toxicity," *Cells*, vol. 13, no. 8, Art. no. 498, 2024, doi: 10.3390/cells13080498.

[13] M. Z. Wiloch, K. Kacprzak, J. M. Witkowski, *et al*., "Redox activity of copper(II) complexes with NSFRY pentapeptide," *PLoS One*, vol. 11, no. 8, Art. no. e0160256, 2016.

[14] R. Lane *et al*., "A primer on copper biology in the brain," *Trends Neurosci.*, vol. 48, no. 2, pp. 85–101, 2025.

[15] H. Uehara *et al*., "Copper supplementation enhances pigmentation and induces dopamine production," *PLoS One*, vol. 20, no. 7, Art. no. e0327352, 2025, doi: 10.1371/journal.pone.0327352.