Artificial Intelligence for Water Resources and Flood Management: Applications, Challenges, And Future Directions

V. Ganesh Ajay Velu
Assistant Professor, Civil Engineering
Narayana Engineering College, Nellore, Andhra Pradesh, India

***Abstract*—Water resources management and flood risk reduction have become major challenges due to climate variability, rapid urbanisation, population growth, changing land use, and increasing water demand. Traditional approaches to hydrological prediction and water management often face limitations associated with insufficient observations, complex environmental processes, and the need for timely decisions. Artificial intelligence (AI), particularly machine learning and deep learning, offers opportunities to improve water-related forecasting, monitoring, planning, and disaster preparedness. This paper presents a review of AI applications in water resources and flood management, focusing on rainfall–runoff modelling, river discharge prediction, flood susceptibility assessment, inundation mapping, reservoir operation, water-quality monitoring, and early-warning systems. Machine-learning algorithms, including random forests, support vector machines, artificial neural networks, and long short-term memory networks, can analyse historical and real-time hydrological data to identify patterns and support predictive decision-making. Integration with geographic information systems, remote sensing, Internet of Things sensors, and numerical hydrological models can further improve situational awareness and operational planning. However, the reliability of AI-based systems depends on data quality, model validation, uncertainty assessment, interpretability, and performance under extreme hydrological conditions. This paper discusses the benefits and limitations of these technologies and proposes a general framework for their implementation. It concludes that combining AI with established hydrological science, engineering judgement, and effective institutional warning systems can contribute to more sustainable water management and improved flood resilience**

***Index Terms*—Artificial Intelligence, Water Resources Management, Flood Forecasting, Machine Learning, Deep Learning, Rainfall–Runoff Modelling, Remote Sensing, Geographic Information System, Early-Warning Systems, Climate Resilience.**

I. INTRODUCTION

Water resources are essential for domestic consumption, agriculture, industry, energy production, and ecosystem sustainability. Effective management of water resources is necessary to support economic development and protect communities from water-related hazards. However, increasing demand, irregular rainfall, changing climatic conditions, catchment degradation, and rapid urbanisation create significant challenges for water planners and civil engineers.

Floods are among the most damaging natural hazards because they can cause loss of life, destruction of infrastructure, agricultural losses, environmental degradation, and disruption of economic activities. In urban areas, the expansion of impervious surfaces and inadequate drainage systems can increase surface runoff and waterlogging. In river basins, intense rainfall, saturated soils, and reservoir releases may contribute to elevated river levels and flooding.

Conventional hydrological methods use rainfall observations, streamflow measurements, hydraulic modelling, statistical analysis, and engineering judgement to support water management. These methods remain important, but their effectiveness may be constrained by limited monitoring networks, nonlinear hydrological processes, and the computational requirements of certain forecasting applications.

Artificial intelligence provides additional tools for analysing complex datasets and identifying relationships between meteorological conditions,

catchment characteristics, and hydrological responses. Machine-learning algorithms can support prediction, classification, anomaly detection, and optimisation. Deep-learning techniques can analyse time-dependent data and spatial imagery.

For example, long short-term memory (LSTM) networks have been investigated for rainfall–runoff modelling, while machine-learning methods have been extensively examined for flood prediction (Kratzert et al., 2018; Mosavi et al., 2018).

The integration of AI with hydrological models, remote sensing, geographic information systems (GIS), and real-time monitoring can support more informed decisions. Nevertheless, AI should be considered a decision-support technology rather than a complete replacement for established engineering methods.

II. OBJECTIVES OF THE STUDY

The primary objectives of this review are:

1. To examine the role of artificial intelligence in water resources planning and management.
2. To review AI techniques used in rainfall–runoff modelling and flood forecasting.
3. To describe applications of AI in flood mapping, reservoir operation, water-quality assessment, and early-warning systems.
4. To explain the integration of AI with GIS, remote sensing, and IoT-based monitoring.
5. To identify the technical, operational, and institutional challenges associated with AI implementation.
6. To propose future research directions for developing reliable and sustainable AI-based water management systems.

III. REVIEW METHODOLOGY

This paper adopts a narrative literature-review approach. It synthesizes selected published research on machine learning, deep learning, hydrological forecasting, flood prediction, and AI-supported water management.

The discussion draws on established research publications and technical literature addressing rainfall–runoff modelling, flood prediction, and the integration of data-driven methods with hydrological science. Particular attention is given to the suitability of different AI techniques, the types of data required, the potential operational benefits, and the limitations of model-based predictions.

The review is organised thematically into water resources management, flood forecasting, flood mapping, early-warning systems, implementation requirements, and future research needs.

This is a narrative review rather than a systematic review or an experimental investigation. No new field measurements, model training, numerical simulations, or original performance tests are claimed. A formal systematic review would require a documented database-search strategy, defined inclusion and exclusion criteria, a reproducible screening process, and a structured assessment of the selected studies.

IV. ARTIFICIAL INTELLIGENCE TECHNIQUES FOR WATER RESOURCES MANAGEMENT

4.1. Machine Learning

Machine learning enables computer systems to learn patterns from historical data and use those patterns to estimate outcomes for new observations. Common machine-learning methods in water resources applications include:

Random Forest:

A tree-based ensemble algorithm suitable for regression and classification. It can support flood-susceptibility mapping, water-quality estimation, and the identification of important environmental variables.

Support Vector Machine:

A supervised learning method that can be used for classification and regression. It may be applied to water-level prediction, rainfall classification, and flood-risk assessment.

Gradient Boosting:

An ensemble-learning approach that combines multiple decision trees to improve predictive performance. It can be useful for streamflow estimation and hydrological risk classification.

Artificial Neural Networks:

These models learn nonlinear relationships between input and output variables. They can be applied to rainfall forecasting, reservoir inflow estimation, and water-demand prediction.

The performance of these methods depends on the availability of representative data, appropriate input-variable selection, model calibration, and independent validation.

4.2. Deep Learning

Deep learning uses multilayer neural networks to identify complex patterns in large datasets.

Long short-term memory networks are particularly relevant to time-series applications because they can learn temporal dependencies in sequential observations. In hydrology, these models may be used to analyse rainfall, river discharge, soil moisture, and historical water levels.

Convolutional neural networks can process satellite imagery and gridded environmental information. They may support the identification of flooded areas, classification of land cover, and extraction of spatial features.

Deep-learning models can provide useful predictions, but their complexity may increase computational requirements and reduce interpretability. Their performance must be evaluated using independent observations and appropriate comparison models.

4.3. Hybrid AI and Hydrological Models

Hydrological and hydraulic models represent physical processes such as infiltration, runoff generation, river routing, and floodplain flow. AI models, in contrast, learn relationships from data.

A hybrid approach combines these complementary capabilities. AI may be used to estimate model parameters, correct systematic prediction errors, or provide rapid approximations of computationally demanding simulations.

For example, a rainfall–runoff model can estimate river discharge, while an AI model can learn residual errors between simulated and observed discharge. The corrected prediction can then be evaluated against independent flood events.

Hybrid approaches can improve efficiency and flexibility, but they require careful calibration and validation to ensure that predictions remain physically plausible.

V. APPLICATIONS OF AI IN WATER RESOURCES MANAGEMENT

5.1. Water Demand Forecasting

Accurate estimation of water demand is essential for municipal supply systems, irrigation planning, and industrial operations.

AI models can analyse historical consumption, population trends, temperature, rainfall, seasonal variation, and agricultural activity to estimate future demand.

These forecasts can assist water authorities in planning distribution, identifying periods of high consumption, and allocating available resources.

For example, a municipal water utility may use historical daily consumption and weather observations to estimate demand during the summer season. The resulting estimates can support pumping schedules and reservoir planning.

However, reliable forecasting also requires consideration of population changes, leakage, infrastructure constraints, and changes in consumer behaviour.

5.2. Reservoir Operation and Water Allocation

Reservoirs support drinking-water supply, irrigation, hydropower generation, environmental flows, and flood control.

AI-based forecasting can estimate reservoir inflows using rainfall, catchment conditions, upstream discharge, and historical inflow records. These predictions can help engineers compare alternative storage and release scenarios.

Optimisation algorithms may also support the allocation of water among competing users.

Nevertheless, AI recommendations must comply with approved operating rules, dam-safety requirements, environmental regulations, and downstream flood-control obligations. Critical reservoir operations should retain appropriate engineering oversight.

5.3. Groundwater Assessment

Groundwater is an important resource for drinking water and agriculture, particularly in regions with seasonal rainfall and limited surface-water availability.

Machine-learning models can analyse groundwater-level records, rainfall, geology, soil characteristics,

land use, and pumping information to estimate groundwater-level changes.

GIS-integrated AI models may also help identify potential groundwater recharge zones by combining information on terrain, drainage, geology, and land cover.

These predictions can support groundwater monitoring and planning. However, groundwater decisions should be based on hydrogeological evidence, field investigations, and appropriate water-balance assessments.

5.4. Water-Quality Monitoring

Water quality can be affected by industrial discharge, agricultural runoff, sewage, sediment transport, and changes in temperature.

AI models can analyse sensor measurements and laboratory observations to estimate selected water-quality indicators or detect unusual patterns.

Potential applications include:

- Estimation of turbidity and dissolved oxygen.
- Identification of unusual sensor readings.
- Monitoring of changing water-quality conditions.
- Prioritisation of sampling locations.
- Analysis of relationships between rainfall and pollution events.

AI predictions should not replace required laboratory testing or regulatory monitoring. Unusual results should be verified using suitable field and laboratory procedures.

5.5. Watershed Management

Watershed management requires an understanding of rainfall, runoff, erosion, land cover, soil properties, vegetation, and drainage characteristics.

AI can assist in identifying areas that may be vulnerable to soil erosion, sediment movement, and excessive surface runoff.

When combined with remote sensing and GIS, these techniques can help prioritise watershed interventions, including vegetation restoration, drainage improvement, soil conservation, and rainwater harvesting.

The final selection of interventions should consider field conditions, local communities, environmental effects, and engineering feasibility.

VI. APPLICATIONS OF AI IN FLOOD MANAGEMENT

6.1. Rainfall Forecasting

Rainfall is one of the most important variables in flood prediction.

AI models can analyse historical rainfall, atmospheric observations, weather-model outputs, and radar or satellite products to estimate rainfall patterns.

Short-term rainfall prediction, often called nowcasting, can support decisions during rapidly developing storms. Longer-term forecasts may assist preparedness and reservoir planning.

The usefulness of AI rainfall forecasts depends on the spatial and temporal resolution of the observations, the quality of the input forecasts, and the ability of the model to represent unusual weather conditions.

AI predictions should therefore be evaluated against observed rainfall and established meteorological forecasting products.

6.2. Rainfall–Runoff Modelling

Rainfall–runoff modelling estimates how precipitation is transformed into streamflow within a catchment.

The response depends on rainfall intensity, soil moisture, infiltration, vegetation, terrain, drainage networks, and land-use conditions.

Machine-learning models can learn relationships between these variables and observed discharge.

LSTM networks are one example of a deep-learning approach investigated for rainfall–runoff modelling. Kratzert et al. (2018) demonstrated the potential of LSTM networks for hydrological modelling across multiple catchments.

In a practical application, rainfall observations and previous discharge measurements can be used to predict river flow at a downstream monitoring station. Such predictions can support flood preparedness, but the model must be tested during both ordinary conditions and significant flood events.

6.3. River Water-Level and Discharge Prediction

Flood management requires timely estimates of river stage and discharge.

AI models can use upstream water levels, rainfall, previous discharge measurements, catchment characteristics, and reservoir information to estimate future river conditions.

For example, if upstream water levels increase rapidly following heavy rainfall, an AI-supported system may estimate the likely downstream response and identify whether predefined warning thresholds could be exceeded. The forecast should communicate its lead time and uncertainty. Predictions must also account for tributary inflows, reservoir releases, embankment conditions, and other factors that may influence river behaviour.

6.4. Flood Susceptibility Mapping

Flood susceptibility mapping identifies locations that may be more prone to flooding based on environmental and geographical characteristics.

Potential input variables include:

- Elevation and slope.
- Distance from rivers and drainage channels.
- Rainfall intensity.
- Soil type and infiltration characteristics.
- Land use and land cover.
- Drainage density.
- Historical flood occurrence.

Machine-learning algorithms such as random forests and support vector machines can analyse these variables to classify relative flood susceptibility.

GIS can then display the resulting classifications as spatial maps.

Flood susceptibility maps are useful for preliminary planning, infrastructure assessment, and prioritisation of detailed investigations. They should not be interpreted as precise predictions of flood depth or timing unless supported by suitable hydraulic analysis and validation.

6.5. Flood Inundation Mapping

Flood inundation mapping estimates the areas covered by floodwater.

Satellite imagery can provide observations of water extent during or after a flood. Image-processing and deep-learning methods can help distinguish water-covered surfaces from other land-cover types.

These observations can support damage assessment, emergency planning, and evaluation of flood models.

Optical satellite imagery may be affected by clouds, while radar imagery has different limitations related to surface characteristics, vegetation, and urban structures.

Combining remote sensing with elevation data and hydraulic modelling can provide a more comprehensive understanding of inundation patterns.

6.6. Urban Flood Management

Urban flooding is influenced by intense rainfall, impervious surfaces, inadequate drainage, construction activities, and obstruction of stormwater channels.

AI systems can help identify waterlogging hotspots, forecast short-duration rainfall, monitor drainage conditions, and prioritise maintenance.

For example, a city may combine rainfall sensors, drainage-level measurements, road-elevation data, and historical complaints to identify locations where waterlogging repeatedly occurs.

These results can support the planning of improved drainage, additional storage, permeable surfaces, and maintenance programmes.

However, reliable urban flood modelling requires sufficiently detailed terrain, drainage-network, and rainfall data.

6.7. AI-Based Flood Early-Warning Systems

Early-warning systems aim to provide timely information that enables communities and authorities to take protective action.

An AI-supported flood-warning system may combine:

1. Weather forecasts and rainfall observations.
2. River-level and discharge measurements.
3. Reservoir-level information.
4. Catchment and terrain data.
5. Predicted flood thresholds.
6. Population and infrastructure exposure.
7. Communication and emergency-response procedures.

When the model indicates that a warning threshold may be exceeded, the system can support the preparation of alerts for the responsible authorities.

Warnings may be communicated through mobile messages, public-address systems, sirens, radio, and local administrative networks.

An effective warning system must consider accessibility, local languages, communication failures, false alarms, and the needs of vulnerable communities. Prediction alone does not guarantee effective disaster response.

VII. INTEGRATION OF AI WITH GIS, REMOTE SENSING, AND IoT

The integration of AI with modern monitoring technologies can improve the availability and interpretation of hydrological information.

7.1. Geographic Information Systems

GIS provides tools for storing, analysing, and displaying spatial information.

When integrated with AI, GIS can support flood susceptibility mapping, watershed prioritisation, infrastructure exposure assessment, and the identification of communities located in flood-prone areas.

Spatial outputs can help planners identify locations requiring additional drainage capacity, protective infrastructure, or emergency access routes.

7.2. Remote Sensing

Remote sensing provides information on rainfall, land cover, vegetation, surface-water extent, soil moisture, and terrain.

Satellite observations are particularly valuable in large or remote catchments where ground monitoring networks are limited. AI can analyse these datasets to extract environmental features and support flood mapping and catchment assessment.

Satellite-derived information should be checked against available ground observations because different sensors have different spatial resolutions, revisit periods, and measurement limitations.

7.3. Internet of Things Sensors

IoT-based systems use connected sensors to collect and transmit measurements.

In water management, sensors can monitor rainfall, river stage, reservoir level, soil moisture, and selected water-quality indicators.

The measurements can be transmitted to a central platform where AI models process them and generate updated estimates.

However, practical systems require sensor calibration, power supply, reliable communications, data-quality checks, cybersecurity, and regular maintenance.

7.4. Integrated Decision-Support Architecture

An integrated AI-enabled water management system can be organised into five layers:

Layer 1: Data collection. Gather meteorological, hydrological, geographical, remote-sensing, and infrastructure data.

Layer 2: Data processing. Check data quality, manage missing observations, align timestamps, and prepare model inputs.

Layer 3: AI modelling. Apply suitable machine-learning or deep-learning models to estimate rainfall, discharge, water demand, or flood susceptibility.

Layer 4: Decision support. Combine predictions with engineering rules, hydraulic models, warning thresholds, and exposure information.

Layer 5: Communication and monitoring. Deliver forecasts and warnings to responsible agencies and communities, monitor performance, and update the system when necessary.

This architecture supports a coordinated approach to water planning and flood preparedness.

VIII. PROPOSED IMPLEMENTATION FRAMEWORK

A practical implementation programme should follow a structured process.

Step 1: Define the Problem

The first step is to specify the intended application, such as river-level forecasting, reservoir inflow prediction, urban waterlogging detection, or flood susceptibility mapping.

The required forecast lead time and decision-making purpose should be clearly defined.

Step 2: Collect Relevant Data

Depending on the application, the dataset may include rainfall, temperature, discharge, water levels, soil characteristics, terrain, land use, and historical flood records.

The period, resolution, and reliability of the available observations should be documented.

Step 3: Prepare the Dataset

Data preparation includes correcting obvious measurement errors, checking missing values, aligning time records, and ensuring consistent units.

Training data should represent the range of conditions under which the model is expected to operate.

Step 4: Select Appropriate Models
Potential models include random forests, gradient boosting, support vector regression, artificial neural networks, and LSTM networks.
A simple baseline model should be established before adopting a complex AI method.

Step 5: Train and Validate
Historical observations can be divided into training, validation, and testing periods.
For time-series prediction, chronological separation is important to avoid using future information during training.
Where possible, validation should also examine performance across different catchments and independent flood events.

Step 6: Evaluate Model Performance
Suitable evaluation measures depend on the task.
For continuous predictions, common measures include mean absolute error (MAE), root mean square error (RMSE), and coefficient of determination (R^2).

For flood classification, precision, recall, F1-score, and precision–recall analysis may be useful.
For operational warning systems, lead time, missed events, false alarms, and the consequences of incorrect warnings should also be evaluated.
No single metric fully describes model performance.

Step 7: Deploy with Engineering Safeguards
The AI model should be integrated with approved operational procedures and clear responsibility for issuing warnings or making infrastructure decisions.
Uncertainty information and fallback procedures should be available when sensor data fail or model performance becomes unreliable.

Step 8: Monitor and Update
After deployment, performance should be monitored continuously.
Changes in climate, catchment land use, monitoring equipment, and river behaviour may affect model reliability. Periodic review and controlled updating are therefore necessary.

IX. BENEFITS OF AI IN WATER RESOURCES AND FLOOD MANAGEMENT

AI offers several potential benefits when implemented appropriately.

Improved forecasting:
AI can analyse historical and real-time data to estimate future hydrological conditions.

Faster information processing:
Automated systems can process large datasets and update predictions as new observations become available.

Improved spatial assessment:
GIS and remote-sensing integration can help identify flood-prone locations and exposed infrastructure.

Better resource planning:
Water-demand and reservoir-inflow predictions can support planning and allocation decisions.

Enhanced preparedness:
Timely forecasts can help authorities prepare emergency resources and issue warnings.
Support for sustainable development: better information can contribute to water conservation, watershed management, and climate resilience.
These benefits are potential outcomes rather than guaranteed improvements. They depend on data quality, model performance, institutional capacity, and the ability to translate predictions into effective action.

X. CHALLENGES AND LIMITATIONS

10.1. Data Availability and Quality

AI models require suitable data for training and evaluation. In many catchments, rainfall and river-level measurements may be sparse, incomplete, or inconsistent.

Sensor failures and differences in measurement practices can introduce errors.
Improving monitoring networks and establishing quality-control procedures are therefore important prerequisites.

10.2. Prediction of Extreme Events

Extreme floods may be poorly represented in historical datasets.

A model that performs well under ordinary conditions may not perform equally well during rare, severe events.

Testing should include significant historical floods where reliable observations are available, together with appropriate uncertainty analysis and conservative operational safeguards.

10.3. Model Interpretability

Complex AI models may not clearly explain why a prediction was generated.

Limited interpretability can reduce confidence among engineers and decision-makers.

Explainable AI techniques, transparent documentation, and comparison with physical models can help improve understanding.

10.4. Generalisation to Different Catchments

A model developed for one catchment may not perform well in another because of differences in climate, topography, soil, drainage, land use, and available observations.

Local calibration and independent validation are essential before transferring a model to a new location.

10.5. Infrastructure and Financial Constraints

Real-time systems require sensors, communication networks, computing facilities, software, maintenance, and trained personnel. These costs may create difficulties for smaller municipalities and resource-constrained regions.

Implementation should therefore consider long-term operating costs and local technical capacity, not only initial development costs.

10.6. Cybersecurity and Data Governance

Connected monitoring systems can be vulnerable to communication failures, unauthorised access, and manipulation of data.

Appropriate access controls, secure communication, backup systems, and clear data-governance procedures are necessary.

10.7. Human and Institutional Factors

Warnings are useful only when authorities and communities receive, understand, trust, and act upon them.

AI systems must therefore be integrated with disaster-management institutions, public communication procedures, evacuation planning, and community awareness programmes.

XI. FUTURE RESEARCH DIRECTIONS

Future research should focus on improving the reliability and practical value of AI-based water management systems.

11.1. Physics-Informed Artificial Intelligence

Combining physical hydrological principles with data-driven learning may improve model consistency and help reduce unrealistic predictions.

Research should examine how physical constraints can be incorporated without unnecessarily limiting the ability to learn complex relationships.

11.2. Explainable and Uncertainty-Aware Models

Future models should communicate not only predicted water levels or discharge but also uncertainty and the conditions under which the prediction may be unreliable.

Explainable methods can help engineers understand the influence of rainfall, soil moisture, catchment conditions, and other variables.

11.3. Climate-Resilient Flood Forecasting

Historical observations may not fully represent future climatic conditions.

Research should evaluate AI models under changing rainfall patterns, altered land use, and different climate scenarios. Models should be tested for robustness rather than assuming that past relationships will remain unchanged.

11.4. AI for Data-Scarce Catchments

Many watersheds lack long-term monitoring records.

Potential research directions include transfer learning, regional modelling, satellite-derived data, and methods for learning from related catchments.

These methods require careful local validation before being used in operational flood management.

11.5. Digital Twins for Water Infrastructure

Digital twins combine digital representations of physical systems with updated observations and simulation tools.

When integrated with AI, digital twins may support reservoir planning, drainage-system assessment, and infrastructure monitoring.

Future studies should evaluate the accuracy, cost, scalability, and practical decision-making value of such systems.

11.6. Community-Centred Early-Warning Systems

Research should examine how forecast information can be converted into clear and actionable warnings.

This includes assessing warning lead time, local-language communication, public response, accessibility, and the needs of vulnerable populations.

The effectiveness of an early-warning system should be assessed through both technical performance and its contribution to reducing risk.

XII. CONCLUSION

Artificial intelligence offers promising opportunities to improve water resources management and flood preparedness through rainfall forecasting, rainfall–runoff modelling, river-level prediction, flood susceptibility assessment, water-quality monitoring, reservoir planning, and early-warning systems.

Machine-learning and deep-learning methods can analyse complex relationships within historical and real-time environmental data. Their integration with GIS, remote sensing, IoT sensors, and hydrological models can support timely information and better-informed engineering decisions.

However, successful implementation depends on reliable observations, representative training data, independent validation, uncertainty assessment, interpretability, and appropriate operational safeguards. Particular care is required when predicting rare extreme floods and applying models to catchments that differ from their training environments.

AI should complement established hydrological and hydraulic modelling, professional engineering judgement, and community-based disaster management. It should not be treated as a substitute for dam-safety procedures, verified measurements, or responsible public-warning systems.

Future developments in physics-informed AI, explainable modelling, climate-resilient forecasting, and integrated digital monitoring may further strengthen these applications. With appropriate validation and governance, AI can contribute to more sustainable water management, improved flood preparedness, and greater resilience to water-related hazards.

REFERENCES

[1] Mosavi, P. Ozturk, and K.-W. Chau, “Flood prediction using machine learning models: Literature review,” *Water*, vol. 10, no. 11, Art. no. 1536, 2018, doi: 10.3390/w10111536.

[2] F. Kratzert, D. Klotz, C. Brenner, K. Schulz, and M. Herrnegger, “Rainfall–runoff modelling using long short-term memory (LSTM) networks,” *Hydrology and Earth System Sciences*, vol. 22, pp. 6005–6022, 2018, doi: 10.5194/hess-22-6005-2018.

[3] F. Kratzert, D. Klotz, M. Herrnegger, *et al.*, “Toward improved predictions in ungauged basins: Exploiting the power of machine learning,” *Water Resources Research*, vol. 55, no. 12, pp. 11344–11354, 2019, doi: 10.1029/2019WR026065.

[4] S. Hochreiter and J. Schmidhuber, “Long short-term memory,” *Neural Computation*, vol. 9, no. 8, pp. 1735–1780, 1997.

[5] United Nations Office for Disaster Risk Reduction, *Sendai Framework for Disaster Risk Reduction 2015–2030*. United Nations, 2015.

[6] World Meteorological Organization, *WMO Guidelines on Multi-Hazard Impact-Based Forecast and Warning Services*. World Meteorological Organization, 2021.

[7] I. Goodfellow, Y. Bengio, and A. Courville, *Deep Learning*. Cambridge, MA, USA: MIT Press, 2016.

[8] World Meteorological Organization, *WMO HydroSOS: Hydrological Status and Outlook System—Concept and Implementation*. World Meteorological Organization, 2021.