Kwatha Kalpana: A Critical Appraisal of Classical Decoction Technology, Pharmaceutical Standardisation and Contemporary Research Perspectives

Dr. Snehal Kishor Dhayafule
Assistant Professor, Department of Rasashastra and Bhaishajya Kalpana, SGR Ayurved Mahavidyalaya, Solapur, Maharashtra, India.

***Abstract*—Background: Kwatha Kalpana is an important traditional pharmaceutical preparation described in classical Ayurvedic literature. It involves the extraction of constituents from medicinal substances using water and controlled heating. Despite its extensive traditional application, variations in raw material characteristics, particle size, solvent proportion, heating conditions and storage practices may influence the quality and reproducibility of the preparation.**

Objective: To critically review the classical principles of Kwatha Kalpana, traditional preparation procedures, pharmaceutical processing variables, standardization approaches and contemporary scientific perspectives.

Methods: A narrative review was undertaken using classical Ayurvedic treatises, authoritative pharmaceutical reference books, official pharmacopoeial publications and contemporary scientific literature. Relevant sources were identified through PubMed, Google Scholar and official publications of the World Health Organization and the Ministry of AYUSH.

Discussion: Classical literature describes specific procedures for decoction preparation, including drug selection, coarse powder preparation, water proportion, controlled heating and volume reduction. Contemporary pharmaceutical science provides additional approaches for assessing physicochemical characteristics, phytochemical composition, microbial quality and stability. Chromatographic fingerprinting and quantitative marker estimation may improve batch-to-batch consistency. Modern extraction technologies offer opportunities for process optimization but require formulation-specific comparative validation.

Conclusion: Kwatha Kalpana represents an established traditional aqueous extraction technology. Integrating classical pharmaceutical principles with validated analytical procedures may improve its quality, reproducibility and scientific evaluation. Further experimental research is required to establish formulation-specific processing standards and clinical evidence.

***Index Terms*—Kwatha Kalpana, Ayurvedic decoction, Kashaya Kalpana, Pharmaceutical standardization, Aqueous extraction, Quality control.**

I. INTRODUCTION

Medicinal plants have been used in traditional healthcare systems for centuries. Different pharmaceutical procedures have been developed to transform crude medicinal substances into suitable dosage forms. Among these, decoction preparation represents one of the oldest and widely practiced methods of extracting medicinal constituents using water as the extraction medium.

In Ayurveda, pharmaceutical preparations are classified according to the nature of the ingredients, processing methods and intended dosage forms. Bhaishajya Kalpana, the pharmaceutical branch of Ayurveda, describes several methods of processing medicinal substances to obtain preparations suitable for therapeutic administration.

Kwatha Kalpana is an important liquid pharmaceutical preparation in which medicinal substances are boiled with water under prescribed conditions, followed by filtration to obtain the decoction. It is traditionally included among the Panchavidha Kashaya Kalpana, together with Swarasa, Kalka, Hima and Phanta. [1,2]

Sharangadhara Samhita provides a systematic description of decoction preparation in Madhyama Khanda, Chapter 2. The classical procedure describes the addition of sixteen parts of water to one part of crushed medicinal material, followed by boiling until one-eighth of the initial liquid remains. [2]

The preparation of Kwatha involves several interconnected pharmaceutical operations, including selection and authentication of raw materials, size reduction, addition of an appropriate quantity of water, heating, volume reduction and filtration.

From a contemporary pharmaceutical perspective, decoction preparation can be understood as thermal aqueous extraction. The extraction process facilitates the transfer of water-soluble constituents from the medicinal plant matrix into the liquid phase. However, the final chemical composition depends on the botanical identity, physical characteristics and chemical stability of the ingredients, together with the processing conditions. [3,4]

With increasing interest in standardized herbal medicines, the scientific evaluation of traditional dosage forms has become important. Reproducible manufacturing procedures, validated analytical methods, appropriate quality-control specifications and stability studies are essential for maintaining product quality.

Although classical texts provide fundamental pharmaceutical principles, their translation into modern manufacturing practices requires systematic evaluation. The integration of traditional pharmaceutical knowledge with contemporary analytical sciences may provide a rational approach to improving the quality and reproducibility of Kwatha preparations.

AIM: To critically review the classical principles, pharmaceutical preparation, standardization and contemporary research perspectives of Kwatha Kalpana.

Objective: To study the classical concept and preparation of Kwatha, understand the factors affecting its quality, review its standardization and quality control, explore modern extraction techniques, and identify future research possibilities.

II. REVIEW METHODOLOGY

The present article was prepared as a narrative review of classical and contemporary literature.

The literature sources included classical Ayurvedic treatises, recognized pharmaceutical reference books, official pharmacopoeial publications, WHO guidelines and peer-reviewed scientific articles.

Relevant publications were identified through searches of PubMed, Google Scholar and scientific journal platforms. Search terms included Kwatha Kalpana, Kashaya Kalpana, Ayurvedic decoction, herbal extraction, aqueous extraction, pharmaceutical standardization, phytochemical evaluation and quality control of herbal formulations.

Particular attention was given to Sharangadhara Samhita, Charaka Samhita, Ashtanga Hridaya, the Ayurvedic Pharmacopoeia of India and contemporary literature on medicinal plant extraction and extract characterization.

Classical sources were examined to identify traditional pharmaceutical principles, whereas modern publications were considered for their relevance to extraction technology, analytical evaluation and quality assurance.

The review is descriptive and interpretative in nature. It does not involve original experimental work, clinical intervention or quantitative meta-analysis. No formal systematic-review protocol or PRISMA-based selection process was undertaken.

III. CLASSICAL CONCEPT AND PHARMACEUTICAL FOUNDATION OF KWATHA KALPANA

Definition and Fundamental Concept

Kwatha is a liquid pharmaceutical preparation obtained by boiling medicinal substances with an appropriate liquid medium, generally water, until the prescribed degree of reduction is achieved. The preparation is subsequently filtered to obtain the liquid extract.

The process involves the following operations:

- Selection and authentication of medicinal ingredients.
- Cleaning and removal of extraneous materials.
- Coarse powder preparation.
- Addition of the prescribed quantity of water.
- Controlled heating and boiling.
- Reduction to the required volume.
- Filtration and collection.

The pharmaceutical significance of this procedure lies in the extraction of constituents capable of dissolving or dispersing in an aqueous medium.

The final composition is influenced by the solubility and stability of individual constituents, the nature of

the plant matrix and the conditions under which extraction is performed. [3,4]

Classical References in Ayurvedic Literature

Classical Ayurvedic literature describes different methods of preparing liquid medicinal formulations.

Charaka Samhita refers to Shrita, a preparation obtained through the application of heat, and describes the pharmaceutical relevance of processed medicinal substances. [1]

Sharangadhara Samhita provides a specific and systematic description of Kwatha preparation in Madhyama Khanda, Chapter 2, verses 1–2. The text describes the addition of sixteen parts of water to one part of crushed medicinal material, boiling in an earthen vessel and reducing the liquid to one-eighth of its initial volume. The resulting preparation is administered lukewarm. [2]

The classical verse is:

पानीयं षोडशगुणं क्षुण्णे द्रव्यपले क्षिपेत्।मृत्पात्रे क्काथयेद् ग्राह्यमष्टमांशावशेषितम्॥१॥
तज्जलं पाययेद्धीमान् कोष्णं मृद्वग्निसाधितम्।श्रुतः क्काथः कषायश्च निर्यूहः स निगद्यते॥२॥

This description represents an important reference for understanding the traditional extraction process.

However, it should not be assumed that this ratio is universally applicable to all decoctions. Classical literature describes variations according to the nature and quantity of the medicinal substances and the purpose of preparation. [2,5]

Panchavidha Kashaya Kalpana

Sharangadhara Samhita describes five fundamental pharmaceutical preparations, commonly known as Panchavidha Kashaya Kalpana:

1. Swarasa – expressed juice.
2. Kalka – medicinal paste.
3. Kwatha – decoction.
4. Hima – cold infusion.
5. Phanta – hot infusion.

These preparations differ in their pharmaceutical processing and extraction conditions.

Swarasa involves mechanical expression, Kalka involves grinding, Kwatha employs boiling, Hima involves cold-water extraction and Phanta involves the use of hot water without prolonged boiling.

The distinction is important because the extraction temperature and processing duration may influence the chemical composition of the final preparation. [2,5]

Table 1. Fundamental characteristics of Panchavidha Kashaya Kalpana

Preparation.	Pharmaceutical procedure	Principal processing condition
Swarasa	Expression of fresh medicinal material	Mechanical extraction
Kalka	Grinding of medicinal material	Mechanical size reduction
Kwatha	Boiling with water	Thermal aqueous extraction
Hima	Soaking in cold water	Cold extraction
Phanta	Addition of hot water	Hot infusion

Classical Variations in Water Proportion

The proportion of water used for decoction preparation is an important pharmaceutical consideration.

Sharangadhara Samhita describes the general procedure of sixteen parts of water with reduction to one-eighth of the original liquid volume. [2]

Other classical pharmaceutical descriptions provide variations according to the nature of the medicinal material and the quantity of ingredients. Kashyapa Samhita, for example, describes different water proportions and reduction endpoints in its pharmaceutical context. [5]

Therefore, different ratios should be interpreted in relation to their original textual context rather than treated as interchangeable standards.

A critical distinction must also be made between:

- The initial drug-to-water proportion.
- The final residual volume.
- The quantity of liquid actually recovered after filtration.

These are separate pharmaceutical parameters and should be documented independently during experimental preparation.

IV. CLASSICAL PHARMACEUTICAL METHOD OF KWATHA PREPARATION

Selection and Authentication of Raw Materials

The quality of the finished decoction depends fundamentally on the identity and quality of the medicinal substances.

Correct botanical identification is essential because substitution, adulteration and misidentification may alter the chemical composition of the preparation.
The raw materials should be evaluated for:

- Correct botanical identity.
- Appropriate plant part.
- Macroscopic and microscopic characteristics.
- Foreign matter.
- Moisture content.
- Insect infestation.
- Fungal contamination.
- Compliance with relevant pharmacopoeial specifications.

The Ayurvedic Pharmacopoeia of India provides official monograph-based requirements for the identification and evaluation of applicable Ayurvedic raw materials and formulations. [6]

Preparation of Coarse Powder

The authenticated and cleaned medicinal substances are subjected to appropriate size reduction.
In traditional pharmaceutical terminology, the material is generally prepared in Yavakuta form, indicating a coarse or moderately crushed condition.
Particle size has a direct influence on the surface area available for extraction. Smaller particles generally provide a greater surface area for solvent contact. However, excessively fine powder may cause filtration difficulties and increase the suspended material in the final preparation.
The official Pharmacopoeial Monograph of AYUSH Kvatha Curna provides a measurable specification for coarse powder:

- All particles should pass through a 710 µm IS sieve (sieve number 22).
- Not more than 10% should pass through a 355 µm IS sieve (sieve number 44).

This specification applies to the AYUSH Kvatha Curna monograph and should not automatically be interpreted as a universal requirement for every classical Kwatha formulation. [6]

Addition of Water

Water acts as the extraction medium in conventional Kwatha preparation.
The classical water proportion is determined according to the relevant formulation instructions and the nature of the medicinal ingredients.
The commonly described Sharangadhara procedure uses sixteen parts of water for one part of crushed medicinal material, followed by reduction to one-eighth of the initial liquid volume. [2]
Water quality is also important. The official AYUSH formulation monograph refers to the applicable pharmacopoeial requirements for Jala when water is used as an ingredient. [6]

Heating and Volume Reduction

The medicinal material and water are subjected to controlled heating in a suitable vessel.
During heating, water penetrates the plant matrix and facilitates the dissolution and diffusion of extractable constituents into the surrounding liquid.
The process is continued until the prescribed reduction is achieved.
The rate of heating, vessel geometry, evaporation surface, environmental conditions and nature of the medicinal ingredients may influence the duration required to reach the endpoint.
Therefore, modern pharmaceutical standardization should preferably document both the prescribed final volume and the relevant processing conditions.

Filtration and Collection

After achieving the prescribed reduction, the preparation is filtered through a suitable filtration medium.
Filtration removes coarse plant residues and produces a comparatively uniform liquid preparation.
The process should be sufficiently controlled to minimise variations in suspended matter and extract recovery.
The final preparation should be evaluated for appearance, colour, odour, pH, total solids and other formulation-specific parameters.

Table 2. Major stages in classical Kwatha preparation

Stage	Pharmaceutical operation	Quality consideration
1	Raw drug selection	Identity and purity
2	Cleaning	Removal of foreign matter
3	Coarse powder preparation	Particle-size distribution
4	Water addition	Drug-to-water proportion

5	Heating	Temperature and duration
6	Volume reduction	Defined extraction endpoint
7	Filtration	Extract recovery and clarity
8	Collection	Final yield and quality

V. PHARMACEUTICAL FACTORS INFLUENCING KWATHA QUALITY

The quality of Kwatha is determined by the combined influence of raw material characteristics and manufacturing conditions. Variations in these parameters may produce differences in the chemical composition and physical characteristics of the finished preparation. [3,4]

Particle Size

Particle size influences the surface area available for solvent contact. Coarse particles may require longer extraction periods, whereas excessively fine particles may create filtration difficulties.

The optimum particle size should be determined according to the characteristics of each formulation.

Drug-to-Water Ratio

The quantity of water determines the availability of the extraction medium.

Insufficient water may restrict extraction, whereas excessive water may increase processing duration and energy requirements.

The final concentration is also influenced by the initial water quantity and the extent of evaporation.

Temperature and Heating Rate

Temperature affects solvent penetration, diffusion and the solubility of different phytochemical constituents.

Higher temperatures may increase extraction rates for certain constituents. However, prolonged heating may cause degradation, oxidation, hydrolysis or other chemical changes in thermolabile substances.

Duration of Extraction

Extraction duration influences the amount and nature of constituents transferred into the aqueous medium.

A shorter duration may result in incomplete extraction, whereas prolonged heating may increase energy consumption and potentially degrade sensitive constituents.

Nature of the Vessel

The vessel used for preparation may influence heat transfer, temperature distribution and possible interactions with the preparation.

Stainless steel vessels are commonly considered for pharmaceutical manufacturing because of their durability, clean ability and suitability for controlled processing.

The suitability of the vessel material should be established through compatibility and cleaning validation.

Quality of Water

Water quality is an essential consideration in pharmaceutical manufacturing.

The extraction medium should meet the relevant pharmaceutical requirements. Variations in hardness, dissolved minerals, microbial contamination and pH may affect extraction characteristics and product quality.

Table 3. Critical pharmaceutical variables and their potential influence

Variable	Potential influence	Suggested control
Raw material identity	Chemical composition	Botanical authentication
Particle size	Extraction rate	Sieve analysis
Water proportion	Extraction efficiency	Gravimetric measurement
Temperature	Solubility and stability	Temperature monitoring
Heating duration	Extract composition	Defined endpoint
Vessel material	Heat transfer and compatibility	Material qualification
Filtration	Extract clarity and yield	Defined filtration procedure
Storage	Chemical and microbial stability	Stability assessment

VI. CONTEMPORARY PHARMACEUTICAL STANDARDISATION OF KWATHA

Standardization is essential for maintaining the quality, consistency and reproducibility of herbal pharmaceutical preparations.

Herbal formulations may contain numerous chemical constituents whose concentrations depend on botanical, geographical, seasonal and processing-related factors.
Consequently, the standardization of Kwatha should involve an integrated assessment of raw materials, manufacturing processes and finished products. [6–8]

Organoleptic Evaluation
Organoleptic evaluation is the preliminary assessment of a formulation through its sensory characteristics.
The principal parameters include:

- Colour.
- Odour.
- Appearance.
- Homogeneity.
- Presence of visible foreign matter.

Taste evaluation should be performed only when appropriate and safe.
These parameters provide preliminary information about batch consistency and possible deterioration, but they cannot establish the complete chemical quality of a preparation.

Physicochemical Evaluation
Physicochemical evaluation provides measurable information about the properties of the finished decoction.
Important parameters include:
a. pH: Measurement of hydrogen-ion activity provides information about the acidic or alkaline nature of the preparation. Changes during storage may indicate chemical or microbial alterations.
b. Total solids: The non-volatile material remaining after removal of water provides an estimate of extract concentration.
c. Specific gravity: This parameter may help assess consistency between batches.
d. Viscosity: Viscosity may provide information about the physical characteristics of preparations containing appreciable quantities of dissolved or dispersed constituents.
e. Ash values: Total ash, acid-insoluble ash and water-soluble ash are primarily used for evaluating crude medicinal materials and applicable powdered formulations. Their interpretation in finished liquid decoctions requires a separately defined and validated procedure.
f. Extractive values: Water-soluble and alcohol-soluble extractive values are relevant to the evaluation of applicable raw medicinal materials. For finished aqueous extracts, total solids and formulation-specific chemical assays may be more directly informative.
WHO describes established procedures for determining ash values and other quality-control parameters of medicinal plant materials. [7]

Phytochemical Screening
Preliminary phytochemical screening may help identify broad classes of constituents present in an aqueous extract.
Depending on the ingredients, evaluation may include:

- Alkaloids.
- Flavonoids.
- Phenolic compounds.
- Tannins.
- Saponins.
- Glycosides.
- Carbohydrates.
- Proteins and amino acids.

These tests provide preliminary qualitative information. They should not be interpreted as definitive identification or quantification of individual chemical compounds.

Chromatographic Fingerprinting
Chromatographic techniques provide a more specific approach to evaluating complex herbal preparations.
Thin-layer chromatography (TLC) and high-performance thin-layer chromatography (HPTLC) can be used to obtain characteristic chemical fingerprints.
High-performance liquid chromatography (HPLC) may be employed for separation and quantitative estimation of selected chemical markers.
Gas chromatography and gas chromatography–mass spectrometry may be useful when volatile or suitably derivatives constituents are relevant to the formulation.
The choice of analytical method should depend on the chemical characteristics of the ingredients and the intended quality-control objective.
The ConPhyMP recommendations emphasize the importance of documenting extraction conditions and using appropriate chemical characterization methods for plant extracts intended for pharmacological and toxicological research. [8]

Microbiological Evaluation

Aqueous decoctions contain substantial moisture and may provide favourable conditions for microbial growth.

Microbial quality is therefore an important consideration, particularly for preparations intended for storage or commercial distribution.

The evaluation may include:

- Total aerobic microbial count.
- Total yeast and mould count.
- Testing for specified objectionable microorganisms.
- Assessment of microbial deterioration during storage.

Applicable pharmacopoeial limits should be selected according to the formulation category and intended route of administration.

Heavy Metals and Other Contaminants

The quality evaluation of herbal medicinal substances should also consider potentially hazardous contaminants.

These may include lead, arsenic, cadmium and mercury, pesticide residues, aflatoxins and microbial contaminants.

The source material, manufacturing environment and water used for extraction may contribute to contamination.

Applicable limits and analytical procedures should be selected according to relevant pharmacopoeial and regulatory requirements. [6,7]

VII. MODERN EXTRACTION TECHNOLOGIES AND THEIR RELEVANCE TO KWATHA KALPANA

Traditional boiling remains the fundamental extraction method for Kwatha. However, contemporary pharmaceutical research has introduced several techniques that may improve extraction efficiency, process control and analytical reproducibility.

These techniques should be evaluated as potential research adaptations rather than assumed substitutes for classical procedures. [4,9]

Conventional Hot-Water Extraction

Conventional decoction involves heating medicinal substances in water for a prescribed duration.

Advantages include:

- Simple equipment requirements.
- Use of water as an extraction medium.
- Compatibility with traditional preparation procedures.
- Suitability for extracting several polar and water-soluble constituents.

Limitations include variable extraction efficiency, prolonged processing and possible degradation of heat-sensitive compounds.

Microwave-Assisted Extraction

Microwave-assisted extraction uses electromagnetic energy to facilitate heating within the extraction medium and plant material.

It may reduce extraction time and improve extraction efficiency under appropriately controlled conditions.

However, temperature distribution, microwave power and exposure duration must be optimized to minimize degradation of sensitive phytochemicals.

Ultrasound-Assisted Extraction

Ultrasound-assisted extraction involves acoustic cavitation, which may facilitate solvent penetration and disruption of plant tissues.

It can improve mass transfer and reduce the time required for extraction.

Its applicability to individual Kwatha formulations requires experimental validation.

Pressurized Hot-Water Extraction

Pressurized hot-water extraction employs water at elevated temperatures under controlled pressure.

Changes in the physical properties of water at higher temperatures may alter its solvent characteristics and extraction behavior.

However, the chemical composition of the resulting extract may differ from that of traditionally prepared Kwatha.

Comparative Evaluation

A modern extraction technique should be considered advantageous only when its performance is demonstrated through appropriate comparative experiments.

Important evaluation criteria include extraction yield, marker-compound concentration, chemical

fingerprint, processing time, energy consumption, degradation products and batch reproducibility.

Scientific literature indicates that the suitability of conventional and non-conventional extraction methods depends on the plant matrix, chemical characteristics of the target constituents and processing conditions. [4,9]

Table 4. Comparison of conventional and contemporary extraction methods

Method	Principal mechanism	Potential advantage	Important limitation
Conventional decoction	Thermal aqueous extraction	Traditional compatibility	Longer processing
Microwave-assisted extraction	Dielectric heating	Faster extraction	Thermal degradation risk
Ultrasound-assisted extraction	Acoustic cavitation	Improved mass transfer	Requires optimization
Pressurised hot-water extraction	Hot water under pressure	Modified solvent properties	Specialized equipment
Reflux extraction	Heating with solvent condensation	Reduced solvent loss	Prolonged thermal exposure

VIII. STABILITY AND SHELF-LIFE CONSIDERATIONS

One important pharmaceutical limitation of traditional aqueous decoctions is their potential susceptibility to deterioration during storage.

High water content and environmental exposure may promote microbial proliferation and chemical changes.

Freshly prepared Kwatha and commercially manufactured, processed or preserved aqueous decoctions should not be considered equivalent in terms of stability.

Factors Affecting Stability

The stability of Kwatha may be influenced by:

1. Initial microbial quality of raw materials.
2. Quality of water used for extraction.
3. Extraction temperature and duration.
4. pH of the finished preparation.
5. Concentration of dissolved solids.
6. Exposure to light and oxygen.
7. Storage temperature.
8. Nature and suitability of packaging.
9. Presence or absence of preservatives.

Chemical Stability

Certain phytochemicals may undergo oxidation, hydrolysis, polymerization or other chemical transformations during storage.

These changes may influence the chemical fingerprint and marker-compound concentration.

Stability studies should therefore include relevant physicochemical and chromatographic parameters.

Microbial Stability

Microbial contamination represents a major concern for aqueous preparations.

Appropriate microbial testing should be conducted at predetermined intervals under the proposed storage conditions.

The results should be used to establish a scientifically justified shelf life rather than assigning a storage duration solely on the basis of traditional assumptions.

Packaging and Preservation

Packaging should protect the preparation from environmental contamination, light and excessive exposure to oxygen.

Preservatives, where required, should be selected after evaluating their compatibility, safety and effectiveness in the particular formulation.

WHO guidelines on good manufacturing practices for herbal medicines provide a relevant framework for quality assurance during manufacturing and storage. [10]

IX. CONTEMPORARY RESEARCH PERSPECTIVES AND SCIENTIFIC VALIDATION

The scientific evaluation of Kwatha Kalpana requires a multidisciplinary approach involving pharmaceutical sciences, pharmacognosy, phytochemistry, analytical chemistry and pharmacology.

Traditional descriptions provide a foundation for preparation, but scientific validation requires reproducible experimental evidence.

Extraction Efficiency and Chemical Characterisation

One important research area is the relationship between traditional extraction conditions and the quantity of phytochemicals obtained.

Studies should evaluate the influence of particle size, water proportion, heating duration and temperature on extraction yield and chemical composition.

Comparative investigations can determine whether traditional preparation conditions provide reproducible extraction of selected marker compounds.

Chemical fingerprinting may further help identify batch-to-batch variations and possible degradation during processing. [8,9]

Pharmacological Evaluation

The pharmacological activity of a Kwatha formulation depends on its constituent medicinal substances, chemical composition, dose and biological availability.

Therefore, experimental evaluation should not be restricted to preliminary phytochemical screening.

Relevant investigations may include:

- In vitro antioxidant studies.
- Anti-inflammatory assays.
- Antimicrobial evaluation.
- Cytotoxicity and cell-based assays.
- Appropriate animal studies.
- Pharmacokinetic investigations.
- Toxicological evaluation.

The selection of experimental models should be based on the traditional therapeutic indications and scientifically justified research hypotheses.

An observed antioxidant or antimicrobial effect in vitro should not automatically be interpreted as evidence of clinical efficacy.

Standardisation of Polyherbal Decoctions

Polyherbal Kwatha preparations present additional standardisation challenges because several medicinal ingredients contribute to the final chemical composition.

The presence of multiple plant species may produce complex chemical interactions during aqueous extraction.

A suitable standardisation strategy should include authentication of every ingredient, controlled processing conditions, selected chemical markers and an overall chromatographic fingerprint.

The ConPhyMP recommendations provide a useful framework for documenting extraction parameters and characterising complex plant extracts intended for experimental research [8].

10.4 Pharmacological Reproducibility

A major requirement for translational herbal research is reproducibility.

Investigators should document:

- Botanical identity and voucher specimen details.
- Plant part and geographical origin.
- Collection season.
- Drying and storage conditions.
- Particle-size distribution.
- Drug-to-water ratio.
- Extraction temperature and duration.
- Final extract volume and yield.
- Chemical fingerprint.
- Experimental dose and route of administration.

Without these details, comparisons between different studies may be unreliable.

Integration of Traditional Knowledge and Modern Technology

The scientific development of Kwatha should maintain a balance between classical pharmaceutical principles and contemporary quality requirements.

Modern analytical methods can provide objective measurements of the chemical characteristics of traditional preparations.

However, modifications in extraction conditions should be justified through comparative studies to establish whether they preserve the relevant characteristics of the classical formulation.

The objective should be to improve consistency and safety without making unsupported assumptions about equivalence between different extraction technologies.

X. CRITICAL APPRAISAL: TRADITIONAL KNOWLEDGE VERSUS MODERN PHARMACEUTICAL PERSPECTIVES

Kwatha Kalpana represents a well-established pharmaceutical procedure in classical Ayurveda. Its continued relevance can be understood through its relatively simple preparation, use of water as an

extraction medium and adaptability to different medicinal ingredients.

Nevertheless, traditional preparation procedures and contemporary industrial requirements are not identical.

Classical descriptions provide general pharmaceutical principles, whereas modern manufacturing requires defined specifications, validated processes and measurable acceptance criteria.

A critical comparison identifies several important areas.

Table 5. Comparative appraisal of classical and contemporary approaches

Aspect	Classical approach	Contemporary pharmaceutical approach
Raw material	Traditional identification	Botanical and pharmacognostic authentication
Particle size	Yavakuta	Standardized sieve analysis
Extraction medium	Water	Qualified pharmaceutical water
Heating	Prescribed reduction procedure	Controlled temperature and time
Endpoint	Prescribed residual volume	Defined volume and analytical endpoint
Filtration	Traditional filtration	Defined filtration procedure
Quality assessment	Sensory and preparation-based evaluation	Physicochemical and chemical analysis
Stability	Traditional preparation and use practices	Validated stability studies
Quality assurance	Classical pharmaceutical principles	GMP and documented quality systems

An important consideration is that standardization should not be restricted to obtaining identical physical characteristics. Two decoctions may have similar colour, pH and total solids but differ in their chemical composition.

Consequently, the integration of physicochemical evaluation with chromatographic fingerprinting and marker-compound estimation is particularly relevant.

Another challenge concerns the interpretation of pharmacological findings. Biological activity demonstrated by an isolated extract or a laboratory model cannot automatically be extrapolated to the traditionally prepared formulation or its clinical application.

Thus, the scientific evaluation of Kwatha requires a sequential approach involving raw material authentication, pharmaceutical standardization, chemical characterization, safety assessment and appropriately designed biological investigations.

XI. CONCLUSION

Kwatha Kalpana is an important classical pharmaceutical preparation that demonstrates the systematic use of aqueous extraction in Ayurveda. Its principles involve the appropriate selection of medicinal substances, size reduction, controlled addition of water, thermal processing, volume reduction and filtration.

Although the traditional method provides a practical foundation for decoction preparation, variations in raw material quality and processing conditions may influence the chemical composition and reproducibility of the finished product.

Contemporary pharmaceutical science offers several opportunities to strengthen the standardization of Kwatha through physicochemical evaluation, phytochemical screening, chromatographic fingerprinting, microbial testing and stability assessment.

Modern extraction technologies may also provide useful approaches for improving process efficiency, provided their suitability is established through formulation-specific comparative investigations.

The integration of classical pharmaceutical knowledge with validated modern analytical procedures represents a rational direction for the future development of Kwatha Kalpana. Such an approach may improve quality assurance, manufacturing consistency and scientific understanding while preserving the fundamental principles of traditional pharmaceutical preparation.

XII. LIMITATIONS OF THE REVIEW

The present article is a narrative review and does not represent a systematic review or meta-analysis. The

available literature includes classical textual descriptions, pharmacopoeial standards, experimental investigations and modern pharmaceutical publications with varying research designs. Therefore, the findings should be interpreted as a qualitative synthesis. Further systematic and experimental studies are necessary to establish formulation-specific extraction standards and their relationship with pharmacological and clinical outcomes.

REFERENCES

[1] Shastri K. and G. Chaturvedi, eds., *Charaka Samhita of Agnivesha with Vidyotini Hindi Commentary by Kashinath Shastri and Gorakhnath Chaturvedi*, Sutra Sthana, ch. 4, ver. 7. Varanasi, India: Chaukhambha Bharati Academy, 1984.

[2] K. R. S. Murthy, trans., *Sharngadhara Samhita of Sharngadhara*, Madhyama Khanda, ch. 2, verses 1–2. Varanasi, India: Chaukhambha Orientalia, 1984.

[3] N. S. Bhatt and M. Deshpande, "A critical review and significance of Ayurvedic preparation Kwatha—herbal decoction," *International Journal of Ayurvedic Medicine*, vol. 11, no. 2, pp. 155–164, 2020.

[4] J. Azmir, I. S. M. Zaidul, M. M. Rahman, K. M. Sharif, A. Mohamed, F. Sahena, *et al*., "Techniques for extraction of bioactive compounds from plant materials: A review," *Journal of Food Engineering*, vol. 117, no. 4, pp. 426–436, 2013, doi: 10.1016/j.jfoodeng.2013.01.014.

[5] H. Sharma, ed., *Kashyapa Samhita of Vriddha Jivaka with Sanskrit Introduction and Hindi Commentary*, Khila Sthana, ch. 3. Varanasi, India: Chaukhambha Sanskrit Series Office, 1994.

[6] Government of India, Ministry of AYUSH, Pharmacopoeia Commission for Indian Medicine and Homoeopathy, *Pharmacopoeial Monograph of AYUSH Kvatha Curna*, *The Ayurvedic Pharmacopoeia of India*, Part II (Formulations). New Delhi, India: Government of India, 2020, UINADS: API-2/2020/KC/1.0.

[7] World Health Organization, *Quality Control Methods for Medicinal Plant Materials*. Geneva, Switzerland: World Health Organization, 1998, ISBN: 92-4-154510-0.

[8] M. Heinrich, B. Jalil, M. Abdel-Tawab, J. Echeverria, Ž. Kulić, L. J. McGaw, *et al*., "Best practice in the chemical characterization of extracts used in pharmacological and toxicological research—the ConPhyMP guidelines," *Frontiers in Pharmacology*, vol. 13, Art. no. 953205, 2022, doi: 10.3389/fphar.2022.953205.

[9] R. Abubakar and M. Haque, "Preparation of medicinal plants: Basic extraction and fractionation procedures for experimental purposes," *Journal of Pharmacy and Bioallied Sciences*, vol. 12, no. 1, pp. 1–10, 2020, doi: 10.4103/jpbs.JPBS_175_19.

[10] World Health Organization, *WHO Guidelines on Good Manufacturing Practices (GMP) for Herbal Medicines*. Geneva, Switzerland: World Health Organization, 2007.