No Salt to Form: Making Piperine More Soluble

Shruti Singh[1], Swarup J. Chatterjee[2]
[1] *Department of Pharmaceutical Chemistry, S. N. College of Pharmacy, Babupur, Lucknow Road, Jaunpur – 222132, Uttar Pradesh, India. Affiliated to Dr. A. P. J. Abdul Kalam Technical University, Lucknow.*
[2] *Professor & Principal, Department of Pharmaceutical Chemistry, S. N. College of Pharmacy, Babupur, Lucknow Road, Jaunpur – 222132, Uttar Pradesh, India.*

***Abstract*—Piperine dissolves in water to about 40 mg per litre, and almost every proposal to fix that begins with the same instinct: make a salt. For this molecule that instinct is wrong, and the reason is visible in the structure. The only nitrogen in piperine is an amide nitrogen, its lone pair delocalised into the carbonyl; the predicted strongest basic pK_a is below zero, the molecule carries no charge at any physiological pH, and it has no hydrogen-bond donor at all. There is no salt to form. This review takes that constraint as its starting point and asks what remains. The general solubility equation — log S = 0.5 − 0.01(mp − 25) − log P, which applies precisely to non-ionisable solutes such as this one — reproduces piperine's measured solubility to within a factor of four and turns the project's structure–solubility question into arithmetic: lowering log P by one unit is worth ten-fold, lowering the melting point by 50 °C is worth about three-fold, and doing both is worth thirty. Three semisynthetic handles are then assessed against that arithmetic. Alkaline hydrolysis to piperic acid installs an ionisable group but raises the melting point by roughly 80 °C; re-amidation with a basic amine such as N-methylpiperazine restores salt formation; saturating the diene lowers the lattice energy but destroys the chromophore used to assay the compound. The methylenedioxyphenyl ring is the one region that should not be touched, because it is the pharmacophore for the cytochrome P450 and P-glycoprotein inhibition that makes piperine interesting in the first place.**

***Index Terms*—Piperine; Piper nigrum; aqueous solubility; general solubility equation; piperic acid; semisynthesis; methylenedioxyphenyl**

I. INTRODUCTION

Piperine is the pungent principle of black pepper, isolated by Ørsted in 1819 and now one of the most heavily studied plant amides in pharmacology [1–3]. Its interest is not mainly as a drug in its own right. It is a bioenhancer: by inhibiting cytochrome P450 3A4 and P-glycoprotein it raises the systemic exposure of co-administered agents, famously increasing the bioavailability of curcumin by a factor of twenty in human volunteers [27–29]. The obstacle to using it, and to studying it, is that it barely dissolves: roughly 40 mg per litre of water at 18 °C.

A project that proposes to fix this by rational semisynthesis is well framed, but it inherits one constraint that has to be stated before any chemistry is planned. The standard first move in solubility enhancement is salt formation, which is used for more than half of all marketed small-molecule drugs and typically buys one to three orders of magnitude [13–15]. Piperine cannot be salted, and the reason is structural rather than practical.

It is worth being precise about why 40 mg L^{-1} is a problem, because piperine is not a compound that struggles to cross membranes. With a log P near 3 and a polar surface area under 40 $Å^2$ it satisfies every criterion of the rule of five and is highly permeable [33]. It is a textbook class II compound in the Biopharmaceutics Classification System: absorption is limited by how much dissolves, not by how much crosses [34]. That classification is good news for a solubility project, because it means the solubility term is the whole of the problem and improving it should translate directly into exposure — a claim that cannot be made for class IV compounds, where both terms fail [35,36].

This review is organised around that fact. Section 2 establishes the physicochemical profile and the ionisation argument. Section 3 deals with isolation, including a stability problem that affects every

measurement made afterwards. Section 4 introduces the quantitative tool that converts a structure–solubility question into arithmetic. Section 5 assesses the three semisynthetic handles the molecule actually offers against that arithmetic, and says what each one costs. Section 6 deals with measuring solubility in a way that will survive scrutiny.

II. THE MOLECULE, MEASURED

2.1 The numbers

Piperine is (2E,4E)-5-(benzo[d][1,3]dioxol-5-yl)-1-(piperidin-1-yl)penta-2,4-dien-1-one, $C_{17}H_{19}NO_3$, molar mass 285.34. It is a pale yellow crystalline solid melting at 128–135 °C depending on the source and purity, with a reported aqueous solubility of 40 mg L^{-1} at 18 °C and estimated log P between 2.7 and 3.4 [1]. It dissolves freely in ethanol, chloroform and dimethyl sulfoxide at around 10 mg mL^{-1}, which is why almost every biological study of piperine is a study of a DMSO stock solution. Table 1 collects the properties that matter for the present purpose and what each one implies.

Two of those entries do most of the work. The polar surface area is only 38.8 Å² and the hydrogen-bond donor count is zero: the amide nitrogen is fully substituted, the two dioxole oxygens and the carbonyl are acceptors only. A molecule with no donor cannot form the hydrogen bonds to water that would offset the energy needed to break its crystal lattice. The other entry is the melting point, which at 130 °C is high for a molecule of this size and reflects efficient packing of a flat, conjugated, rod-shaped molecule.

The methylenedioxy bridge is worth one note of origin, since it recurs throughout this review. It is installed late in the plant by a dedicated cytochrome P450, CYP719A37, acting on a methoxy-hydroxy precursor [7]. The same five-membered acetal that the plant works to build is what makes the compound flat, lipophilic and, as section 5.4 argues, pharmacologically interesting.




Fig. 1. Piperine and three semisynthetic directions. (a) The parent: the only nitrogen is an amide nitrogen, so the molecule is neutral at every physiological pH and has no hydrogen-bond donor. (b) Hydrolysis gives an ionisable acid but a much higher melting point. (c) Re-amidation with a basic amine restores salt formation. (d) Saturating the diene lowers the lattice energy and removes the chromophore.

Table 1. Physicochemical profile of piperine. The last three rows are the ones that decide the project's strategy [1,9].

Property	Value	Basis	What it implies for solubility
Molecular formula, mass	$C_{17}H_{19}NO_3$, 285.34	Structure	Small enough that solubility is not a size problem

Property	Value	Basis	What it implies for solubility
Melting point	128–135 °C	Measured, several sources	High for the mass; the lattice term costs about 1.05 log units
Aqueous solubility	40 mg L^{-1} (18 °C)	Measured	log S = −3.85; practically insoluble
log P	2.66–3.37	Estimated	The dominant term in the solubility equation
Hydrogen-bond donors	0	Structure	No donor to hydrate the solute; cannot be improved without adding one
Polar surface area	38.8 $Å^2$	Computed	Low; consistent with high membrane permeability and poor solubility
Strongest basic pK_a	−0.1 to −0.7 (computed)	Amide nitrogen	No protonation at usable pH; salt formation unavailable
Physiological charge	0	Computed	Non-ionisable from pH 2 to 13, so the GSE applies without correction

2.2 Why there is no salt to form

Piperine looks basic. It contains a piperidine ring, and piperidine itself is a strong base that is miscible with water. But in piperine that ring is acylated: the nitrogen is an amide nitrogen, and its lone pair is delocalised into the carbonyl π system. Amide nitrogens are not protonated at accessible pH; the conjugate acid of a tertiary amide has a pK_a around −0.5, and computed values for piperine fall between −0.1 and −0.7, with a physiological charge of zero [1]. Protonation under those conditions would require concentrated strong acid and would occur at oxygen, not nitrogen.

This matters because catalogue entries and secondary sources circulate a pK_a of about 12 for piperine, and some reviews describe it as a weak base forming acid salts. A value near 12 cannot describe protonation of an N-acyl piperidine, and a student who plans a hydrochloride salt on that basis will produce either unchanged starting material or hydrolysis products. The chemically defensible statement is the one in Fig. 1a: piperine is a neutral, non-ionisable amide across the whole pharmaceutically relevant pH range, which is exactly the class of compound to which the general solubility equation of Section 4 applies [9].

Ruling salt formation out is not a negative result. It narrows the design space to three honest options — reduce log P, reduce the melting point, or install an ionisable group that the parent does not have — and section 5 shows that the semisynthesis can deliver all three.

III. ISOLATION, AND WHAT IT DOES TO THE MATERIAL

3.1 Getting the compound

Black pepper contains roughly 2–9% piperine by weight, and the classical teaching preparation extracts the ground fruit with ethanol or dichloromethane, concentrates the oleoresin, triturates it with alcoholic potassium hydroxide to remove acidic and resinous material, and allows the piperine to crystallise from the decanted filtrate as yellow needles [4,5]. The procedure is reliable and is the right starting point.


Two cautions belong in the experimental section. First, the alkaline treatment that removes the resin is the same chemistry that hydrolyses piperine (Scheme 1); it is a matter of concentration and time, so the potassium hydroxide step should be brief and cold, and the mother liquor should be checked for piperic acid rather than discarded. Second, commercial pepper varies, and a yield expressed as a percentage of starting material is a statement about the pepper as much as about the method; purity should be established by melting point and chromatography rather than inferred from appearance.

3.2 The isomers that form in daylight

Piperine is photolabile. The 2E,4E dienamide interconverts with its 2Z,4E, 2E,4Z and 2Z,4Z geometric isomers — isopiperine, isochavicine and chavicine — under ordinary laboratory lighting and much faster under ultraviolet light (Scheme 3). The kinetics have been measured by HPLC and LC-MS,

and all four isomers are detectable in ground black pepper itself [6].

For a solubility project this is not a footnote. A sample that has stood in a clear flask on a bench is a mixture, and a mixture of geometric isomers melts lower and dissolves better than either pure component — so the apparent solubility improvement from almost any handling step can be an isomerisation artefact. The controls are inexpensive: amber glassware or foil throughout, a melting range rather than a single figure, and an HPLC trace of the actual sample used for each solubility determination, reported alongside the result. A sharp melting point near 130 °C and a single chromatographic peak are the evidence that the number refers to piperine.



piperine (2E,4E) chavicine (2Z,4Z)

daylight and UV interconvert the four geometric isomers; the mixture melts lower and dissolves better, but it is a mixture

Scheme 3. Photoisomerisation of the dienamide. All four geometric isomers are present in ground pepper and interconvert in ordinary laboratory light, so a solubility measured on an uncharacterised sample may be the solubility of a mixture.

IV. SOLUBILITY IS A PROPERTY OF THE CRYSTAL AS MUCH AS THE MOLECULE

4.1 The general solubility equation

The relationship the project's title calls a structure–solubility relationship already exists in quantitative form. Yalkowsky and Valvani showed that the aqueous solubility of a non-electrolyte is governed by two terms, one for how hard it is to melt the crystal and one for how unwilling the resulting liquid is to mix with water [8]. In the revised form given by Jain and Yalkowsky,

$$\log S = 0.5 - 0.01(mp - 25) - \log P,$$

where S is the intrinsic molar solubility, mp is the Celsius melting point and log P the octanol–water partition coefficient [9]. The equation has no fitted parameters, has been validated on more than a thousand compounds, and carries a root-mean-square error of roughly 0.7 to 0.8 log units [10,11]. Critically for this project, it is derived for compounds that are non-ionisable between pH 2 and 13 — which is precisely what Section 2.2 established piperine to be.

Applied to piperine with mp 130 °C and log P 2.66, the equation gives log S = −3.21, or 176 mg L^{-1}. With the higher reported log P of 3.37 it gives 34 mg L^{-1}. The measured value, 40 mg L^{-1}, sits between them and within the equation's stated error. That agreement is the licence to use the equation predictively (Fig. 2a).

Two caveats keep the tool honest. The equation predicts intrinsic solubility of the stable crystalline form, so it will under-predict for an amorphous or metastable sample and cannot describe a salt. And its accuracy depends on the inputs: a calculated log P introduces its own error, which is why the two literature values for piperine bracket the measurement rather than agreeing with it. More elaborate computational methods exist and perform somewhat better on diverse sets, but none is as transparent, and for a design decision transparency is worth more than a tenth of a log unit [37,41].




Fig. 2. (a) The general solubility equation plotted for four values of log P, with piperine's predicted and measured solubility marked; the two differ by less than the equation's stated error. (b) Predicted log S for piperine and for three hypothetical modifications, computed from the same equation; figures in mg L^{-1} above each bar, measured piperine solubility shown as the dotted line.

4.2 What each change is worth

Because the two terms are additive and their coefficients are fixed, the value of any proposed modification can be read off directly, and Fig. 2b does so. Removing one log unit of lipophilicity multiplies solubility by ten. Lowering the melting point by 50 °C multiplies it by about three. Doing both multiplies it by thirty, taking a predicted 176 mg L^{-1} to roughly 5.6 g L^{-1}. Table 2 applies this to the specific modifications available here.

Two lessons follow that are not obvious from inspection of the structure. The first is that polarity is worth more per unit than crystal disruption, so adding a polar group usually beats branching a chain. The second, and the more useful one, is that the two can oppose each other. Piperic acid is far more polar than piperine, but it melts around 206–217 °C rather than 130 °C, because the carboxylic acid forms hydrogen-bonded dimers in the lattice. Its intrinsic solubility as the neutral acid is predicted at only about 108 mg L^{-1} — the melting point term has eaten most of the polarity gain. The acid wins anyway, but it wins by ionising, not by being more polar.

Two further routes change the solid without changing the molecule at all, and should be named even though they fall outside a synthetic project: cocrystallisation with a coformer, and preparation of the amorphous form, both of which raise apparent solubility by removing or weakening the lattice term [16–18]. Cyclodextrin complexation does something different again, solubilising rather than dissolving, and has been applied to piperine with measurable success [19,20,38]. Particle-size reduction to a nanosuspension raises dissolution rate and apparent solubility without touching the molecule either, and has been shown to increase piperine's oral bioavailability [39]. None of these is a competitor to the synthetic work; they are the comparators against which it should be judged, because a chemist who spends six months making an analogue that performs no better than milling the parent has learned something important and should report it.

There is also a reason to care about the melting point beyond the solubility equation. Across large data sets, melting point correlates with oral absorption independently of solubility, with high-melting compounds absorbed less well than their calculated solubility would suggest [12]. A modification that lowers the melting point therefore tends to help twice.

Table 2. Semisynthetic options assessed against the general solubility equation. Changes in log P are estimates from fragment contributions and should be recalculated for each actual analogue [9–11].


Modification	Δ log P	Δ mp	Predicted effect	What it costs
None (piperine)	—	—	log S −3.21 (176 mg L^{-1})	—
Hydrolysis to piperic acid	≈ −0.6	+ 80 °C	Neutral form barely improved; the salt is freely soluble	Loses the amide and most of the activity
Morpholine amide	≈ −1.0	small	About ten-fold on the neutral form	Minimal; closest analogue of the parent
N-Methylpiperazine amide	≈ −1.0	small	Ten-fold, plus salt formation at the distal nitrogen	Adds a basic centre that may alter distribution
Amino-alcohol amide	≈ −1.5	variable	Thirty-fold or more; adds a hydrogen-bond donor	Possible loss of planarity-dependent binding
Saturation of the diene	≈ 0	− 30 to − 50 °C	Two- to three-fold	Chromophore and part of the pharmacophore
Opening the methylenedioxy bridge	≈ −0.5	variable	Several-fold	The CYP3A4 and P-gp inhibition; not recommended

V. THE THREE HANDLES THE MOLECULE OFFERS

5.1 Hydrolysis to piperic acid

Refluxing piperine with 20% ethanolic potassium hydroxide for 16–22 hours and acidifying to pH below 1 gives piperic acid and piperidine (Scheme 1); reported yields range from 18% to 72% depending on conditions [23,24]. Piperic acid is the key intermediate for everything that follows, and it is also a modification in its own right. Its carboxylic acid has a pK_a near 4.5, so at intestinal pH it is largely ionised and its sodium or potassium salt is freely water soluble. This is the salt the project can legitimately make — just not from the nitrogen.

The cost is the amide. Removing the piperidine ring removes most of what distinguishes piperine from a cinnamic acid derivative, and the reported biological activity of piperic acid is generally much weaker than that of the parent [23]. Piperic acid is best regarded as an intermediate and a reference point rather than as a destination.



piperine
mp 130 °C, log P 2.7, no ionisable group

piperic acid
mp 206–217 °C, pK_a ≈ 4.5


piperidine

Scheme 1. Alkaline hydrolysis of piperine to piperic acid and piperidine. This is both the route to every analogue in Scheme 2 and the side reaction that the alkaline wash during isolation can cause.

5.2 Re-amidation: the handle that restores the salt

Converting piperic acid to its acid chloride with thionyl or oxalyl chloride, or activating it with a coupling reagent, and treating it with an amine gives piperamide analogues in good yield (Scheme 2) [23,25,26]. This is the most productive handle the scaffold has, because the amine is a free variable while the dienamide and the methylenedioxyphenyl ring are untouched.

For a solubility project the choice of amine is the whole experiment, and it can be made with the arithmetic of Section 4 in advance. A morpholine amide lowers log P by roughly one unit relative to the piperidine amide through the ring oxygen. An N-methylpiperazine amide (Fig. 1c) does the same and adds something the parent does not have: the distal nitrogen is a genuine amine, not an amide, with a pK_a near 7, so it protonates in the stomach and forms isolable hydrochloride, citrate and mesylate salts. An amino acid or aminoethanol amide introduces a hydrogen-bond donor, which both lowers log P and tends to disrupt the planar packing. Each of these is a single-step modification of a common intermediate, which makes the series an unusually clean structure–solubility experiment: one variable, several points, and a predicted value for each before it is measured.



Scheme 2. Activation of piperic acid and re-amidation. The amine is the only variable, which makes a series prepared this way a clean structure–solubility experiment.

5.3 Saturating the diene

Hydrogenation of the 2,4-diene gives tetrahydropiperine (Fig. 1d), a compound already used commercially as a skin-penetration enhancer. The conformationally flexible chain packs less efficiently than the rigid conjugated one, so the melting point falls and, by the arithmetic above, solubility rises; log P changes little. Two costs attach. The conjugated chromophore disappears, so the ultraviolet method used to assay piperine at around 343 nm no longer applies and the analytical method must be rebuilt. And the diene is part of the pharmacophore for several of piperine's reported activities, so the analogue is a different compound pharmacologically, not a more soluble piperine [26].

5.4 The region to leave alone

One part of the molecule should be excluded from the design space on principle. The methylenedioxyphenyl group is the structural basis of

piperine's inhibition of cytochrome P450 3A4 and P-glycoprotein [27,28]. Methylenedioxyphenyl compounds are oxidised by P450 to a carbene that coordinates the haem iron, forming a stable metabolic intermediate complex and inactivating the enzyme; this mechanism is shared with the insecticide synergists of the same class and is well characterised [30–32]. It is the reason piperine raises curcumin exposure twenty-fold [29].

A series that opens the methylenedioxy bridge to a catechol or a dimethoxy group will almost certainly be more soluble, because both modifications add donors or reduce planarity. It will also almost certainly have lost the bioenhancer activity, which was the point. The general principle is worth stating plainly in the thesis: in a solubility-optimisation project on a natural product, the pharmacophore has to be identified first and placed out of bounds, or the project will succeed on its stated endpoint and fail on its purpose.

VI. MEASURING THE SOLUBILITY PROPERLY

The number that the whole project turns on is the easiest thing in it to get wrong, and Table 3 lists the common failures. Four points are worth making directly.

Equilibrium solubility is measured by the saturation shake-flask method: excess solid in buffer, agitated at controlled temperature for 24 to 72 hours, separated by filtration or centrifugation, and assayed against a calibration curve, with the residual solid examined afterwards to confirm it has not changed form [21,22]. Apparent solubility obtained by diluting a DMSO stock is a different quantity, usually higher, and the two must not be compared.

The medium must be stated. Since piperine is non-ionisable, its solubility should be the same in pH 1.2 and pH 6.8 buffers; measuring both and finding them equal is a useful positive control, and finding them different indicates ionisation, degradation or an artefact. For the piperamide analogues that are ionisable, solubility becomes pH-dependent and both media must be reported.

The assay must be specific. Because piperine photoisomerises and the isomers absorb at similar wavelengths, a single-wavelength ultraviolet reading cannot distinguish them; a validated HPLC method with a stated retention time and purity check should be used, following the usual validation parameters [40].

Finally, the comparison must be like for like. The useful statement is the fold increase in equilibrium solubility of a pure, characterised analogue over pure piperine measured in the same medium on the same day by the same method, with the melting point and log P of both reported so that the result can be checked against the prediction. A table of analogues with measured melting points, measured or calculated log P, predicted log S and measured log S is the deliverable that would make this project genuinely useful — and it is a table nobody has published for this scaffold.

Table 3. Common failures in solubility measurement and their remedies [21,22,40].

Practice	Why it misleads	What to do instead
Diluting a DMSO stock into buffer and reading absorbance	Measures kinetic, not equilibrium, solubility; cosolvent raises the result and the value depends on dilution order	Saturation shake-flask with excess solid, 24–72 h, controlled temperature
Reporting one number with no medium stated	Solubility in water, saline and buffer differ, and for ionisable analogues so does pH	State buffer, pH, ionic strength and temperature; for piperine, measure at pH 1.2 and 6.8 and expect them to agree
Single-wavelength UV assay	Geometric isomers and degradation products absorb in the same region	Validated HPLC with retention time and peak purity reported
No check on the residual solid	The excess solid may have converted to a hydrate or an amorphous form during equilibration	Recover and examine it by melting point, infrared or powder diffraction

Practice	Why it misleads	What to do instead
Comparing a new analogue with a literature value for piperine	Different method, medium, temperature and purity	Measure both compounds side by side, same day, same method
No light protection	Photoisomerisation lowers the melting point and raises apparent solubility	Amber glassware throughout; report the chromatogram of the actual sample

VII. CONCLUSION

Piperine is a good molecule for a solubility project precisely because it is simple. It is neutral across the whole physiological range, so the general solubility equation applies to it without correction; its measured solubility matches the prediction; and the equation then says, quantitatively and in advance, what any proposed modification is worth. The instinct to make a salt has to be abandoned, because the only nitrogen is an amide — but the semisynthesis supplies a better route to the same end, since hydrolysis and re-amidation put a genuinely basic nitrogen in the molecule for the first time.

What distinguishes a useful dissertation here from a routine one is discipline about two things. The isolated material must be shown to be the 2E,4E isomer at the moment the solubility is measured, because daylight alone will otherwise supply a spurious improvement. And the methylenedioxyphenyl ring must be left intact, because it carries the activity that makes the solubility worth improving. Within those two constraints the design space is small, well defined, and quantitatively predictable — which is an unusually comfortable position for a synthetic project to be in.

REFERENCES

[1] Gorgani L, Mohammadi M, Najafpour GD, Nikzad M. Piperine — the bioactive compound of black pepper: from isolation to medicinal formulations. Compr Rev Food Sci Food Saf. 2017;16(1):124–140.

[2] Srinivasan K. Black pepper and its pungent principle-piperine: a review of diverse physiological effects. Crit Rev Food Sci Nutr. 2007;47(8):735–748.

[3] Meghwal M, Goswami TK. Piper nigrum and piperine: an update. Phytother Res. 2013;27(8):1121–1130.

[4] Ikan R. Natural Products: A Laboratory Guide. 2nd ed. San Diego: Academic Press; 1991.

[5] Epstein WW, Netz DF, Seidel JL. Isolation of piperine from black pepper. J Chem Educ. 1993;70(7):598.

[6] Kozukue N, Park MS, Choi SH, Lee SU, Ohnishi-Kameyama M, Levin CE, Friedman M. Kinetics of light-induced cis-trans isomerization of four piperines and their levels in ground black peppers as determined by HPLC and LC/MS. J Agric Food Chem. 2007;55(17):7131–7139.

[7] Schnabel A, Cotinguiba F, Athmer B, Vogt T. Piper nigrum CYP719A37 catalyzes the decisive methylenedioxy bridge formation in piperine biosynthesis. Plants. 2021;10(1):128.

[8] Yalkowsky SH, Valvani SC. Solubility and partitioning I: solubility of nonelectrolytes in water. J Pharm Sci. 1980;69(8):912–922.

[9] Jain N, Yalkowsky SH. Estimation of the aqueous solubility I: application to organic nonelectrolytes. J Pharm Sci. 2001;90(2):234–252.

[10] Ran Y, Yalkowsky SH. Prediction of drug solubility by the general solubility equation (GSE). J Chem Inf Comput Sci. 2001;41(2):354–357.

[11] Ran Y, Jain N, Yalkowsky SH. Prediction of aqueous solubility of organic compounds by the general solubility equation (GSE). J Chem Inf Comput Sci. 2001;41(5):1208–1217.

[12] Chu KA, Yalkowsky SH. An interesting relationship between drug absorption and melting point. Int J Pharm. 2009;373(1–2):24–40.

[13] Serajuddin ATM. Salt formation to improve drug solubility. Adv Drug Deliv Rev. 2007;59(7):603–616.

[14] Stahl PH, Wermuth CG, editors. Handbook of Pharmaceutical Salts: Properties, Selection and Use. 2nd ed. Weinheim: Wiley-VCH; 2011.
[15] Bastin RJ, Bowker MJ, Slater BJ. Salt selection and optimisation procedures for pharmaceutical new chemical entities. Org Process Res Dev. 2000;4(5):427–435.
[16] Babu NJ, Nangia A. Solubility advantage of amorphous drugs and pharmaceutical cocrystals. Cryst Growth Des. 2011;11(7):2662–2679.
[17] Good DJ, Rodríguez-Hornedo N. Solubility advantage of pharmaceutical cocrystals. Cryst Growth Des. 2009;9(5):2252–2264.
[18] Hancock BC, Zografi G. Characteristics and significance of the amorphous state in pharmaceutical systems. J Pharm Sci. 1997;86(1):1–12.
[19] Loftsson T, Brewster ME. Pharmaceutical applications of cyclodextrins. 1. Drug solubilization and stabilization. J Pharm Sci. 1996;85(10):1017–1025.
[20] Higuchi T, Connors KA. Phase-solubility techniques. Adv Anal Chem Instrum. 1965;4:117–212.
[21] Baka E, Comer JEA, Takács-Novák K. Study of equilibrium solubility measurement by saturation shake-flask method using hydrochlorothiazide as model compound. J Pharm Biomed Anal. 2008;46(2):335–341.
[22] Avdeef A, Fuguet E, Llinàs A, Ràfols C, Bosch E, Völgyi G, et al. Equilibrium solubility measurement of ionizable drugs — consensus recommendations for improving data quality. ADMET DMPK. 2016;4(2):117–178.
[23] Wansri R, Lin ACK, Pengon J, Kamchonwongpaisan S, Srimongkolpithak N, Rattanajak R, et al. Semi-synthesis of N-aryl amide analogs of piperine from Piper nigrum and evaluation of their antitrypanosomal, antimalarial, and anti-SARS-CoV-2 main protease activities. Molecules. 2022;27(9):2841.
[24] Choochana P, Moungjaroen J, Jongkon N, Gritsanapan W, Tangyuenyongwatana P. Development of piperic acid derivatives from Piper nigrum as UV protection agents. Pharm Biol. 2015;53(4):477–482.
[25] Synthesis of piperine analogues as AChE and BChE inhibitors for the treatment of Alzheimer's disease. Phytochem Lett. 2023. [Author list, volume and page range to be completed.]
[26] Venkatasamy R, Faas L, Young AR, Raman A, Hider RC. Effects of piperine analogues on stimulation of melanocyte proliferation and melanocyte differentiation. Bioorg Med Chem. 2004;12(8):1905–1920.
[27] Bhardwaj RK, Glaeser H, Becquemont L, Klotz U, Gupta SK, Fromm MF. Piperine, a major constituent of black pepper, inhibits human P-glycoprotein and CYP3A4. J Pharmacol Exp Ther. 2002;302(2):645–650.
[28] Atal CK, Dubey RK, Singh J. Biochemical basis of enhanced drug bioavailability by piperine: evidence that piperine is a potent inhibitor of drug metabolism. J Pharmacol Exp Ther. 1985;232(1):258–262.
[29] Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas PS. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. Planta Med. 1998;64(4):353–356.
[30] Murray M. Mechanisms of inhibitory and regulatory effects of methylenedioxyphenyl compounds on cytochrome P450-dependent drug oxidation. Curr Drug Metab. 2000;1(1):67–84.
[31] Correia MA, Hollenberg PF. Inhibition of cytochrome P450 enzymes. In: Ortiz de Montellano PR, editor. Cytochrome P450: Structure, Mechanism, and Biochemistry. 4th ed. Cham: Springer; 2015. p. 177–259.
[32] Franklin MR. Methylenedioxyphenyl insecticide synergists as potential human health hazards. Environ Health Perspect. 1976;14:29–37.
[33] Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. Adv Drug Deliv Rev. 1997;23(1–3):3–25.
[34] Amidon GL, Lennernäs H, Shah VP, Crison JR. A theoretical basis for a biopharmaceutic drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability. Pharm Res. 1995;12(3):413–420.
[35] Williams HD, Trevaskis NL, Charman SA, Shanker RM, Charman WN, Pouton CW, Porter CJH. Strategies to address low drug solubility in discovery and development. Pharmacol Rev. 2013;65(1):315–499.

[36] Di L, Fish PV, Mano T. Bridging solubility between drug discovery and development. Drug Discov Today. 2012;17(9–10):486–495.
[37] Yalkowsky SH. Solubility and Solubilization in Aqueous Media. Washington, DC: American Chemical Society; 1999.
[38] Ezawa T, Inoue Y, Murata I, Takao K, Sugita Y, Kanamoto I. Characterization of the dissolution behavior of piperine/cyclodextrins inclusion complexes. AAPS PharmSciTech. 2018;19(2):923–933.
[39] Zafar F, Jahan N, Bhatti HN. Increased oral bioavailability of piperine from an optimized Piper nigrum nanosuspension. Planta Med. 2019;85(3):249–257.
[40] International Council for Harmonisation. Validation of Analytical Procedures Q2(R2). Geneva: ICH; 2023.
[41] Bergström CAS, Larsson P. Computational prediction of drug solubility in water-based systems: qualitative and quantitative approaches used in the current drug discovery paradigm. Int J Pharm. 2018;540(1–2):185–193.